

Investigation of the relationship between immune checkpoints and mismatch repair deficiency in recurrent and nonrecurrent glioblastoma

Emel ÇAKIR* , İsmail SAYGIN , Mustafa Emre ERCİN 

Department of Pathology, Faculty of Medicine, Karadeniz Technical University, Trabzon, Turkey

Received: 14.10.2020 • Accepted/Published Online: 18.02.2021 • Final Version: 30.08.2021

Background/aim: Microsatellite instability tests and programmed cell death-1 (PD-1)/programmed cell death ligand-1 (PD-L1) in the immune checkpoint pathway are the tests that determine who will benefit from immune checkpoint inhibitor therapy. We aimed to show the expression of DNA mismatch repair proteins and PD-1/PD-L1 molecules that inhibit immune checkpoints, to explain the relationship between them, and to demonstrate their predictive role in recurrent and nonrecurrent glioblastoma.

Materials and methods: We analyzed 27 recurrent and 47 nonrecurrent cases at our archive. We performed immunohistochemical analysis to determine expressions of PD-1, PD-L1, and mismatch repair proteins in glioblastoma. We evaluated the relationship between these two group and compared the results with the clinicopathological features.

Results: The mean age of diagnosis was significantly lower in recurrent glioblastoma patients. Median survival was longer in this group. We found that PD-L1 expression was reduced in recurrent cases. Additionally, recurrent cases had a significantly higher rate of microsatellite instability. Loss of PMS2 was high in both group but was substantially higher in recurrent cases.

Conclusion: The presence of microsatellite instability and low PD-L1 levels, which are among the causes of treatment resistance in glioblastoma, were found to be compatible with the literature in our study, with higher rates in recurrent cases. In recurrent cases with higher mutations and where immunotherapy resistance is expected less, low PD-L1 levels thought that different combinations with other immune checkpoint inhibitors can be tried as predictive and prognostic marker in GBM patients.

Key words: Glioblastoma, mismatch repair, programmed cell death protein ligand 1 (PD-L1), programmed cell death protein 1 (PD-1), microsatellite instability

1. Introduction

Glioblastoma (GBM) is the most common primary brain tumor with a poor prognosis (47.1%) [1]. GBMs are fatal tumors with a median survival time of 12 months. Approximately 3%–5% of cases live more than 3 years [2]. Many studies are showing the importance of genetic susceptibility, exogenous factors, age and clinical parameters at the time of diagnosis, as the reason for aggressive behavior [3–6]. Other important factors are surgical procedures and multimodal treatment options. Immunotherapy is a promising treatment method that shows a synergistic effect with radiotherapy [7].

Immune checkpoints can be grouped into two main groups as immunostimulating and blocking. The main molecules that inhibit immune checkpoints are the cytotoxic t-lymphocyte-associated protein-4 (CTLA-4), programmed death-1 (PD-1) receptor, and its two ligands (PD-L1 and PD-L2). These molecules block control signals

that lead to the T cell response against the tumor. New treatments are aimed at inhibiting PD-L1 on the tumor cell or PD-1 receptor on the T cell which will produce an antitumoral response. These treatment agents are called immune checkpoint inhibitors (ICIs). CTLA-4, one of the major inhibitory molecules in GBMs, is released from T cells, binding to its ligands (CD80, CD86) reduces the activation and proliferation of effector T cells and increases the activation of regulatory T cells (Treg) in the GBM microenvironment [8,9].

PD-1 is released from T cells and other immune cells. One of its ligands, PD-L1 is found in Tregs in the GBM microenvironment, tumor-associated macrophages, and other cells in the tumor microenvironment, including tumor cells [8,10]. The predictive markers in PD-1/PD-L1 antibody therapy are mainly the number of cytotoxic T-lymphocytes inside tumor tissues and the expression level of PD-L1 in cancer cells [11,12]. In extensive studies,

* Correspondence: emelnur7@gmail.com

PD-L1 expression levels in GBMs were found between 61%–88% [13–15]. High PD-L1 mRNA expression level was found to be associated with shorter overall survival in glioma patients [12,16,17].

Treating recurrent GBMs is more difficult and resistance develops more frequently to treatment. These tumors have been treated several times (radiotherapy and temozolomide) and therefore contain a greater number of mutations. Due to the presence of potential new antigens, these tumors are assumed to be more suitable for the recognition and attack of the immune system. In contrast to the hypothesis that the tumor assumes that it will increase the release of PD-L1 with the immune-escape mechanism developed to protect itself from stronger immune response, in some studies, this rate was found to be significantly lower in recurrent tumors [13,14,18].

PD-1/PD-L1 expression, tumor mutation load, and DNA mismatch repair (MMR) defects are thought to be related to the treatment response. Some clinical studies revealed that defective MMR is associated with clinical responses to immune checkpoint inhibitors (ICI) [5,19]. Tumors with high microsatellite instability (MSI) and high immunogenicity benefit from immunotherapy more and have a better clinical course. Therefore, it is recommended to use MSI status as a marker for response to PD-1/PD-L1 blockade in cancer patients [20].

In the last 6 years, significant results have been obtained in immunotherapy in various tumors (melanoma, renal cell cancer, lung cancer, head, and neck cancers) with anti-PD-1/PD-L1 antibodies. While the response to PD-1 inhibitors is significantly high in lymphoma subtypes (87% in Hodgkin's lymphoma), this rate ranges from 15% to 40% in solid organ cancers [21,22]. There are several studies showing that PD-1 inhibition increases the antitumor responses and survival rate on animal glioma models [23,15]. Especially combined immune checkpoint blockade resulted in 100% long survivors [23]. However, there is insufficient clinical evidence to support its effectiveness in GBM patients. There are some "case reports" in the literature showing that anti-PD-1 therapy (nivolumab) have significant therapeutic effects on GBM patients [19,24,25].

2. Material and methods

2.1 Patients and clinical information

In our study, 74 cases (recurrent and nonrecurrent GBM) diagnosed in our department between 2007–2019 were selected. Twenty-seven of these cases showed recurrence. Hypercellular tumors including palisading necrosis and vascular endothelial proliferation were accepted as 'original tumor'. Diffuse necrosis, vascular hyalinization, gliosis, and the presence of rare atypical cells were accepted as 'radiotherapy effect'. High-grade glioma with

mitosis and minimal evidence of radiation effect is defined as 'recurrent tumor' [26]. The first resection materials of the recurrent cases were defined as "Group 1", the 2nd resection materials were "Group 2" and the nonrecurrent cases were defined as "Group 3". When calculating rates in recurrent cases, cases were considered "positive" if any of the cases in group 1 or 2 were positive. The results of recurrent and nonrecurrent GBM patients were compared. In addition, in recurrent cases, the expression rates in the 1st and 2nd resection materials were compared. For each case, the patient's age at the time of diagnosis, sex, time for recurrence, and survival were recorded. Clinical information was obtained from patient files on the computer.

2.2 Immunohistochemical study

Immunohistochemically, the relationship between MLH1 (MutL Homolog 1), MSH2 (MutS Homolog 2), MSH6 (MutS Homolog 6), and PMS2 (PostMeiotic Segregation increased 2) results and immune checkpoint inhibitors PD-1 and PD-L1 were examined and compared with the clinicopathological features. Sections stained with immunohistochemical antibodies were examined under a light microscope (Olympus BX50). Normal colon tissue was used as a control of the immunohistochemical MLH1, MSH2, MSH6, and PMS2 markers. Expressions of MMR proteins were evaluated as follows: Nuclear staining in more than 80% neoplastic cells was accepted as score 4; 51%–80% staining score 3; 10%–50% staining score 2; less than 10% staining score 1 and no nuclear staining score 0 [27]. Expressions of MMR proteins were shown in Figure 1 (Figure 1: a) Loss of nuclear staining with MLH1 ($\times 200$), b) Nuclear staining with MLH1 ($\times 100$), c) Loss of nuclear staining with MSH2 ($\times 200$), d) Nuclear staining with MSH2 ($\times 100$), e) Loss of nuclear staining with MSH6 ($\times 200$), f) Nuclear staining with with MSH6 ($\times 200$), g) Loss of nuclear staining with with PMS2 ($\times 100$), and h) Nuclear staining with PMS2 ($\times 100$). Loss of nuclear expression of one or more MMR proteins was accepted as deficient mismatch repair [28].

SP263 clone of the PD-L1 antibody was used. Diffuse fibrillar/membranous staining in tumor tissue was evaluated. According to previous studies, no staining in nonnecrotic tumor tissue was accepted as score 0; <25% staining score 1; 25%–50% staining score 2; 50%–75% staining score 3 and > 75% staining score 4 [13]. Membranous staining in epitheloid tumor cells was defined as (+) if > 5% staining in tumor cells as in previous studies [13].

PD-1 expression was seen in lymphocytes in tumor tissue, and perivascular space. It was scored as sparsely, moderately and intensively according to the staining rates in large magnification (200 \times –400 \times). The staining patterns of PD-L1 and PD-1 were shown in Figure 2 (Figure 2: a) Diffuse fibrillary PD-L1 staining in tumor matrix ($\times 200$),




Figure 1. Expressions of MMR proteins.

b) Membranous PD-L1 staining on tumor cells ($\times 400$), c) Diffuse fibrillary and membranous PD-L1 staining ($\times 200$), d) PD-1 staining on tumor infiltrating lymphocytes ($\times 400$).

2.3 Statistical analysis

Descriptive analyses of the study group were given as numbers and percentages. SPSS Statistics v: 22.0 (IBM SPSS, 2013, Armonk, NY, USA) was used for statistical analysis. Comparisons between recurrent and nonrecurrent cases were made using the Chi-square and Fisher's Exact test, and the comparison of the Group 1 and Group 2 with

the Mc Nemar test. The mean and standard deviation values were compared with student t test. In all analyses, the statistical significance level was taken as $p = 0.05$. The effects of recurrence, PD-L1 expression, MMR status, and loss of PMS2 on survival in patients were examined using log-rank test. Survival rates were calculated using the Kaplan–Meier survival analysis.

3. Results

Of the 74 GBM patients, 29 (39.2%) were female and 45 (60.8%) were male with a median age of 58.4 (range,

4–85). However, the mean age of diagnosis in recurrent cases was 52.1 (4–75) and 62.1 (42–85) in nonrecurrent cases and was significantly lower in recurrent GBM cases ($p = 0.007$). Twenty-seven cases (36.5%) recurred, 62 cases (83.8%) died, and 12 cases (25.5%) are still alive. The mean recurrence time was 8.2 months (0.8–39.7). Median survival was 9.9 months (6.1–13.8) in all cases from the time of diagnosis, and 12.5 months in recurrent cases; 6 months for nonrecurrent ones. The clinicopathological characteristics of the patients are given in Table 1.

On immunohistochemical study, PD-L1 expression was observed in 36 (48.6%) of the 74 cases. Expression was detected in 12 (44.5%) of the recurrent cases and 24 (51.0%) of the nonrecurrent cases. Median survival was 7.1 months in patients with PD-L1 (+) and 10 months in patients with PD-L1(-). The effect of PD-L1 expression on survival was

not significant ($p = 0.300$), shown in Figure 3 (Figure 3: a) Recurrent cases b) Nonrecurrent cases c) All cases). Thirteen (17.6%) patients exhibited loss of expression for at least one MMR protein and they were considered to MSI. Nine of the cases with MSI were recurrent (33.3%) and 4 were nonrecurrent GBM (8.5%). The details of the immunohistochemical expression of MMR proteins are given in Tables 2 and 3. Median survival for tumors with MSI was 7 months, and 10 months for those with MSS. However, MMR status did not have a significant effect on survival ($p = 0.953$). Among the MMR proteins, loss of PMS2 was noted in 9 recurrent cases (33.3%) and in 4 nonrecurrent cases (8.5%). Median survival was 7 months in patients with PMS2 loss and 10 months in patients without loss. However, there was no significant effect of PMS2 loss on survival ($p = 0.953$). Loss of PMS2 was found to be significantly effective




Figure 2. The staining patterns of PD-L1 and PD-1.

Table 1. The clinicopathological characteristics of the recurrent and nonrecurrent cases.

Clinical parameters	Recurrent cases (n = 27)	Nonrecurrent cases (n = 47)	Total (n = 74)
Sex (female/male) n (%)	11 (40.7)/16 (59.3)	18 (38.3)/29 (61.7)	29 (39.2)/45 (60.8)
Mean age (± SD)	52.1 (± 15.349)	62.1 (± 10.693)	58.4 (± 13.392)
Mean recurrence time (month) (95% CI)	8.2 (0.8–39.7)	-	-
Median survival time (month) (95% CI)	12.5 (7.4–17.5)	6.0 (3.7–8.2)	9.9 (6.1–13.8)
Death n (%)	27 (100)	35 (74.5)	62 (83.8)

Student t test: $p = 0.02$.

Median survival time: Kaplan–Meier.




Figure 3. The effect of PD-L1 expression on survival.

in relapsed cases ($p = 0.003$). The median recurrence time was 10.1 months in those with PMS2 loss, and 39.7 months in those without. PD-1 expression was observed in 6 cases

(8.1%). It was observed in 3 recurrent cases (11.1%) and 3 nonrecurrent cases (6.3%). The expressions of PD-L1, PD-1 and presence of MSI are given in Table 4.

Table 2. Loss of MMR proteins in the recurrent cases (n = 27, p < 0.05).

MMR proteins	Resection material	n	%	p
MLH1	1st resection	1	3.7	-
	2nd resection	0	0	
MSH2	1st resection	0	0	-
	2nd resection	0	0	
MSH6	1st resection	0	0	-
	2nd resection	2	7.4	
PMS2	1st resection	5	18.5	1.000
	2nd resection	6	22.2	

MMR: Mismatch repair.
McNemar Exact test.

Table 3. Loss of MMR proteins in all cases (n = 74, p < 0.05).

MMR proteins	Cases	n	%	p
MLH1	Recurrent	1	3.7	0.635
	Nonrecurrent	1	2.1	
MSH2	Recurrent	0	0	-
	Nonrecurrent	0	0	
MSH6	Recurrent	2	7.4	0.530
	Nonrecurrent	2	4.3	
PMS2	Recurrent	9	33.3	0.011*
	Nonrecurrent	4	8.5	

MMR: Mismatch repair.
*Statistically significant.
Chi-Square Fisher's Exact test.

Clinical features of Group 1 and 2 (recurrent) cases: 11 (40.7%) were female and 16 (59.3%) were male with the median age was 52.1 (range, 4–75). All of them died. The number of cases with PD-L1 (+) in Group 1 was 8 (29.6%). MSI was detected in 6 of 27 cases (22.2%). The number of the cases with PD-1 (+) in Group 1 was 1 (3.7%). Loss of PMS2 was observed in 5 cases (18.5%). The number of cases with PD-L1 (+) in Group 2 was 8 (29.6%), of which 4 were the same cases in Group 1. MSI was detected in 6 of 27 cases (22.2%). Three of these cases were the same as Group 1. The number of cases with PD-1 (+) was 2 (7.4%), both cases were different from Group 1. Six cases had PMS2 loss (22.2%). Two cases were the same as Group 1 and PMS2 loss was observed in 9 cases totally. Clinical characteristics of Group 3 cases: 18 (38.3%) were

Table 4. The expressions of PD-L1, PD-1, and presence of MSI in the recurrent and non-recurrent cases (n = 74, p < 0.05).

Protein expression	Cases	n	%	p
PD-L1	Recurrent	12	44.5	0.060
	Nonrecurrent	24	51.0	
PD-1	Recurrent	3	11.1	0.536
	Nonrecurrent	3	6.4	
MSI	Recurrent	9	33.3	0.011*
	Nonrecurrent	4	8.5	

MSI: Microsatellite instability.
*Statistically significant.
Chi-Square Fisher's Exact test.




Figure 4. The relationship between MMR status and PD-L1 expression in recurrent and nonrecurrent GBM

women and 29 (61.7%) were men with the median age was 62.1 (42–85). Thirty five cases (74.5%) died and 12 cases (25.5%) are still alive.

The relationship between MMR status and PD-L1 expression in recurrent and nonrecurrent cases was shown in Figure 4 (p = 0.448 and p = 0.348, respectively, Chi-square Fisher's exact test). There was no significant difference between PMS2 and PD-L1 expression in patients with or without recurrence (p = 1.000 and p = 0.348, respectively, Chi-square Fisher's exact test). There was no significant difference between PD-L1 expression and survival in patients with or without recurrence (p = 0.136, log rank (Mantel Cox) test). There was no significant difference between MMR status and survival in patients with or without recurrence (p = 0.133, log rank (Mantel Cox) test). There was no significant difference between

loss of PMS2 and survival in patients with or without recurrence ($p = 0.133$, log rank (Mantel Cox) test).

4. Discussion

The ability of GBM to cause local and systemic immunosuppression limits the innate defense and adaptive immunotherapy effect against the tumor, and thus prevents the development of new therapies. The process of immunosuppression is not only related to abnormal PD-L1 expression in GBM cells, but also to the tumor microenvironment.

In one of the most recent studies to elucidate immunotherapy resistance mechanisms in GBM, it was seen that low PD-L1 expression, low tumor mutation burden and T lymphocytes, which are largely depleted in the tumor, are indicators of decreased antitumor immunity [21].

However, PD-L1 expression levels are observed in a highly variable range in GBMs. Although it was seen between 61% and 88% in two studies with large patient groups, the median percentage of PD-L1 expression on tumor cells in the study of Nudom et al. was 2.7% [13–15]. Berghof et al. said that the rate of PD-L1-positive cases in GBM was quite higher (72% in recurrent; 88% in newly diagnosed GBM) than melanoma cases (30%) and nonsmall cell lung cancer cases (25%–36%) [13]. In recurrent GBM cases with higher mutations and where immunotherapy resistance is expected less, we also found PD-L1 expression lower than those which are nonrecurrent (44.5% and 51.0%, respectively), similar to the studies of Berghof, Heynckes, and Ndom [13,14,18].

While low PD-L1 expression levels are associated with treatment resistance, some studies have shown that high PD-L1 levels are associated with shorter overall survival in glioma patients [12,17,29]. On the other hand, in several other studies no significant relationship was found between PD-L1 expression and survival [12,13,16,30]. In our study, no significant difference was found between PD-L1 expression level and survival.

In tumors with deficient MMR, 10 to 100 times more somatic mutations were found compared to those which are proficient [21,31]. Microsatellite instability is not high in GBM. Patients with MSI are generally young and have colorectal cancer at the same time [32]. In a study conducted with 30 different tumors, it was stated that the neontigen burden in GBM was in the lower third section [21,33]. In GBM, mutations in MMR genes are thought to be associated with resistance to therapy and thus tumor recurrence [8]. Martine et al. observed that the presence of MSI was at a significantly higher rate in patients with recurrent GBM and stated that this may be associated with malignant progression [34]. We have also found the rate of MSI significantly higher in recurrent patients than the nonrecurrent ones (33.3%, and 8.5% respectively).

GBM specimens containing MSH6 mutations have been described as hypermutator phenotypes [35,36]. Shinsato et al. found reduced levels of MLH1 and PMS2 related to therapy resistance and recurrence [37]. We have also found a significant elevation in PMS2 loss in all groups (Table 2–3). PMS2 loss, which was observed more clearly in recurrent cases, suggested that this change might be a marker for malignant progression. However, we could not find a significant relationship between the loss of PMS2 neither with survival nor PD-L1 expression.

In one of the studies on the role of the status of MSI in predicting immunotherapy response, it was observed that colorectal cancers with MMR deficiency had a high response to PD-1 inhibitor therapy [30]. In another study, the research was expanded and the effectiveness of PD-1 blockade was evaluated in 12 different tumor types with MMR deficiency at an advanced stage and it was seen that 21% of the patients had a complete response and 53% of the patients had an objective radiological response [38]. In high-grade urothelial carcinomas, it was shown that MMR deficiency (loss of MSH2 and MSH6) is associated with increased PD-L1 expression [8,39]. PD-L1 and PD-L1 expressions were found high in colorectal and endometrial cancers with microsatellite instability (MSI) [8,40].

In conclusion, MMR status is suggested as a marker for response to PD-1/PD-L1 blockade in other cancer types. However, we found that PD-L1 expression was low and MSI rate was higher in recurrent GBM than nonrecurrent cases and we could not find a significant relationship between these two entities. The presence of higher MSI in the patients with recurrent GBM in this study indicates the importance of these proteins as a predictive markers. However, low PD-L1 expression levels suggest that this antibody may not be a good predictive marker for determining the group of patients who will receive immunotherapy.

In recent years, an increasing number of clinical trials are available to try different combinations in GBM treatment. We also think that different combinations with other immune checkpoint proteins can be tried in GBM patients to determine both prognostic and therapeutic efficacy.

The limitation of our study was that the clinical data about treatment modalities are incomplete and, therefore, were not included in the article.

Acknowledgment and/or disclaimers

This project (Project no: TSB-2019-7796) was supported by the Karadeniz Technical University Scientific Research Projects Support Fund. Approval of Karadeniz Technical University Faculty of Medicine Ethics Committee was obtained before starting the study (26.07.2018- Protocol no: 2018/167).

References

1. Ostrom QT, Gittleman H, Liao P, Vecchione-Koval T, Wolinsky Y et al. CBTRUS Statistical Report: primary brain and other central nervous system tumors diagnosed in the United States in 2010-2014. *Neuro-Oncology* 2017; 19 (5):1-88. doi: 10.1093/neuonc/now158
2. Krex D, Klink B, Hartmann C, Von Deimling A, Pietsch T et al. Long-term survival with glioblastoma multiforme. *Brain* 2007; 130 (10): 2596-2606. doi: 10.1093/brain/awm204
3. Curran WJ Jr, Scott CB, Horton J, Nelson JS, Weinstein AS et al. Recursive partitioning analysis of prognostic factors in three Radiation Therapy Oncology Group malignant glioma trials. *Journal of the National Cancer Institute* 1993; 85 (9): 704-710. doi: 10.1186/1471-2407-9-450
4. Zheng T, Cantor KP, Zhang Y, Chiu BC, Lynch CF. Risk of brain glioma not associated with cigarette smoking or use of other tobacco products in Iowa. *Cancer Epidemiology, Biomarkers & Prevention* 2001; 10 (4): 413-414.
5. Huncharek M, Kupelnick B, Wheeler L. Dietary cured meat and the risk of adult glioma: a meta-analysis of nine observational studies. *Journal of Environmental Pathology, Toxicology and Oncology* 2003; 22 (2): 129-137. doi: 10.1615/JEnvPathToxOncol.v22.i2.60
6. Kleihues P, Schäuble B, Zur Hausen A, Estève J, Ohgaki H. Tumors associated with p53 germline mutations: a synopsis of 91 families. *The American Journal of Pathology* 1997; 150 (1): 1-13.
7. Reardon DA, Gokhale PC, Klein SR, Ligon KL, Rodig SJ et al. Glioblastoma eradication following immune checkpoint blockade in an orthotopic, immunocompetent model. *Cancer Immunology Research* 2016; 4 (2): 124-135. doi: 10.1158/2326-6066.CIR-15-0151
8. Kurz SC, Wen PY. Quo Vadis-do immunotherapies have a role in glioblastoma? *Current Treatment Options in Neurology* 2018; 20 (5): 14. doi: 10.1007/s11940-018-0499-0
9. Wing K, Onishi Y, Prieto-Martin P, Yamaguchi T, Miyara M et al. CTLA-4 control over Foxp3+ regulatory T cell function. *Science* 2008; 322 (5899): 271-275. doi: 10.1126/science.1160062
10. Bloch O, Crane CA, Kaur R, Safaee M, Rutkowski MJ et al. Gliomas promote immunosuppression through induction of B7-H1 expression in tumor-associated macrophages. *Clinical Cancer Research* 2013; 19 (12): 3165-3175. doi: 10.1158/1078-0432
11. Chen PL, Roh W, Reuben A, Cooper ZA, Spencer CN et al. Analysis of immune signatures in longitudinal tumor samples yields insight into biomarkers of response and mechanisms of resistance to immune checkpoint blockade. *Cancer Discovery* 2015; 6 (8): 827-837. doi: 10.1158/2159-8290
12. Chen RQ, Liu F, Qiu XY, Chen XQ. The Prognostic and therapeutic value of PD-L1 in glioma. *Frontiers in Pharmacology* 2019; 9: 1503. doi: 10.3389/fphar.2018.01503
13. Berghoff AS, Kiesel B, Widhalm G, Rajky O, Ricken G et al. Programmed death ligand 1 expression and tumor-infiltrating lymphocytes in glioblastoma. *Neuro-Oncology* 2015; 17 (8): 1064-1075. doi: 10.1093/neuonc/nou307
14. Nduom EK, Wei J, Yaghi NK, Huang N, Kong LY et al. PD-L1 expression and prognostic impact in glioblastoma. *Neuro-Oncology* 2016; 18 (2): 195-205. doi: 10.1093/neuonc/nov172
15. Preusser M, Berghoff AS, Wick W, Weller M. Clinical Neuropathology mini-review 6-2015: PD-L1: emerging biomarker in glioblastoma? *Clinical Neuropathology* 2015; 34 (6): 313-321. doi: 10.5414/NP300922
16. Zeng J, Zhang XK, Chen HD, Zhong ZH, Wu QL et al. Expression of programmed cell death-ligand 1 and its correlation with clinical outcomes in gliomas. *Oncotarget* 2016; 7 (8): 8944-8955. doi: 10.18632/oncotarget.6884
17. Xue S, Song G, Yu J. The prognostic significance of PD-L1 expression in patients with glioma: a meta-analysis. *Scientific Reports* 2017; 7 (1): 4231. doi: 10.1038/s41598-017-04023-x
18. Heynckes S, Gaebelein A, Haaker G, Grauvogel J, Franco P et al. Expression differences of programmed death ligand 1 in de-novo and recurrent glioblastoma multiforme. *Oncotarget* 2017; 8 (43): 74170-74177. doi: 10.18632/oncotarget.18819
19. Bouffet E, Larouche V, Campbell BB, Merico D, De Borja R et al. Immune checkpoint inhibition for hypermutant glioblastoma multiforme resulting from germline biallelic mismatch repair deficiency. *Journal of Clinical Oncology* 2016; 34 (19): 2206-2211. doi: 10.1200/JCO.2016.66.6552
20. Llosa NJ, Cruise M, Tam A, Wicks EC, Hechenbleikner EM et al. The vigorous immune microenvironment of microsatellite instable colon cancer is balanced by multiple counter-inhibitory checkpoints. *Cancer Discovery* 2015; 5 (1): 43-51. doi: 10.1158/2159-8290
21. Adhikaree J, Moreno-Vicente J, Kaur AP, Jackson AM, Patel PM. Resistance mechanisms and barriers to successful immunotherapy for treating glioblastoma. *Cells* 2020; 9 (2): 263. doi: 10.3390/cells9020263
22. Zou W, Wolchok JD, Chen L. PD-L1 (B7-H1) and PD-1 pathway blockade for cancer therapy: mechanisms, response biomarkers, and combinations. *Science Translational Medicine* 2016; 8 (328): 328rv4. doi: 10.1126/scitranslmed.aad7118
23. Zeng J, See AP, Phallen J, Jackson CM, Belcaid Z et al. Anti-PD-1 blockade and stereotactic radiation produce long-term survival in mice with intracranial gliomas. *International Journal of Radiation Oncology Biology, Physics* 2013; 86 (2): 343-349. doi: 10.1016/j.ijrobp.2012.12.025
24. Roth P, Valavanis A, Weller M. Long-term control and partial remission after initial pseudo progression of glioblastoma by anti-PD-1 treatment with nivolumab. *Neuro-Oncology* 2017; 19 (3): 454-456. doi: 10.1093/neuonc/now265

25. Simonelli M, Di Tommaso L, Baretta M, Santoro A. Pathological characterization of nivolumab-related liver injury in a patient with glioblastoma. *Immunotherapy* 2016; 8 (12): 1363-1369. doi: 10.2217/imt-2016-0057
26. Tihan T, Barletta J, Parney I, Lamborn K, Sneed PK et al. Prognostic value of detecting recurrent glioblastoma multiforme in surgical specimens from patients after radiotherapy: should pathology evaluation alter treatment decisions?. *Human Pathology* 2006; 37 (3): 272-282. doi: 10.1016/j.humpath.2005.11.010
27. Felsberg J, Thon N, Eigenbrod S, Hentschel B, Sabel MC et al. Promoter methylation and expression of MGMT and the DNA mismatch repair genes MLH1, MSH2, MSH6 and PMS2 in paired primary and recurrent glioblastomas. *International Journal of Cancer* 2011; 129 (3): 659-670. doi: 10.1002/ijc.26083
28. Collage of American pathologist (2018). Template for Reporting Results of DNA Mismatch Repair Testing in Patients Being Considered for Checkpoint Inhibitor Immunotherapy (online). Website: <https://documents.cap.org/protocols/cp-general-dnamismatchrepair-18biomarker-1001.pdf> [accessed 09.10.2020].
29. Wang Z, Zhang C, Liu X, Wang Z, Sun L et al. Molecular and clinical characterization of PD-L1 expression at transcriptional level via 976 samples of brain glioma. *Oncoimmunology* 2016; 5 (11): e1196310. doi: 10.1080/2162402X.2016.1196310
30. Miyazaki T, Ishikawa E, Matsuda M, Akutsu H, Osuka S et al. Assessment of PD-1 positive cells on initial and secondary resected tumor specimens of newly diagnosed glioblastoma and its implications on patient outcome. *Journal of Neuro-Oncology* 2017; 133 (2): 277-285. doi: 10.1007/s11060-017-2451-7
31. Le DT, Uram JN, Wang H, Bartlett BR, Kemberling H et al. PD-1 blockade in tumors with mismatch-repair deficiency. *The New England Journal of Medicine* 2015; 372 (26): 2509-2520. doi: 10.1056/NEJMoa1500596
32. Leung SY, Chan TL, Chung LP, Chan AS, Fan YW et al. Microsatellite instability and mutation of dna mismatch repair genes in gliomas. *The American Journal of Pathology* 1998; 153 (4): 1181-1188. doi: 10.1016/S0002-9440(10)65662-3
33. Schumacher TN, Schreiber RD. Neoantigens in cancer immunotherapy. *Science* 2003; 348 (6230): 69-74. doi: 10.1126/science.aaa4971
34. Martinez R, Schackert HK, Plaschke J, Baretton G, Appelt H et al. Molecular mechanisms associated with chromosomal and microsatellite instability in sporadic glioblastoma multiforme. *Oncology* 2004; 66: 395-403. doi: 10.1159/000079488
35. Hodges TR, Ott M, Xiu J, Gatalica Z, Swensen J et al. Mutational burden, immune checkpoint expression, and mismatch repair in glioma: implications for immune checkpoint immunotherapy. *Neuro-Oncology* 2017; 19 (8): 1047-1057. doi: 10.1093/neuonc/nox026
36. Cahill DP, Levine KK, Betensky RA, Codd PJ, Romany CA et al. Loss of the mismatch repair protein MSH6 in human glioblastomas is associated with tumor progression during temozolomide treatment. *Clinical Cancer Research* 2007; 13 (7): 2038-2045. doi: 10.1158/1078-0432.CCR-06-2149
37. Shinsato Y, Furukawa T, Yunoue S, Yonezawa H, Minami K et al. Reduction of MLH1 and PMS2 confers temozolomide resistance and is associated with recurrence of glioblastoma. *Oncotarget* 2013; 4 (12): 2261-2270. doi: 10.18632/oncotarget.1302
38. Le DT, Durham JN, Smith KN, Wang H, Bartlett BR et al. Mismatch-repair deficiency predicts response of solid tumors to PD-1 blockade. *Science* 2017; 357 (6349): 409-413. doi: 10.1126/science.aan6733
39. Castro MP, Goldstein N. Mismatch repair deficiency associated with complete remission to combination programmed cell death ligand immune therapy in a patient with sporadic urothelial carcinoma: immunotheranostic considerations. *Journal for Immunotherapy of Cancer* 2015; 3: 58. doi: 10.1186/s40425-015-0104-y
40. Yamashita H, Nakayama K, Ishikawa M, Nakamura K, Ishibashi T et al. Microsatellite instability is a biomarker for immune checkpoint inhibitors in endometrial cancer. *Oncotarget* 2018; 9 (5): 5652-5664. doi: 10.18632/oncotarget.23790