

Hematological status of avian species along a metal pollution gradient at Sialkot, Pakistan

Ajmal SHEHZAD¹, Khalid Mahmood ANJUM^{1,*}, Atif YAQUB²,

Muhammad Zubair YOUSAF³, Sarwar Allah DITTA², Junaid NASEER⁴

¹Department of Wildlife and Ecology University of Veterinary and Animal Sciences Lahore, Pakistan

²Department of Zoology Government College University Lahore, Pakistan

³School of life sciences Forman Christian College Lahore, Pakistan

⁴Department of Forestry & Range Management University of Agriculture Faisalabad, Pakistan

Received: 26.11.2020

Accepted/Published Online: 04.11.2021

Final Version: 18.01.2022

Abstract: The concentration of metals and effects of these metals on hematological parameters were examined in avian species by using feathers and blood samples collected from different sites of Sialkot. The major reason for significant differences in hematological parameters and concentrations of metal contents between different species was the difference in feeding habitat. Metal concentrations in feathers and blood samples indicated that the chromium and lead were not found at a safe level. The results did not revealed a significant negative association between metal contents and hematological parameters but concluded that increased metal contents exert negative effects on hematological status especially lead and chromium, while white blood cells showed a significant positive correlation with increased metal contents except cadmium. The feathers accumulate highest concentration of metals than blood, the most probable reason behind that is the feathers reflect exogenous contamination and blood only represents the immediate dietary sources of contamination. It concluded that the feathers of avian species can be used as an efficient bio-indicator to assess the metal contamination level in an environment, and the assessment of hematological parameters can be used to find out the negative impact of metal contamination on health status. The findings of this study will help to manage environmental pollution and its impacts on wildlife.

Key words: Environmental pollution, feathers, heavy metals, hematological parameters, sialkot

1. Introduction

Environmental degradation takes place due to the increasing human population and the development of technology (Ayyamperumal et al., 2006; Qadir and Malik, 2011; Xu et al., 2013). The environment is contaminated with the passage of time by different types of contaminants, out of which metals are important pollutants because metals are non-biodegradable. Both natural and anthropogenic activities are sources of metal contamination. The metals are persistent, so these metals accumulate in the body and the level of these metals increases at each trophic level. The presence of metallic contaminants is the major threat to the natural ecosystem (Deng et al., 2007; Naccari et al., 2009). Various previous studies revealed that the accumulation of metallic contaminants causes many adverse effects such as disorders of endocrine and nervous system, mutations in genes, and abnormalities in physiology and behavior (Burger and Gochfeld, 2000; Dauwe et al., 2004; Martin et al., 2003). The number of red blood cells, amount of hemoglobin, and haematocrit represents the oxygen binding capacity of blood (Ots, et al. 1998). The exposure

to lead is directly associated with the inhibition of enzyme δ -aminolevulinic acid dehydratase (ALAD), which affects the formation of hemoglobin (Papanikolaou, et al. 2005). Zinc is an essential element to perform the normal functioning of ALAD, but lead replaces this element, which affects the normal functioning of this enzyme. ALAD is a useful biomarker for the study of metal contamination and its effects. Several studies concluded that cadmium and lead cause anemia by decreasing the level of hemoglobin, hematocrit, and mean cell volume (Iolascon, et al. 2009). Cadmium and lead are considered as most dangerous heavy metals, which come into the environment by burning fossil fuels, vehicles, and industries (Kenntner et al., 2003). Several previous studies revealed that lead and cadmium are very toxic for avian fauna such as causing abnormalities in endocrine system, molting, reproduction, growth rate, and enzymes involved in hemoglobin formation (Cheney et al., 1981; Eisler, 1988; Honda et al., 1986). Nickel is used as a catalyst in different industries is deposited into the natural water bodies, which are taken up by different organisms and damage their respiratory organs and DNA.

* Correspondence: khalid.mahmood@uvas.edu.pk

Zinc is used in various kinds of industries such as cosmetic industry, electric industry, and agrochemical industry causes nausea, stomach disorders, and fainting (Peplow, 2000; Pérez-López et al., 2008). In south Asia, the metal pollution has increased in a few years (Hardoy et al., 1992).

Due to the increasing metal pollution and its adverse effects, we need to assess the environmental status, but we cannot estimate all of the parts of the environment, so we can use a bio-indicator, which represent the environmental conditions (Brait and Antoniosi Filho, 2011; Falq et al., 2011; Scheifler et al., 2006). The appropriate bio-indicator is required to estimate the concentration and hazardous effects of these metals in the environment. The bio-indicators are living organisms, which represent their environmental status correctly (Burger and Gochfeld, 2007; Kalisińska et al., 2004). Avian species are considered as an important and appropriate bio-indicator still from 1960s. Birds are considered as good bio-indicators because they feed on diverse types of food and can represent primary fluctuations in food chain (Boncompagni et al., 2003). Avian species can be used to estimate environmental contamination because they are widely distributed in the environment and take food from different types of sources (Furness, 1993; Llabjani et al., 2012). The use of the internal organs of birds has proven to be a good bio-indicator, but it is a destructive method because of many ethical and conservative issues related to the use of internal organs (Jaspers et al., 2005; Van den Steen et al., 2006). Birds are liable to environmental pollutants either from contaminated air, drinking water, and dietary intake. The metals accumulate in the body and become the part of bloodstream, and metals are integrated with the keratin structure in the feathers (Tsipoura et al., 2008). In Europe and United States, avian species are used as bio-indicator to estimate environmental pollution (Burger 2013; Frantz et al., 2012). Feathers can be used as an alternative bio-indicator without sacrificing the bird because they are easy to collect, transport and store; moreover, feathers can be collected without affecting the fitness of birds (Spahn and Sherry, 1999).

The estimation of metal pollution does not give us any information about the biological stress occurring due to metal contamination. Therefore, we need an approach to estimate the negative effects of metal contamination on health parameters of avian fauna (Peakall, 1992). The assessment of hematological parameters can be used to assess the health status and biological stress. The normal hematological status is compulsory for the survival of avian fauna (Kilgas et al., 2006; Nadolski et al., 2006). Many birds do not show the symptoms of diseases at early stages of illness. Hematological parameters play an important role to evaluate the health status, so we can recognize many diseases at early stages by the use of hematological assays (Scope et al., 2005). Many features are related with

hematological values such as source of food, day length, molting period, and environmental conditions (Millaku et al., 2000). The estimation of metal contamination is very useful tool to assess the environmental status. The evaluation of health risks associated with metal contamination is very necessary for the conservation of flora and fauna (Herrera-Duenas et al., 2014). The abnormalities in hematological parameters are associated with breeding capabilities of avian species such as low level of hematocrit associated with low body conditions (Svensson and Merila, 1996), infectious diseases by parasites of blood (Booth and Elliott, 2002), and also flying abilities (Saino et al., 1997).

Like other developing countries, Pakistan is also facing the problem of population development, urbanization, and industrialization in recent years, which has resulted in environmental contamination (Khan, 1991; Qadir et al., 2008; Zaman and Ara, 2000). The contamination of environment by metal pollutants is more in industrial cities (Hashmi et al., 2013). The use of agrochemicals, fertilizers, and industrial effluents are the major sources of metal contamination in Pakistan (Baluch, 1995). Several studies revealed higher concentration of heavy metals in water bodies of Pakistan (Tariq et al., 1994; Tehseen et al., 1994). Punjab is most developed, urbanized, and industrialized province of Pakistan. During the last few years, the environmental pollution has increased due to human activities. The industrial and agricultural activities are dangerous for water bodies, fishes, birds, soil, and sediments (Eqani et al., 2012; Hashmi et al., 2013; Qadir and Malik, 2011; Syed and Malik, 2011). Sialkot city is internationally famous for its sports, leather, and surgical industries. The effluents, solid wastes, and air particles emitted from leather industries are the major sources of environmental contamination. The leather industries use seventeen different types of tanning agents, but chromium usage is ninety percent (Iqbal et al., 1998). This research was conducted to estimate the concentration of metals and the negative impact of metals contaminants on hematological parameters in the selected avian fauna from Sialkot to assess the environmental contamination and biological stress resulted due to the metal contamination.

2. Materials and methods

2.1. Study area

Samples were collected from different sites of Sialkot, Pakistan. In the last decade, due to increase in population size of the city, development of industries has taken place. These include tanneries, steel factories, leather industry, garments industry, and sports industry. Sialkot generated nineteen million-meter cubes per year of wastewater. There is no water treatment facility available in Sialkot, and this water is drained into natural water bodies.

2.2. Sampling

Fifty-eight birds were captured from different sites of Sialkot by using mist net with mesh size of 30–38 mm (Gushit et al., 2016). The chest feathers samples were collected from house sparrow ($n = 21$), common myna ($n = 24$) and common quail ($n = 13$) and stored in a paper envelop for the analysis of metal contaminants. Twenty-seven blood samples were collected by using 1mL syringe from the jugular or ulnar vein of house sparrow ($n = 9$), common myna ($n = 11$) and common quail ($n = 7$) and then shifted into EDTA containing vacutainer for hematological and metal contents analysis. The blood samples were stored at 4 °C immediately (Geens et al., 2010). All of the methods, which were used in this study, are approved by the ethical committee of the University of veterinary and animal sciences Lahore-Pakistan.

2.3. Metal analysis

The feather samples were washed with tap water and then cleaned with distilled water and acetone (3 times) for the removal of any external contaminants. The feather samples were air dried before drying in an oven. Feather samples transferred into 4 mL polypropylene vials and dried in oven for 24 h at 60 °C. The feathers were cut into small pieces by using stainless steel scissor to allow easy digestion with acid. The 0.01 mg dry weight of feather samples was calculated by digital balance. The remaining blood samples after hematological analysis were taken in polypropylene vials. The blood samples were dried in an oven at 60 °C for 24 h. A 100 mL stock solution was prepared with a (1:1) ratio by volume with a mixture of seventy percent HNO₃ (Merck, Germany) and thirty percent H₂O₂ (Merck, Germany), respectively per 10 µg dry sample. The digestion of samples by microwave heating continued until the mixture became pale yellow. The completely digested samples were filtered and diluted with 2 mL of double distilled water (MilliQ, Milipore). The samples were stored at -20 °C in a sample bottle before further analysis. The concentrations of chromium, nickel, copper, zinc, cadmium and lead were analyzed by atomic absorption spectrophotometer (SHIMADZU AA-7000F/AAC). The limits of quantification for all metals were set at 0.01 µg/L. For quality control blanks and certified reference material (VMK 102, Bureau of Certified References) were used in analyses. The concentration of metals in certified samples were within 10% of the certified values. The concentrations of metals were represented on fresh weight basis in µg/g (Geens et al., 2010).

2.4. Hematological analysis

The following hematological parameters were analyzed: the number of erythrocytes (RBC in 10⁶/µL counting of leucocytes (WBC in 10³/µL), differential leukocytes count, mean cell volume (MCV in fL), mean cell hemoglobin (MCH in pg), and mean cell hemoglobin concentration

(MCHC in g/dL) by using automated hematology analyzer NIHON KOHDEN, MEK-6550.

2.5. Statistical analysis

SPSS software was used to analyze data. The descriptive analysis was used to find out the means and standers error. One way analysis of variance ANOVA was applied to check the significance level for mean values among different species. The significant differences were analyzed by performing Tukey HST test. Pearson correlation was performed to find out the relationship between metals in feathers and blood. The linear regression was performed to estimate the correlation between metals in blood and hematological parameters. $P < 0.05$ was considered significant for all analyses.

3. Results

3.1. Metals concentration

The results of this study revealed significant variations for concentrations of metals in feathers between different species. The concentrations of copper and nickel in blood not varied significantly, but all other metals vary significantly among house sparrow, common myna, and quail. The highest concentrations of lead, chromium, nickel, and copper in feathers were examined in common myna. The Quail revealed highest concentration of cadmium and zinc both in feathers and blood. The highest concentration of chromium and lead were found in blood of common myna. The mean concentrations of metals in blood and feathers with ANOVA significance are given in (Table 1). The concentrations of metals at different study spots in feather samples are represented in Figure 1 and the ones in blood samples are represented in Figure 2. We also find the relationship between metal levels in feather and blood samples by performing Pearson correlation. The results of this study founded a significant positive association between Cu ($r = 0.972$, $p = 0.000$), Zn ($r = 0.979$, $p = 0.000$), Ni ($r = 0.761$, $p = 0.001$), Cd ($r = 0.897$, $p = 0.000$) Cr ($r = 0.911$, $P = 0.006$) and Pb ($r = 0.879$, $p = 0.001$) levels in blood and feather samples of avian species.

3.2. Hematological parameters

The one way analysis of variance revealed the significant variation in hematological parameters among house sparrow, common myna, and quail. The mean concentrations of hematological parameters and ANOVA significances are given in (Table 2). The highest concentration of RBCs, MCH, and MCHC were observed in quail than other species. The level of WBCs was higher in common myna. The sparrows revealed the highest concentration of MCH among three species. The results of this research did not show significant variations between house sparrow and common myna; however, both were significantly different from common quail in terms of the level of red blood cells ($p = 0.000$, Tukey HST test p

Table 1. Mean concentration ($\mu\text{g/g}$ fresh weight \pm S.E) and one way ANOVA results for metal contents in feathers and blood samples.

Metals	Sample	Species			ANOVA Sig.
		House Sparrow (<i>Passer domesticus</i>)	Common Myna (<i>Acridotheres tristis</i>)	Quail (<i>Coturnix coturnix</i>)	
Copper	Feathers	19.49 \pm 1.59 ^a	23.95 \pm 2.28 ^a	11.87 \pm 1.42 ^b	0.002**
	Blood	0.62 \pm 0.21 ^a	0.63 \pm 0.19 ^a	0.22 \pm 0.04 ^a	0.200
Zinc	Feathers	72.85 \pm 5.87 ^b	66.74 \pm 5.90 ^b	139.63 \pm 12.89 ^a	0.000**
	Blood	6.02 \pm 0.50 ^b	5.18 \pm 0.81 ^b	14.18 \pm 1.67 ^a	0.000**
Nickel	Feathers	7.09 \pm 0.71 ^{ab}	8.69 \pm 1.25 ^a	3.87 \pm 0.74 ^{bc}	0.011*
	Blood	0.47 \pm 0.22 ^a	0.68 \pm 0.23 ^a	0.43 \pm 0.18 ^a	0.696
Cadmium	Feathers	1.52 \pm 0.29 ^b	1.92 \pm 0.33 ^b	5.53 \pm 0.85 ^a	0.000**
	Blood	0.30 \pm 0.18 ^b	0.11 \pm 0.04 ^b	1.32 \pm 0.14 ^a	0.000**
Chromium	Feathers	24.38 \pm 2.85 ^b	45.35 \pm 2.69 ^a	16.47 \pm 2.28 ^b	0.000**
	Blood	1.80 \pm 0.15 ^b	2.92 \pm 0.37 ^a	1.11 \pm 0.20 ^b	0.001**
Lead	Feathers	9.16 \pm 1.17 ^a	14.93 \pm 1.54 ^a	4.45 \pm 0.63 ^b	0.000**
	Blood	0.36 \pm 0.11 ^b	1.30 \pm 0.20 ^a	0.29 \pm 0.17 ^b	0.002**

< 0.05) and mean corpuscular hemoglobin concentration ($p = 0.00$, Tukey HST test $p < 0.05$). Common myna and common quail did not reveal significant variations, while both varies significantly from house sparrow in terms of the numbers of white blood cells ($p = 0.007$, Tukey HST test $p < 0.05$) and mean cell volume ($p = 0.000$, Tukey HST test $p < 0.05$). We also estimated the percentage of differential leukocyte count in white blood cells. The house sparrow and common myna shows significant differences with quail for the percentages of lymphocytes, monocytes, heterophils, and basophils as shown in (Table 3).

3.3. Relation between metal pollution and hematological parameters

The results of this research not reported significant correlation between metals concentrations in blood and hematological parameters. However, we find some negative correlations between blood metal contents level and hematological parameters. The cadmium showed negative relation with WBCs and MCV. The results revealed negative association between MCV, MCH, and chromium. The lead showed negative relation with all hematological parameters except white blood cells. According to ANOVA results of linear regression, white blood cells revealed significance positive correlation with metal contaminants. The results of correlation between metals and hematological parameters are given in (Table 4).

4. Discussion

Copper is required in small amount to perform some normal metabolic activities. However, the exposure of copper in large amount may lead to serious damages such

as abnormalities of the reproductive system, respiratory system, digestive system, and endocrine system. The accumulation of copper in large amount may also damage blood cells and may cause cancer (Chen et al., 1993). The main sources of copper in the environment are mining activities, sewage sludge, and agricultural activities such as usage of fungicides (Ullah et al., 2014). The results of this study concluded that the concentration of copper in blood and feathers were present in the following descending order between three species common myna > house sparrow > quail. According to analysis of variance the concentration of copper in feathers varies significantly between species. However, the results for the concentration of copper in blood do not vary significantly between house sparrow, common myna, and quail. The level of copper was not differing significantly between feathers of sparrow and common myna, but sparrow and common myna revealed significant variations with quail. According to our research, the increased level of copper in blood results in increased level of copper in feathers. The concentration of copper not revealed significant relation with hematological parameters. The results of house sparrow for concentration of copper in feathers were comparable with the study of Manjula et al., 2015; Swaileh and Sansur, 2006. The copper concentration in blood samples of sparrow and common myna were comparable with the one (Lester and van Riper, 2014) in blood of song sparrow. The results of (Manjula et al., 2015) for rural area were comparable with the copper contents in common myna used in the present study; however, the concentration of copper in urban area were higher than the one found in the present study. The





Figure 1. Maps showing concentrations of metal ($\mu\text{g/g}$) in feather samples on different sampling sites of Sialkot.

concentration of copper in our study was higher than the results of (Abbasi et al., 2015; Mustafa et al., 2015).

Zinc is an essential element for many metabolic reactions. The zinc originated in environment as a result of both human and natural activities. The burning of solid wastes, construction activities, and marble industries are also sources of zinc pollution (Ullah et al., 2014). The usage of zinc salts is the major source of zinc contamination in our study area (Abdullah et al., 2015). There are no sides effects of zinc are known, but some studies reported 1200 $\mu\text{g/g}$ of Zn as threshold limit (Taggart et al., 2009). The highest level of zinc was founded in quail birds between three species. The level of zinc contents was different significantly between house sparrow, common myna, and quail. According to our estimations, zinc does not affect significantly hematological parameters. However, zinc only showed positive association with white blood cells. The concentration of zinc in house sparrow and common myna feathers was comparable with the results of Manjula et al., 2015; Abbasi et al., 2015. The results of our study detected higher concentration of Zn in house sparrow

than the concentration found by Mustafa et al., 2015; Swaileh and Sansur, 2006. The uptake of zinc in common myna feathers were higher in the study of Muralidharan et al., 2004. The zinc level in quail was almost similar with the results of Malik and Zeb, 2009, which were reported for cattle egret. The level of zinc in our blood samples was too much lower than the results of Bounagua et al., 2014. However, the zinc level in blood samples of house sparrow and common myna were similar with the results of Geens et al., 2010 for great tit and Lester and van Riper, 2014 for song sparrow. However, zinc was found at safe level in our study samples. The nickel exposure may cause some allergic reactions, damages to bronchitis, digestive abnormalities and as well nickel is a carcinogen (Abdullah et al., 2015). The industrial activities such as ghee production, batteries manufacturing, leather production, and Ni-Cr plating are the major reasons of nickel contamination (Ullah et al., 2014). According to (Outridge and Scheuhammer, 1993) the birds from unpolluted area accumulate nickel in range of 0.1–5 $\mu\text{g/g}$. The level of nickel was quite different in feather samples between three species, but, in blood



different sampling sites of Sialkot.

and devices coated with metals (Malik and Malik, 2011). The accumulation of cadmium in the body of organisms with the passage of time. The higher concentration of cadmium causes thinning, damages of kidney and calcium metabolism, reproductive system, immune system (Burger, 2008). The concentration of cadmium in different species were founded in following order: quail > common myna > house sparrow. The levels of cadmium in quail blood and feathers were found to be different from sparrow and common myna. According to our results, cadmium was not significantly associated with hematological parameters. There was no significant negative relation between cadmium concentration in blood and white blood cells. The results

concentration in sparrow and common
st similar with the results of (Lester and
Manjula et al., 2015; Muralidharan et al.,
al., 2015). The level of cadmium in blood

and devices coated with metals (Malik and Malik, 2011). The accumulation of metals in the body of organisms with the passage of time. The higher concentration of cadmium in quail causes thinning, damages of kidney and calcium metabolism, reproductive system, and immune system (Burger, 2008). The concentration of metals in different species were founded in following order: quail > common myna > house sparrow. The levels of cadmium in quail blood and feathers were found to be different from sparrow and common myna. According to our results, cadmium was not significantly associated with hematological parameters. There was no significant negative relation between cadmium concentration in blood and white blood cells. The results of cadmium concentration in sparrow and common myna were found to be similar with the results of (Lester and Manjula et al., 2015; Muralidharan et al., 2015; Manjula et al., 2015). The level of cadmium in blood

Table 2. Results of one way ANOVA means and standard errors of hematological parameters.

Parameters	Species			ANOVA Sig.
	House Sparrow (<i>Passer domesticus</i>)	Common Myna (<i>Acridotheres tristis</i>)	Quail (<i>Coturnix coturnix</i>)	
RBCs ($10^6/\mu\text{l}$)	1.34 ± 0.06^b	1.87 ± 0.02^b	4.678 ± 0.70^a	0.000**
WBCs ($10^3/\mu\text{l}$)	14.79 ± 0.71^b	23.80 ± 0.71^a	21.52 ± 2.76^a	0.007**
MCV (fl)	168.64 ± 3.93^a	141.32 ± 1.37^b	128.54 ± 6.14^b	0.000**
MCH (pg)	68.62 ± 2.70^{ab}	56.67 ± 1.10^b	72.90 ± 5.66^a	0.023*
MCHC (g/dl)	40.93 ± 1.02^b	38.80 ± 1.20^b	64.29 ± 7.52^a	0.003**

Table 3. Results of differential leukocytes count in percentage and one way ANOVA.

Parameters	Species			ANOVA Sig.
	House Sparrow (<i>Passer domesticus</i>)	Common Myna (<i>Acridotheres tristis</i>)	Quail (<i>Coturnix coturnix</i>)	
Lymphocytes	41.41 ± 1.07^b	44.43 ± 2.34^b	63.46 ± 2.48^a	0.000**
Monocytes	1.40 ± 0.03^b	1.49 ± 0.02^b	2.63 ± 0.36^a	0.002**
Heterophils	38.51 ± 2.00^a	37.04 ± 1.70^a	22.50 ± 4.45^b	0.004**
Eosinophils	18.31 ± 1.77^a	16.92 ± 1.81^a	10.40 ± 3.88^a	0.126
Basophils	0.58 ± 0.01^b	0.54 ± 0.02^b	1.59 ± 0.38^a	0.009**

Different Alphabetic letters indicate significant differences in metal levels among species derived through post-hoc (tukey) test.

Table 4. Relation between concentration of metals in blood and hematological parameter.

Independents	Dependents				
	RBCs	WBCs	MCV	MCH	MCHC
Cu	0.440	0.846	0.999	0.578	0.593
Zn	0.282	0.011*	-0.058	-0.058	0.352
Ni	0.116	0.531	0.400	0.360	0.135
Cd	0.053	-0.621	-0.300	0.139	0.165
Cr	0.625	0.816	-0.231	-0.484	0.909
Pb	-0.384	0.006*	-0.629	-0.192	-0.202
ANOVA Sig.	0.059	0.009**	0.136	0.076	0.073

and feather samples of house sparrow and common myna were lower than the study of (Geens et al., 2010) reported from contaminated areas for great tits and (Malik and Zeb, 2009; Ullah et al., 2014) reported from Pakistan for cattle egret. The permissible range of cadmium for birds is $2\mu\text{g/g}$ (Burger and Gochfeld, 2000). The level of cadmium in quail feathers was slightly greater than the permissible range.

Chromium is a micro essential element and may cause hazardous effects on reproductive system (Malik and Zeb,

2009). The major source of chromium contamination in our study area is leather industry, which use various chromium salts as a tanning agent (Qadir et al., 2008). Various avian species secrete chromium salts through nasal glands, which can bind with the keratin of feathers. The highest accumulation of chromium was recorded in common myna during our findings. The common myna revealed significant variations with quail and house sparrow for the concentration of lead in feathers and blood. We did not find significant negative impact of

Table 5. Comparison of metals ($\mu\text{g/g}$) in feathers and blood between sparrow, common myna, and quail from previous and present study.

Authors	Species	Sample	Study area	Zn	Cu	Ni	Cd	Cr	Pb
world wide									
(Swaileh and Sansur, 2006)	House sparrow	Feathers	Palestine, Qalqilia, Ramallah city	54.9	19.5		0.02		8.1
(Manjula et al., 2015)	House sparrow	Feathers	India Tiruchirapalli (rural area)	75.91	21.04	8.13	1.71	55.4	
		Feathers	India Tiruchirapalli (urban area)	98.16	148.29	76.6	1.64	53.1	
(Bounagua et al., 2014)	House sparrow	Blood	Morocco (Abdullah <i>et al.</i>)	331.07			3.78		35.42
		Blood	Morocco (Town center)	326.12			5.63		40.12
		Blood	Morocco (Agdal)	212.74			3.88		32.03
		Blood	Morocco (Oulja)	284.6			8.34		33.72
		Blood	Morocco (Bahraoui)	403.7			3.74		2.7
(Manjula et al., 2015)	Common myna	Feathers	India Tiruchirapalli (rural area)	65.1	25.85	7.54	1.59	55.1	
		Feathers	India Tiruchirapalli (urban area)	70.65	80.32	8.51	1.7	85.7	
(Muralidharan et al., 2004)	Common myna	Primary feathers	India Nilgiris	137.98	1.86		1.89	0.58	2.2
		Secondary feathers		179.68	5.69		3.08	0.27	
Pakistan									
(Mustafa et al., 2015)	House sparrow	Feathers	Sargodha	14.78	0.14		1.81		2.67
(Abbasi et al., 2015)	Common myna	Feathers	Baroghil valley	60.83	1.04	1.73	0.35	0.6	0.95
			Soanvally	91.05	2.19	1.88	0.78	1.11	2.15

chromium pollution on hematological parameters, but the results revealed some negative association between chromium contamination with mean cell volume and mean cell hemoglobin constantans. The concentration of chromium in our feathers and blood samples were greater than the findings of (Lester and van Riper, 2014) reported for song sparrow from Arizona and of (Muralidharan et al., 2004) reported for common myna from India. Our results revealed slight similarities with the estimations of (Ullah et al., 2014) reported for cattle egret from Pakistan. According to (Burger and Gochfeld, 2000) the permissible range of chromium in birds is $2.8\mu\text{g/g}$. So, the concentrations of chromium in all studied species were greater than safe level, which can exert negative impact on the health status of avian species.

Lead is a highly toxic element; the major sources of lead in environment are vehicles, burning fossil fuel, paints, furniture polishing, and tanning industries. The lead is not

required for any metabolic process, and lead showed great affinity to accumulate in feathers and calcareous tissues (Jerez et al., 2011; Markowski et al., 2013). According to the study of Burger and Gochfeld, (2000) the level of lead more than $4\mu\text{g/g}$ leads to the abnormalities in thermoregulation, locomotion reproductive system, and feeding behavior. The threshold value of lead in blood is $2\mu\text{g/g}$ (Geens et al., 2010). The accumulation of lead in feathers and blood samples were founded in the following descending order common myna > house sparrow and quail. There were significant differences occurred between different species for lead accumulation. There was no significant relation founded between lead concentration and hematological parameters, but we find some negative relations with hematological parameters except white blood cells. The increased level of lead results in decreases in hematological parameters. The lead concentrations in other studies reported from Pakistan and other parts of the world for

Table 6. Comparison of metals ($\mu\text{g/g}$) in feathers and blood of different species with present study.

Author	Species	Sample	Study area	Zn	Cu	Ni	Cd	Cr	Pb
world wide									
(Geens et al., 2010)	Great tit	Feathers	South Antwerp (UM)	240	37	13	11		140
			South Antwerp (F8)	216	36.5	14	17		110
			South Antwerp (F7)	142	9.8	7.2	0.39		8.1
			South Antwerp (F4)	184	9.8	5.7	0.19		4.1
		Blood	South Antwerp (UM)	5.5	0.13	0.11	0.016		0.28
			South Antwerp (F8)	6.9	0.09	0.14	0.011		0.17
			South Antwerp (F7)	7.9	0.2	0.13	0.007		0.03
			South Antwerp (F4)	6.9	0.13	0.18	0.007		0.02
(Lester and van Riper, 2014)	Song sparrow	Blood	Arizona (LOCH)	195.1	11.69	1.22	0.109	0.418	0.706
			NOWA	9.606	0.811	0.033	0.005	0.047	0.031
			PLSP	6.425	1.118	0.043	0.004	0.023	0.039
			SOCR	6.831	0.578	0.035	0.002	2.145	0.031
			NIWWTP	7.509	0.699	0.083	0.004	0.12	0.108
			TUMA	8.219	0.74	0.025	0.004	0.018	0.027
		Feathers	LOCH	7.132	0.672	0.136	0.002	0.035	0.041
			NOWA	215.6	15.35	1.144	0.362	0.431	1.191
			PLSP	196.3	11.4	0.752	0.366	0.238	2.283
			SOCR	189	10.86	0.736	0.127	0.387	2.273
			NIWWTP	206.3	11.87	3.457	1.022	0.763	1.069
			TUMA	171.6	10.73	1.323	0.422	0.364	1.05
Pakistan									
(Malik and Zeb, 2009)	Cattle egret	feathers	chenab	133.8	4	9.1	3.1	6.6	37.5
			ravi river	155.2	3.7	8	2.4	7.1	76.5
			raval lake	138.4	4	7.8	2.7	5.38	60.2
Ullah et al., s 2014)	Cattle egret	feathers	Trimium headwork	18.5	0.9	0.1	3.3	26.2	30
			Shorkot	18.8	0.2	0.2	3	30.8	32.5
			Mailsi	10.7	0.2	0.2	1.7	35.8	43.1

house sparrow feathers (Mustafa et al., 2015; Swaileh and Sansur, 2006), and for common myna feathers (Abbasi et al., 2015; Muralidharan et al., 2004) were lower than our measured concentrations. Malik and Zeb, 2009 and Ullah et al., 2014 reported higher concentrations of lead than our results in cattle egret feathers from Pakistan. The results for lead in blood were slightly higher in our studied species than the findings of Lester and van Riper, 2014. The level of lead in common myna and house sparrow feathers and blood was not at safe level according to the threshold limits of Burger and Gochfeld, 2000; Geens et al., 2010.

The feathers accumulate highest concentration of metals than blood; the most probable reason behind that is the feathers reflect exogenous contamination, and blood only represent the immediate dietary sources

of contamination (Veerle et al., 2004). So, the feathers can be used as an appropriate approach to find out the environmental contamination level. The major reason of differences between different species was difference in feeding habitat (Abbasi et al., 2015). The house sparrow and common myna are omnivores, but house sparrow mostly feed on grains. The quail is less mobile and feed 95% on plants so the metal contamination was founded at fewer rates in quail bird. So, the metal contamination was founded in the following descending order common myna> house sparrow> quail. The comparison of our study with other studies at Pakistan and international level are given in (Table 5, 6).

The hematological parameters were significantly varied between species. According to our results there

were no significant or non-significant negative association observed between copper and Zinc concentrations in blood and hematological parameters. The results of (Samanta et al., 2011) for broiler chicken revealed that dietary copper more than 150mg/Kg negatively reduce the concentration of hemoglobin. However, there were no effect of copper were observed on total leukocyte count and different leukocyte count. The study of (Aksu et al., 2010) concluded that there are no negative effects of copper so the appropriate concentration of copper from organic sources can be used in the diet of broilers. According to (Dönmez et al., 2002) the level of hemoglobin, hematocrit and erythrocyte, leukocyte is not affected by exposure of zinc in broiler chicks. (Gupta et al., 1985) reported that the leukocyte count increase in rats by supplementation of zinc in diet. However, Donmez and Keskin, 1999 report no significant positive or negative effects of zinc on leukocyte count in rats and goat. Our results concluded that the cadmium was negatively correlated with MCV and WBCs; however, the numbers of RBCs, MCH, and MCHC were increased by increasing cadmium level. The results of (Gabol et al., 2014) can be compared with our study for the impact of cadmium on hematological parameters. The study of (Ali et al., 2020; Guttyj et al., 2019) represent opposite results with our research which concluded a negative relation of cadmium exposure with HGB, MCHC and RBCs while show positive relation with WBCs. The dietary sources of chromium may have positive relation with hematological parameters in Japanese quails which reared under heat stress conditions (El-Kholy et al., 2017). Nickel does not show any significant association with hematological parameters. The results revealed that the chromium and lead were negatively correlated with hematological parameters and chromium and lead were not founded at safe level, which can exert negative impact on health status of birds. The highest level of chromium and lead were observed in common myna, so hematological parameters were most affected in common myna as compared to house sparrow and quail. However, the level of white blood cells was significantly increased by increasing heavy metals in blood. The results for differential leukocyte count showed significant variations among different species except eosinophils. The increased level of chromium causes decreased numbers of MCH and MCV. The chromium is positively associated with the concentration of red blood cells and mean cell hemoglobin concentration. Some other studies also revealed positive association between chromium and red blood cells such as (Cui et al., 2005; Gabol et al., 2014). We founded that

by increasing lead concentration, the number of RBCs, MCV, MCH and MCHC decreased, but the numbers of WBCs increase. Various previous studies (Gabol et al., 2014; Geens et al., 2010) founded the negative impact of lead on hematological status of avian species. The accumulation of lead causes anemia in birds by breaking down of blood cells (Katavolos et al., 2007). The lead damage red blood cells and causes break down of plasma membrane (Leggett, 1993). ALAD activity, hematocrits level, and hemoglobin concentration decreased by increasing lead concentration (Cid et al., 2018). Several changes in blood cells such as erythrocytes agglutination, nucleus displacement in erythrocytes, absence of nucleus in erythrocytes, lymphocytes enlargement, immature erythrocytes, rotation of nucleus, large macrophage, large size lymphocytes, and aggregation of thrombocytes were examined in Japanese quails, which were exposed to lead. The numbers of healthy erythrocytes, lymphocytes, and heterophils were decreased, and there were no significant changes observed in numbers of leucocytes by exposure of lead chloride (Suljević et al., 2020). The study of Rogival et al., 2006 revealed negative effects of lead and cadmium exposure on hematological parameters in wood mice. The study of Li et al., 2021 reported that the air pollutants are negatively associated with hematological parameters.

According to our estimations among three species common myna were most affected by metal contamination. The findings of our research were concluded that the feathers of avian species can be used as an efficient bio-indicator to assess the metal contamination level in an environment; however, feathers did not revealed the internal contamination level. The results revealed that the hematological parameters were affected by increasing metal contents, especially lead and chromium. Therefore, the assessment of hematological parameters can be used to find out the negative impact of metal contamination on health status. We concluded from our research that the level of chromium and lead were not at a safe level, and they have exert negative impact on health status in avian fauna of Sialkot. The findings of this study will be helpful in environmental management and to understand the environmental status of study area.

Acknowledgments

The authors would like to acknowledge the department of Wildlife and Ecology University of veterinary and animal sciences, Lahore, Pakistan for providing assistance in lab work and Mr. Haris nadeem for providing GIS maps.

References

- Abbasi NA, Jaspers VLB, Chaudhry MJ, Ali S, Malik RN et al. (2015). Influence of taxa, trophic level, and location on bioaccumulation of toxic metals in bird's feathers: a preliminary biomonitoring study using multiple bird species from Pakistan. *Chemosphere* 120: 527-537.
- Abdullah M, Fasola M, Muhammad A, Malik SA, Bostan N et al. (2015). Avian feathers as a non-destructive bio-monitoring tool of trace metals signatures: a case study from severely contaminated areas. *Chemosphere* 119: 553-561.
- Agarwal S (2002) *Pollution Management: Vol-IV: Heavy Metal Pollution*. APH Publishing Corporation.
- Aksu DS, Aksu T, Ozsoy B (2010). The effects of lower supplementation levels of organically complexed minerals (zinc, copper and manganese) versus inorganic forms on hematological and biochemical parameters in broilers. *Kafkas Üniversitesi Veteriner Fakültesi Dergisi* 16 (4): 553-559.
- Ali S, Bashir S, Mumtaz S, Shakir HA, Ara C et al. (2020). Evaluation of Cadmium Chloride-Induced Toxicity in Chicks Via Hematological, Biochemical Parameters, and Cadmium Level in Tissues. *Biological trace element research* 1-13.
- Ayyamperumal T, Jonathan M, Srinivasalu S, Armstrong-Altrin J, Ram-Mohan V et al. (2006). Assessment of acid leachable trace metals in sediment cores from River Uppanar, Cuddalore, Southeast coast of India. *Environmental Pollution* 143 (1): 34-45.
- Baluch U (1995) *Pesticide monitoring programme*. WWF Pakistan. In. Lahore, Mimeo
- Boncompagni E, Muhammad A, Jabeen R, Orvini E, Gandini C et al. (2003). Egrets as monitors of trace-metal contamination in wetlands of Pakistan. *Archives of Environmental Contamination and Toxicology* 45 (3): 399-406.
- Booth CE, Elliott PF (2002). Hematological responses to hematozoa in North American and neotropical songbirds. *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology* 133 (3): 451-467.
- Bounagua M, Bellaouchou A, Benabbou A, El Abidi A, Ben-aakam R et al. (2014). Using bloods *Passer domesticus* as a possible bio-indicator of urban heavy metals pollution in Rabat-Salé (Morocco). *Journal of Materials and Environmental Science* 5 (3): 937-944.
- Brait CHH, Antoniosi Filho NR (2011). Use of feathers of feral pigeons (*Columba livia*) as a technique for metal quantification and environmental monitoring. *Environmental monitoring and assessment* 179 (1-4): 457-467.
- Burger J (2008). Assessment and management of risk to wildlife from cadmium. *Science of the total environment* 389 (1): 37-45.
- Burger J (2013). Temporal trends (1989–2011) in levels of mercury and other heavy metals in feathers of fledgling great egrets nesting in Barnegat Bay, NJ. *Environmental research* 122: 11-17.
- Burger J, Gochfeld M (2000). Metal levels in feathers of 12 species of seabirds from Midway Atoll in the northern Pacific Ocean. *Science of the Total Environment* 257 (1): 37-52.
- Burger J, Gochfeld M (2007). Metals and radionuclides in birds and eggs from Amchitka and Kiska Islands in the Bering Sea/Pacific Ocean ecosystem. *Environmental monitoring and assessment* 127 (1-3): 105-117.
- Chen R, Wei L, Huang H (1993). Mortality from lung cancer among copper miners. *Occupational and Environmental Medicine* 50 (6): 505-509.
- Cheney MA, Hacker CS, Schroder GD (1981). Bioaccumulation of lead and cadmium in the Louisiana heron (*Hydranassa tricolor*) and the cattle egret (*Bubulcus ibis*). *Ecotoxicology and Environmental Safety* 5 (2): 211-224.
- Cid FD, Fernández NC, Pérez-Chaca MV, Pardo R, Caviedes-Vidal E, Chediack JG (2018). House sparrow biomarkers as lead pollution bioindicators. Evaluation of dose and exposition length on hematological and oxidative stress parameters. *Ecotoxicology and environmental safety* 154: 154-161.
- Cui H, Yang G, Peng X (2005). Effect of Copper Toxicity on Blood Biochemical Parameters in Broilers. *Acta Veterinaria Et Zootechnica Sinica* 36 (12): 1329.
- Dauwe T, Janssens E, Kempenaers B, Eens M (2004). The effect of heavy metal exposure on egg size, eggshell thickness and the number of spermatozoa in blue tit *Parus caeruleus* eggs. *Environmental Pollution* 129 (1): 125-129.
- Deng H, Zhang Z, Chang C, Wang Y (2007). Trace metal concentration in great tit (*Parus major*) and greenfinch (*Carduelis sinica*) at the Western Mountains of Beijing, China. *Environmental Pollution* 148 (2): 620-626.
- Dönmez N, Dönmez HH, Keskin E, Çelik I (2002). Effects of zinc supplementation to ration on some hematological parameters in broiler chicks. *Biological trace element research* 87 (1): 125-131.
- Donmez N, Keskin E (1999). Ankara Kecilerinde Rasyona Cinko İlavlesinin Bazi Hematolojik Parametreler Ozerine Etkisi. *Veteriner Bilimler Dergisi* 15:125-131.
- Eisler R (1988) *Lead hazards to fish, wildlife, and invertebrates: a synoptic review*. US Fish and Wildlife Service, Patuxent Wildlife Research Center.
- El-Kholy MS, El-Hindawy MM, Alagawany M, Abd El-Hack ME, Abd El SAE-G et al. (2017). Dietary supplementation of chromium can alleviate negative impacts of heat stress on performance, carcass yield, and some blood hematology and chemistry indices of growing Japanese quail. *Biological trace element research* 179 (1): 148-157.
- Eqani SA-M-A-S, Malik RN, Katsoyiannis A, Zhang G, Chakraborty P et al. (2012). Distribution and risk assessment of organochlorine contaminants in surface water from River Chenab, Pakistan. *Journal of Environmental Monitoring* 14 (6): 1645-1654.

- Falq G, Zeghnoun A, Pascal M, Vernay M, Le Strat Y et al. (2011). Blood lead levels in the adult population living in France the French Nutrition and Health Survey (ENNS 2006–2007). *Environment international* 37 (3): 565-571.
- Frantz A, Pottier M-A, Karimi B, Corbel H, Aubry E et al. (2012). Contrasting levels of heavy metals in the feathers of urban pigeons from close habitats suggest limited movements at a restricted scale. *Environmental Pollution* 168 : 23-28.
- Furness RW (1993) Birds as monitors of pollutants. In: *Birds as monitors of environmental change*. Springer, pp 86-143.
- Gabol K, Khan MZ, Khan MUA, Khan P, Fatima F et al. (2014). Induced effects of lead, chromium and cadmium on gallus domesticus. *Canadian Journal of Pure and Applied Sciences* 8 (3): 3035-3042.
- Geens A, Dauwe T, Bervoets L, Blust R, Eens M et al. (2010). Haematological status of wintering great tits (*Parus major*) along a metal pollution gradient. *Science of the Total Environment* 408 (5): 1174-1179.
- Gupta R, Verma P, Gupta RP (1985). Experimental zinc deficiency in guinea-pigs: clinical signs and some haematological studies. *British journal of nutrition* 54 (2): 421-428.
- Gushit J, Turshak L, Chaskda A, Abba B, Nwaeze U et al. (2016). Avian Feathers as Bioindicator of Heavy Metal Pollution in urban degraded woodland.
- Gutyj B, Ostapyyuk A, Sobolev O, Vishchur VJ, Gubash O, Kurtyak B, Kovalskiy Y, Darmohray L, Hunchak A, Tsisaryk O (2019). Cadmium burden impact on morphological and biochemical blood indicators of poultry. *Ukrainian Journal of Ecology* 9: 1.
- Hardoy JE, Mitlin D, Satterthwaite D (1992) *Environmental problems in Third World cities*. London: Earthscan.
- Hashmi MZ, Malik RN, Shahbaz M (2013). Heavy metals in eggshells of cattle egret (*Bubulcus ibis*) and little egret (*Egretta garzetta*) from the Punjab province, Pakistan. *Ecotoxicology and Environmental safety* 89:158-165.
- Herrera-Duenas A, Pineda J, Antonio MT, Aguirre JI (2014). Oxidative stress of house sparrow as bioindicator of urban pollution. *Ecological Indicators* 42: 6-9.
- Honda K, Min BY, Tatsukawa R (1986). Distribution of heavy metals and their age-related changes in the eastern great white egret, *Egretta alba modesta*, in Korea. *Archives of Environmental Contamination and Toxicology* 15 (2): 185-197.
- Iolascon A, De Falco L, Beaumont C (2009). Molecular basis of inherited microcytic anemia due to defects in iron acquisition or heme synthesis. *haematologica* 94 (3): 395-408.
- Iqbal M, Haque I, Berns J (1998). *The Leather Sector, Environmental Report*. Environmental Technology Programme For Industry (ETPI), Federation of Pakistan Chambers of Commerce and Industry, Federation House, Karachi, Pakistan.
- Jaspers V, Covaci A, Maervoet J, Dauwe T, Voorspoels S et al. (2005). Brominated flame retardants and organochlorine pollutants in eggs of little owls (*Athene noctua*) from Belgium. *Environmental Pollution* 136 (1): 81-88.
- Jerez S, Motas M, Palacios MJ, Valera F, Cuervo JJ et al. (2011). Concentration of trace elements in feathers of three Antarctic penguins: geographical and interspecific differences. *Environmental Pollution* 159 (10): 2412-2419.
- Kalisińska E, Salicki W, Myslek P, Kavetska KM, Jackowski A et al. (2004). Using the mallard to biomonitor heavy metal contamination of wetlands in North-Western Poland. *Science of the Total Environment* 320 (2-3): 145-161.
- Katavolos P, Staempfli S, Sears W, Gancz A, Smith D et al. (2007). The effect of lead poisoning on hematologic and biochemical values in trumpeter swans and Canada geese. *Veterinary clinical pathology* 36 (4): 341-347.
- Kenntner N, Krone O, Altenkamp R, Tataruch F (2003). Environmental contaminants in liver and kidney of free-ranging northern goshawks (*Accipiter gentilis*) from three regions of Germany. *Archives of Environmental Contamination and Toxicology* 45 (1): 0128-0135.
- Khan FK (1991) *A geography of Pakistan: environment, people and economy*. Oxford University Press.
- Kilgas P, Tilgar V, Mänd R (2006). Hematological health state indices predict local survival in a small passerine bird, the great tit (*Parus major*). *Physiological and Biochemical Zoology* 79 (3): 565-572.
- Leggett RW (1993). An age-specific kinetic model of lead metabolism in humans. *Environmental Health Perspectives* 101 (7): 598-616.
- Lester MB, van Riper C (2014). The distribution and extent of heavy metal accumulation in song sparrows along Arizona's upper Santa Cruz River. *Environmental monitoring and assessment* 186 (8): 4779-4791.
- Li M, Nabi G, Sun Y, Wang Y, Wang L et al. (2021). The effect of air pollution on immunological, antioxidative and hematological parameters, and body condition of Eurasian tree sparrows. *Ecotoxicology and environmental safety* 208:111755.
- Llabjani V, Malik RN, Trevisan J, Hoti V, Ukpebor J et al. (2012). Alterations in the infrared spectral signature of avian feathers reflect potential chemical exposure: A pilot study comparing two sites in Pakistan. *Environment international* 48: 39-46.
- Malik RN, Jadoon WA, Husain SZ (2010). Metal contamination of surface soils of industrial city Sialkot, Pakistan: a multivariate and GIS approach. *Environmental geochemistry and health* 32 (3): 179-191.
- Malik RN, Zeb N (2009). Assessment of environmental contamination using feathers of *Bubulcus ibis* L., as a biomonitor of heavy metal pollution, Pakistan. *Ecotoxicology* 18 (5): 522-536.
- Manjula M, Mohanraj R, Devi MP (2015). Biomonitoring of heavy metals in feathers of eleven common bird species in urban and rural environments of Tiruchirappalli, India. *Environmental monitoring and assessment* 187 (5): 267.
- Markowski M, Kaliński A, Skwarska J, Wawrzyniak J, Bańbura M et al. (2013). Avian feathers as bioindicators of the exposure to heavy metal contamination of food. *Bulletin of environmental contamination and toxicology* 91 (3): 302-305.

- Martin MB, Reiter R, Pham T, Avellanet YR, Camara J et al. (2003). Estrogen-like activity of metals in MCF-7 breast cancer cells. *Endocrinology* 144 (6): 2425-2436.
- Millaku L, Imeri R, Trebicka A (2000). Histopathological changes in testes of house sparrow (*Passer domesticus*).
- Muralidharan S, Jayakumar R, Vishnu G (2004). Heavy metals in feathers of six species of birds in the district Nilgiris, India. *Bulletin of environmental contamination and toxicology* 73 (2): 285-291.
- Mustafa I, Ghani A, Arif N, Asif S, Khan MR et al. (2015). Comparative metal profiles in different organs of house sparrow (*Passer domesticus*) and black kite (*Milvus migrans*) in Sargodha District, Punjab, Pakistan. *Pakistan Journal of Zoology* 47 (4).
- Naccari C, Cristani M, Cimino F, Arcoraci T, Trombetta D et al. (2009). Common buzzards (*Buteo buteo*) bio-indicators of heavy metals pollution in Sicily (Italy). *Environment International* 35 (3): 594-598.
- Nadolski J, Skwarska J, Kaliński A, Bańbura M, Śniegula R et al. (2006). Blood parameters as consistent predictors of nestling performance in great tits (*Parus major*) in the wild. *Comparative Biochemistry and Physiology Part A: Molecular and Integrative Physiology* 143 (1): 50-54.
- Ots I, Murumägi A, Horak P (1998). Haematological health state indices of reproducing great tits: methodology and sources of natural variation. *Functional Ecology* 12 (4): 700-707.
- Outridge P, Scheuhammer A (1993). Bioaccumulation and toxicology of nickel: implications for wild mammals and birds. *Environment Reviews* 1 (2): 172-197.
- Papanikolaou NC, Hatzidaki EG, Belivanis S, Tzanakakis GN, Tsatsakis AM (2005). Lead toxicity update. A brief review. *Medical science monitor* 11 (10): 329-336.
- Peakall D (1992). Biomarkers of the nervous system. *Animal Biomarkers as Pollution Indicators*. In: New York: Springer.
- Peplow D (2000). Environmental impacts of hard-rock mining in Eastern Washington. University of Washington Fact Sheet 8.
- Pérez-López M, de Mendoza MH, Beceiro AL, Rodríguez FS (2008). Heavy metal (Cd, Pb, Zn) and metalloid (As) content in raptor species from Galicia (NW Spain). *Ecotoxicology and environmental safety* 70 (1): 154-162.
- Qadir A, Malik RN (2011). Heavy metals in eight edible fish species from two polluted tributaries (Aik and Palkhu) of the River Chenab, Pakistan. *Biological Trace Element Research* 143 (3): 1524-1540.
- Qadir A, Malik RN, Husain SZ (2008). Spatio-temporal variations in water quality of Nullah Aik-tributary of the river Chenab, Pakistan. *Biological Trace Element Research* 140 (1-3): 43-59.
- Raihan S, Sarwar F, Azim M, Khan O, Ahmed N et al. (1995). Isolation, characterization and assessment of nickel and cadmium accumulation of bacterial isolates from industrial waste. *Biotechnology for Environment and Agriculture BCC and T Press, Univ of Karachi* 143-152.
- Rogival D, Scheirs J, De Coen W, Verhagen R, Blust R (2006). Metal blood levels and hematological characteristics in wood mice (*Apodemus sylvaticus* L.) along a metal pollution gradient. *Environmental Toxicology and Chemistry: An International Journal* 25 (1): 149-157.
- Saino N, Cuervo JJ, Krivacek M, de Lope F, Møller AP (1997). Experimental manipulation of tail ornament size affects the hematocrit of male barn swallows (*Hirundo rustica*). *Oecologia* 110 (2): 186-190.
- Samanta B, Ghosh P, Biswas A, Das S (2011). The effects of copper supplementation on the performance and hematological parameters of broiler chickens. *Asian-Australasian Journal of Animal Sciences* 24 (7): 1001-1006.
- Scheifler R, Coeurdassier M, Morilhat C, Bernard N, Faivre B et al. (2006). Lead concentrations in feathers and blood of common blackbirds (*Turdus merula*) and in earthworms inhabiting unpolluted and moderately polluted urban areas. *Science of the Total Environment* 371 (3): 197-205.
- Scope A, Frommlet F, Schwendenwein I (2005). Circadian and seasonal variability and influence of sex and race on eight clinical chemistry parameters in budgerigars (*Melopsittacus undulatus*, Shaw 1805). *Research in veterinary science* 78 (1): 85-91.
- Spahn S, Sherry T (1999). Cadmium and lead exposure associated with reduced growth rates, poorer fledging success of little blue heron chicks (*Egretta caerulea*) in south Louisiana wetlands. *Archives of Environmental Contamination and Toxicology* 37 (3): 377-384.
- Suljević D, Handžić N, Fočak M, Lasić I, Sipović F et al. (2020). Lead Exposure Influences Serum Biomarkers, Hepatocyte Survival, Bone Marrow Hematopoiesis, and the Reproductive Cycle in Japanese Quails. *Biological Trace Element Research* 199 (4): 1574-1583.
- Svensson E, Merilä J (1996). Molt and migratory condition in blue tits: a serological study. *The Condor* 98 (4): 825-831.
- Swaileh K, Sansur R (2006). Monitoring urban heavy metal pollution using the House Sparrow (*Passer domesticus*). *Journal of Environmental Monitoring* 8 (1): 209-213.
- Syed JH, Malik RN (2011). Occurrence and source identification of organochlorine pesticides in the surrounding surface soils of the Ittehad Chemical Industries Kalashah Kaku, Pakistan. *Environmental Earth Sciences* 62 (6): 1311-1321.
- Taggart MA, Green AJ, Mateo R, Svanberg F, Hillström L et al. (2009). Metal levels in the bones and livers of globally threatened marbled teal and white-headed duck from El Hondo, Spain. *Ecotoxicology and Environmental Safety* 72 (1): 1-9.
- Tariq J, Jaffar M, Ashraf M (1994). Trace metal concentration, distribution and correlation in water, sediment and fish from the Ravi River, Pakistan. *Fisheries Research* 19 (1-2): 131-139.
- Tehseen W, Hansen L, Wood S, Hanif M (1994). Assessment of chemical contaminants in water and sediment samples from Degh Nala in the province of Punjab, Pakistan. *Archives of Environmental Contamination and Toxicology* 26 (1): 79-89.

- Tsipoura N, Burger J, Feltes R, Yacabucci J, Mizrahi D et al. (2008). Metal concentrations in three species of passerine birds breeding in the Hackensack Meadowlands of New Jersey. *Environmental research* 107 (2): 218-228.
- Ullah K, Hashmi MZ, Malik RN (2014). Heavy-metal levels in feathers of cattle egret and their surrounding environment: a case of the Punjab Province, Pakistan. *Archives of environmental contamination and toxicology* 66 (1): 139-153.
- Van den Steen E, Dauwe T, Covaci A, Jaspers V, Pinxten R et al. (2006). Within-and among-clutch variation of organohalogenated contaminants in eggs of great tits (*Parus major*). *Environmental Pollution* 144 (1): 355-359.
- Veerle J, Tom D, Rianne P, Lieven B, Ronny B et al. (2004). The importance of exogenous contamination on heavy metal levels in bird feathers. A field experiment with free-living great tits, *Parus major*. *Journal of Environmental Monitoring* 6 (4): 356-360.
- Xu L, Wang T, Ni K, Liu S, Wang P et al. (2013). Metals contamination along the watershed and estuarine areas of southern Bohai Sea, China. *Marine pollution bulletin* 74 (1): 453-463.
- Zaman A, Ara I (2000). Rising urbanization in Pakistan, some facts and suggestions. *The journal NIPA* 7: 31-46.