

1 **Leukocyte esterase reagent strips: Rapid, economical and reliable cow side test for**
2 **subclinical endometritis**

3 **Arshaq ASFAR^{1,*}, Khursheed Ahmad SOFI², Mudasir Ahmad BHAT³, Abrar**
4 **Ahad MALIK⁴, Asloob Ahmad MALIK¹**

5 ¹Division of Animal Reproduction Gynaecology and Obstetrics, FVSc & AH, Sher-e-
6 Kashmir University of Agricultural Sciences and Technology of Kashmir, J&K, India

7 ²Division of Veterinary Clinical Complex, FVSc & AH, Sher-e-Kashmir University of
8 Agricultural Sciences and Technology of Kashmir, J&K, India

9 ³Division of Animal Nutrition, FVSc & AH, Sher-e-Kashmir University of Agricultural
10 Sciences and Technology of Kashmir, J&K, India

11 ⁴Division of Animal Biotechnology, FVSc & AH, Sher-e-Kashmir University of
12 Agricultural Sciences and Technology of Kashmir, J&K, India

13 *Corresponding author: arshaqasfar@gmail.com; +917006357052, +91901827769

14 ORCID's:

15 1- Arshaq Asfar: [0000-0002-6587-7773](https://orcid.org/0000-0002-6587-7773)

16 2- Khursheed Ahmad Sofi: 0000-0002-0794-7008

17 3- Mudasir Ahmad Bhat: 0000-0003-2422-2874

18 4- Abrar Ahad Malik: 0000-0003-0089-7332

19 5- Asloob Ahmad Malik: 0000-0002-9006-0482

20
21 **Abstract**

22 This study was carried out in crossbred cows (>60 DIM) to evaluate leukocyte esterase
23 (LE) strip test, protein strip test, pH strip test, Whiteside test (WST) of estrual mucus, in
24 comparison to endometrial cytology (EC) using cytobrush (CB) as reference diagnostic
25 method for subclinical endometritis (SCE). Endometrial Cytology was done in 78 cows
26 with $\geq 4\%$ PMNs as cut-off value for SCE. Leukocyte Esterase strip test was done by
27 contacting estrual mucus with multi-reagent urinalysis reagent strips and results recorded

1 after 10 minutes. Similarly, protein and pH strip tests were conducted using the same
2 strips and results recorded after 60 seconds. Whiteside test of estrual mucus was
3 conducted simultaneously. Leukocyte Esterase, protein and pH strip tests were evaluated
4 and analyzed statistically using Receiver Operator Characteristic (ROC). LE strip test at
5 cut-off value $\geq \pm$ acquired from ROC analysis showed sensitivity (Se) of 86.67% and
6 specificity (Sp) of 66.67%. Similarly, protein and pH strip tests showed Se of 53.33% &
7 66.67% and Sp of 66.67% & 77.78% respectively. Strip tests were also combined in three
8 combinations (LE + protein, LE + pH and LE + protein + pH) and overall decrease in Se
9 was observed in all three combinations (46.67%, 60% and 33.33%, respectively); while,
10 a slight increase in Sp (77.78%) was also observed. Sensitivity and Sp of WST were
11 63.33% and 55.56%; respectively, and these values for WST were lower compared to
12 those in LE test. In conclusion, cow side LE test can be used as an economic and easy test
13 with acceptable Se and Sp compared to WST instead of EC for SCE in cattle. However,
14 no improvement in performance of LE strip test was observed when combined with pH
15 and protein strip tests.

1 **Key words:** Subclinical endometritis, endometrial cytology, leukocyte esterase strip test

2 **1. Introduction**

3 The foundation for optimum fertility and high conception rates in dairy cows
4 require healthy uterine milieu [1]. Corpus luteal function, decreased sperm motility and
5 reduced embryonic viability are the impairments of blooming reproduction at various
6 steps, leading to decline in pregnancy rate and occur due to inflammatory atmosphere
7 being present in-utero [2]. In this aspect, subclinical endometritis (SCE) is important as
8 it produces negative outcome of reproduction and considerable economic sufferings by
9 being the chief cause of repeat breeding and all this by being asymptomatic as well as
10 immensely prevalent disease in cows [3, 4].

11 SCE is known to occur when the uterine interior coating (endometrium) is
12 inflamed. Term cytological endometritis has been used for SCE due to association with
13 elevated fraction (proportion) of poly-morpho-nuclear cells (PMN) in samples of
14 endometrial cytology (EC) acquired by cytobrush [5] or small-volume uterine lavage or
15 from biopsy samples by histology [4, 6]. Also bacteriology and ultrasonography [7, 8]
16 have been engaged to diagnose it. Although cytobrush (CB) technique is presently being
17 considered the reference and most efficient method for cytological identification of SCE
18 [9] but the practical applicability of this method for field situations is not realistic due to
19 the reasons that this method is time consuming and most importantly demands technical
20 expertise for collecting in-utero samples and then to stain as well as to identify individual
21 cells in samples.

22 Leukocyte esterase (LE) reagent test using LE coated strips is an alternative mode
23 to detect inflammatory cells in the diseased uterus [10]. The leukocyte esterase activity
24 of neutrophils' is mainly evaluated by this test and the number of leukocytes is related by
25 the intensity of color change [11]. Diagnoses of UTIs in humans is usually being done by

1 this test but recently have been used for diagnosis of inflammation in the uterus using
2 cervico-vaginal mucus [12] and projected as an alternative, innovative and creative cow-
3 side test for identification of SCE [10].

4 A practical cow-side and rapid clinical diagnostic test to inspect and manage SCE
5 under field conditions is very much desired. Therefore, the objective of this study was to
6 determine the efficacy of various diagnostic tests (LE, protein, pH, or combination of
7 reagent strip tests and Whiteside test (WST) in comparison to reference test (Endometrial
8 cytology) in search of a potential cow side test for SCE.

9 **2. Materials and Methods**

10 This investigation was conducted at Veterinary Clinical Complex (VCC), FVSc
11 & AH, Shuhama, Srinagar and Mountain Livestock Research Institute (MLRI),
12 Manasbal, Ganderbal. Crossbred postpartum cows with more than 60 days in milk (DIM)
13 presented for artificial insemination (AI) at VCC and MLRI were initially enrolled in the
14 study. Investigation was then conducted in 78 clinically healthy cows with no apparent
15 clinical signs of any disease, clear estrual mucus discharge and without any gross
16 reproductive tract pathology detected per-rectally. In the present study, diagnosis of SCE
17 was done by: 1-Endometrial cytology as reference method using cytobrush, 2-Leukocyte
18 esterase strip test, 3-Protein and pH strip test and 4-White side test.

19 **2.1 Endometrial cytology**

20 Uterine cytology samples were collected by cytobrush (Care India Surgical,
21 Mayfair Surgical Corporation, India) using cytobrush assembly as described by Singh et
22 al. [13] with slight modification. Base of the CB was fixed on the stylette and retracted
23 into the steel catheter. The CB assembly was then covered by sanitary sheath (IMV
24 Technologies, India) for protection against vaginal and cervical contamination. Before
25 sampling, perineum including vulva was washed with water and soap and dried with soft

1 absorbent cotton. The introduction of CB assembly into the uterine body through cervix
2 and vagina was achieved while maintaining sleeved arm per-rectally for cervical
3 manipulation. CB progressed out of steel catheter by puncturing the sanitary sheath after
4 crossing the cervical internal Os, approx 1-2 cm into uterine body. CB was rotated gently
5 once and then again in opposite direction while being in contact with adjacent
6 endometrium for cellular material collection. The whole apparatus was then removed
7 from genital tract after retracting the CB into the steel tube.

8 Slides were prepared immediately by rolling the CB on glass slides. Slides were
9 air dried, fixed in methanol (Merck Specialities Pvt Ltd, India) for 5-10 minutes and after
10 drying, staining was done with Giemsa stain (HiMedia Laboratories Pvt Ltd, India).
11 Cytologic evaluation was made by light microscopy at magnification of 1000x. Cells were
12 counted up-to 300 (endometrial cells and PMNs) and average percentage of PMN cells
13 (Neutrophils) was calculated and $\geq 4\%$ of PMNs was taken as threshold level for positive
14 SCE as per Singh et al. [13].

15 **2.2 Leukocyte esterase reagent strip test**

16 In LE strip test, about one ml of collected estrual mucus was contacted with multi-
17 reagent LE test strip (Mission Urinalysis Reagent Strips ACON Laboratories, Inc., U.S.A)
18 for 10-15 seconds. The results for LE strip test were then recorded based on the strip
19 colour change (according to the manufacturer recommendations) after 10 minutes
20 according to Hajibemani et al. [12].

21 **2.3 Protein strip test**

22 About one ml of collected EM was contacted with multi-reagent urinalysis reagent
23 strip (Mission Urinalysis Reagent Strips ACON Laboratories, Inc., U.S.A) for 10-15
24 seconds. Protein (mainly albumin) content of EM was recorded based on colour change
25 after 60 seconds (according to the manufacturer recommendations).

1 **2.4 pH strip test**

2 About one ml of collected EM was contacted with multi-reagent urinalysis reagent
3 strip (Mission Urinalysis Reagent Strips ACON Laboratories, Inc., U.S.A) for 10-15
4 seconds. pH of EM was recorded based on colour change after 60 seconds (according to
5 the manufacturer recommendations).

6 **2.5 Whiteside test**

7 In WST, equal volume (1ml each) of estrual mucus and 5% Sodium Hydroxide
8 (Merck Specialities Pvt Ltd, India) was taken in a sterile test tube, properly mixed and
9 then heated using spirit lamp to boiling point. The test tube was then cooled under tap
10 water and color change observed. Color change from transparent towards yellow was
11 taken as positive and no color change was taken as negative as described by
12 Bhattacharyya et al. [14].

13 **2.6 Collection of estrual mucus**

14 Collection of estrual mucus (EM) from study animals was carried out at approx
15 6-12 hours post commencement of behavioural oestrus as described by Bhat et al. [15]
16 with slight modification. The study animals during EM collection were properly
17 restrained in the service crate. Lubricated (with paraffin) full sleeve gloved left hand was
18 inserted per rectum for evacuation of rectal faeces. Perineal region including vulva was
19 thoroughly washed with running tap water and then dried with soft absorbent cotton
20 followed by swabbing with ethyl alcohol. While holding both the vulvar lips apart by an
21 assistant, a sterile AI sheath over conventional AI gun was inserted into the vagina, then
22 guided to the cervix and then to uterine body by the hand already introduced per rectum.
23 The external end of the sheath after removing AI gun, was connected to a 20 ml disposable
24 sterile syringe for aspiration of EM from body of uterus and deep cervix. After aspiration,
25 syringe was detached and sheath wiped with ethyl alcohol soaked cotton swab after its

1 removal from genital tract. Sheath containing EM was then transported immediately to
2 lab for further processing.

3 **2.7 Statistical analysis**

4 Receiver Operator Characteristic (ROC) was use to analyze the data of LE, protein
5 and pH strip tests statistically and the area under the curve (AUC) and P values were
6 reported as displayed on the graph generated by ROC analysis. The AUC was used to
7 find optimal cut-off point. The cut-off with the highest sum of sensitivity and specificity
8 was selected as optimal. Data of strip test combinations and data of WST were analysed
9 using diagnostic test evaluation calculator to calculate sensitivity and specificity. ROC
10 and diagnostic test evaluation calculator were both utilized from MedCalc Statistical
11 Software version 19.0.4 (“MedCalc Software bvba, Ostend, Belgium;
12 <https://www.medcalc.org>; 2019”).

13 **3. Results**

14 **3.1 Endometrial cytology**

15 EC was used as a standard diagnostic method to estimate the diagnostic precision
16 of other tests. Based on EC, sixty (60) cows ($\geq 4\%$ PMNs) were found to be positive and
17 18 cows ($< 4\%$ PMNs) were negative (control) for SCE.

18 **3.2 Leukocyte esterase strip test**

19 LE test results were recorded in five categories: – or (negative), \pm or (15 Leu/ μ l),
20 + or (70 Leu/ μ l), ++ or (125 Leu/ μ l), and +++ or (300 Leu/ μ l). Area covering the ROC
21 curve (AUC) was 0.811 with significance level P (Area = 0.5) value less than 0.0001
22 (Figure-1). Based on ROC analysis, the cut-off point selected was greater than 15 Leu/ μ l
23 ($> \pm$) to be used for positive diagnosis of SCE. At the cut-off point, LE test showed
24 sensitivity (Se) of 86.67% and specificity (Sp) of 66.67%. Se and Sp of LE test based on
25 ROC analysis at different cut-off valves is shown in Table-1.

1 **3.3 Protein strip test**

2 Protein results were recorded in five categories: – or (negative), ± or (15 mg/dL),
3 + or (30 mg/dL), ++ or (100 mg/dL), and +++ or (300 mg/dL). Area covering the ROC
4 curve (AUC) was 0.633 with significance level P (Area = 0.5) value of 0.0587 (Figure-
5 2). Based on ROC analysis, the cut-off point selected was greater than 30 mg/dL (> +) to
6 be used for positive diagnosis of SCE. At this cut-off point, protein showed Se of 53.33%
7 and Sp of 66.67%. Se and Sp of protein based on ROC analysis at different cut-off valves
8 is given in Table-2.

9 **3.4 pH strip test**

10 pH results were recorded in six categories: 6.5, 7.0, 7.5, 8.0, 8.5 and 9. Mean pH
11 was found to be 8.23 ± 0.84 . Area covering the ROC curve (AUC) was 0.694 with
12 Significance level P (Area = 0.5) value of 0.0027 (Figure-3). Based on ROC analysis,
13 pH>8 was used as cut off point for positive diagnosis of SCE. At this cut-off point, pH
14 had Se of 66.67% and Sp of 77.78%. Se and Sp of pH based on ROC analysis at different
15 cut-off values is displayed in Table-3.

16 Combination of LE test (cut-off value > ±) and pH (cut-off value > 8) showed Se
17 of 60.00% and Sp of 77.78%. Similarly LE test and protein when used in combination
18 with cut-off value > ± for LE test and > + for protein, showed Se of 46.67% and Sp of
19 77.78% in comparison with EC. When all the three reagent strip test results i.e LE, protein
20 and pH were used in combination using > ± for LE, > + for protein and > 8 for pH as cut-
21 off values, showed Se of 33.33% and Sp of 77.78% in comparison with EC (Table-4).

22 **3.5 Whiteside test**

23 WST results were recorded in four categories: transparent (no infection), light
24 yellow (mild infection), yellow (moderate infection), and dark yellow (severe infection).
25 Out of 78 animals, 46 (59%) showed positive reaction and 32 (41%) showed negative

1 reaction to WST. However, out of 46 positive samples for WST, 39.1% (18) showed light
2 yellow, 43.5% (n=20) showed yellow and 17.4% (n=8) showed dark yellow reaction. In
3 comparison to EC, WST showed Se of 63.33% and Sp of 55.56% (AUC = 0.594).

4 In this study, against EC, LE strip test at cut-off $> \pm$ showed highest Se (86.67%)
5 and 2nd highest Sp (66.67%) among all the diagnostic tests used. Maximum Sp (77.78%)
6 was shown by pH test at cut-off > 8 and its Se (66.67%) was also 2nd highest after LE test.
7 WST also exhibited a near similar Se (63.33%) as that of pH strip test but the Sp (55.56%)
8 of WST was much lower than that of pH strip test as well as LE strip test. Lowest Se
9 (53.33%) was shown by protein at cut-off $> +$ among all the tests used but Se (66.67%)
10 was similar to LE (Figure-4).

11 When the three tests (LE, protein and pH) were used in combination, improvement
12 was seen in Sp (77.78%) and was found to be same in all three combinations; however a
13 severe decline in Se was observed in all combinations. LE strip test and pH in
14 combination showed Se (60.00%) while combination of LE, protein and pH showed
15 lowest Se (33.33%) as illustrated in Figure-4.

16 **4. Discussion**

17 Endometrial cytology using CB is considered as the best minimally invasive
18 diagnostic method for the detection of SCE. This method is believed to be without any
19 negative effect on ensuing conception rate, even if uterine cytology sample is taken after
20 few hours of insemination [16]. EC is neither complex nor costly but it is time consuming
21 and requires equipment and expertise [17], thus cannot act as cow side test, therefore,
22 need for an efficient practical cow side test arises having Se and Sp at par with cytology
23 that can be utilized at field level.

24 LE strip test is one such test which has shown promising results. In this study, LE
25 strip test results showed Se of 86.67% and Sp of 66.67% in comparison to EC and nearly

1 parallel results were stated by Hajibemani et al. [12], with Se of 78% and Sp of 60% for
2 LE strip test with respect to EC in dairy cows. Santos et al. [18] reported similar Se (83%)
3 and contrasting higher Sp (94%) than the present study, which could be due to use of
4 different LE strips (Multistix 10 SG, Bayer Health Care L.L.C., Elkhart, IN, USA) and
5 sampling method i.e. uterine lavage by the investigator. In contrast to the present findings,
6 slightly weak performance of LE strips was reported by Cheong et al. [19] with Se of
7 77% and Sp of 51.8%. Couto et al. [10] also reported lesser Se of 68.8% but demonstrated
8 higher Sp of 72.6% in comparison to present study. These differences may be owing to
9 use of different reagent strip (Multistix 10 SG, Bayer Health Care L.L.C., Elkhart, IN,
10 USA), higher cut-off for EC ($\geq 10\%$ PMN) and LE test ($\geq ++$) compared to the present
11 study (cut-off for EC $\geq 4\%$ PMN and for LE $> \pm$). Further, Couto et al. [10] contacted LE
12 test strips with 0.9% saline in which CB was plunged after rolling it on slides in contrast
13 with estrual mucus in the present study.

14 Fluid accumulation during SCE increases in uterus which could lead to increase
15 in protein concentration [19] and could be used in SCE diagnosis. EM from cows with
16 SCE had markedly elevated intensity of total protein at 21 DPP compared to healthy cows
17 [20]. In the current study, protein strip test results showed Se of 53.33% and Sp of 66.67%
18 in comparison to EC and comparable Se of 58.3% was reported by Cheong et al. [19] but
19 reported lower Sp of 55.8% for protein strip test with respect to EC. It was also reported
20 by Cheong et al. [19] that there is weak association between protein and SCE. This low
21 performance for protein strip test in both studies could be due to reagent strips being
22 highly sensitive for albumin but much less sensitive for other proteins and in addition
23 false positive results may be obtained from samples with high alkalinity and false negative
24 outcome may arise due to high specific gravity [21]. Hajibemani et al. [12] reported that

1 protein concentrations were increased in SCE but protein assessment of discharges could
2 not be recommended for evaluation as a diagnostic means for SCE.

3 pH strip test results of the present study showed Se of 66.67% and Sp of 77.78%
4 with mean pH of 8.23 ± 0.84 in SCE. Results reported by Cheong et al. [19] showed
5 similar Sp (78.4%) but very low Se (44.9%) which could be due to use of different reagent
6 strip (Multistix 10 SG, Bayer Health Care L.L.C., Elkhart, IN, USA), higher threshold
7 level of EC ($\geq 10\%$ PMN) and uterine lavage as sampling method. Bhat et al. [22] and
8 Kumar et al. [23] reported increased pH (8.19 ± 0.06 and 7.95 ± 0.096 ; respectively) in
9 cows with subclinical uterine infection which was in concurrence with this study. Cheong
10 et al. [19] also reported increase in pH of uterine fluid in cows with SCE.

11 Multi-reagent strip tests were used in three different combinations (LE + pH, LE
12 + protein and LE + protein + pH). In all the three combinations Se decreased drastically
13 (60.0%, 46.7% and 33.3% respectively) while Sp increased to 77.8% in comparison to
14 LE test (Se of 86.67% and Sp of 66.67%) alone. The only probable reason that we were
15 able to consider for the relation between the combination of the three tests and the
16 decrease of Se was that pH and protein tests had much lower Se than LE. When the data
17 of the 3 tests was combined and subjected to statistical analysis against standard test (EC),
18 it further decreased the Se due to presence of low number of true positives than true
19 negatives and false negatives concurrently. However, on comparing the three
20 combinations with each other, better performance was shown by combination of LE and
21 pH strip tests with Se of 60.0% and Sp of 77.8%. Cheong et al. [19] reported that LE and
22 pH strip tests in combination increased overall Sp (96.8%) but drastically decreased Se
23 (18.6%) in comparison to LE strip test which was in harmony with this study. Cheong et
24 al. [19] also reported that on combining protein with LE only or with both LE and pH had

1 no effect on increasing the performance of the strip test results, which was in accord with
2 the outcome of present study.

3 In this present study, 59% animals showed positive reaction to WST with Se of
4 63.3% and Sp of 55.6% in comparison to EC. Bhat et al. [15] reported bacteriological
5 growth present in 75% (60/80) animals in which 71.25% (57/80) were positive for WST.
6 Similarly, Gupta et al. [24] used WST for detection of SCE and reported 45.8% (44/96)
7 asymptomatic animals positive to WST.

8 **5. Conclusions**

9 Leukocyte esterase strip test using estrual mucus showed acceptable Se and Sp in
10 comparison to EC by cytobrush; thereby, it can be used as non-invasive, simple,
11 inexpensive and reliable cow side method for screening SCE in cattle under field
12 conditions. However, inclusion of protein and pH strip tests along with LE strip test does
13 not improve the performance of strip tests. WST showed less sensitivity and specificity
14 than LE test, and EC may be replaced by LE test the field for screening SCE in cows.

15 Faster intervention and treatment of SCE is possible by accurate cow-side
16 diagnoses. Further modifications and refinements are required in the field to increase the
17 validity of the test. Moreover, modification of the test strips is necessary to optimize them
18 for diagnosis of SCE as these strip tests are not specific for uterine discharge testing.

19 **Acknowledgements/ Funding/ Conflict of Interest:**

20 The authors are thankful to all those involved in successful completion of this
21 investigation at MLRI Manasbal and FVSc & AH, SKUAST-K, J&K, India.

22 This research did not receive any specific grant from funding agencies in public,
23 commercial, or not-for-profit sectors. There are no benefits in any form from a
24 commercial party or any of the authors related directly or indirectly to the subject of this
25 manuscript.

1 The authors declare that they have no conflict of interest.

2

3 **References**

- 4 1. Nebel RL. Optimizing fertility in the dairy herd. In: Proceedings of the Annual
5 Conference Soc for Therio Nashville, TN, USA 1999; 443–452.
- 6 2. Gilbert RO. The effects of endometritis on the establishment of pregnancy in
7 cattle. *Reproduction, Fertility and Development* 2011; 24: 252-257.
- 8 3. Noakes DE, Parkinson TJ, England GCW. *Arthur’s Veterinary Obstetrics*. 8th ed.
9 St. Louis, Missouri, USA: Saunders; 2001.
- 10 4. Gilbert RO, Shin ST, Guard CL, Erb HN, Frajblat M. Prevalence of endometritis
11 and its effects on reproductive performance of dairy cows. *Theriogenology* 2005;
12 64: 1879-1888.
- 13 5. Kasimanickam R, Duffield TF, Foster RA, Gartley CJ, Leslie KE et al.
14 Endometrial cytology and ultrasonography for the detection of subclinical
15 endometritis in postpartum dairy cows. *Theriogenology* 2004; 62: 9-23.
- 16 6. Bonnett BN, Martin SW, Meek AH. Associations of clinical findings,
17 bacteriological and histological results of endometrial biopsy with reproductive
18 performance of postpartum dairy cows. *Preventive Veterinary Medicine* 1993;
19 15: 205-220.
- 20 7. Williams EJ, Fischer DP, Pfeiffer DU, England GCW, Noakes DE et al. Clinical
21 evaluation of postpartum vaginal mucus reflects uterine bacterial infection and the
22 immune response in cattle. *Theriogenology* 2005; 63: 102-117.
- 23 8. Barlund C, Carruthers T, Waldner C, Palmer C. A comparison of diagnostic
24 techniques for postpartum endometritis in dairy cattle. *Theriogenology* 2008; 69:
25 714-723.

- 1 9. Bajaj NK, Jain SK, Swamy M, Sharma V, Shrivastava OP. Serum haptoglobin
2 concentration to monitor recovery from postpartum sub-clinical endometritis in
3 murrah buffaloes. *Journal of Animal Research* 2015; 5 (4): 753-759.
- 4 10. Couto GB, Vaillancourt DH, Lefebvre RC. Comparison of a leukocyte esterase
5 test with endometrial cytology for diagnosis of subclinical endometritis in
6 postpartum dairy cows. *Theriogenology* 2013; 79: 103-107.
- 7 11. Kutter D, Figueiredo G, Klemmer L. Chemical detection of leukocytes in urine
8 by means of a new multiple test strips. *Journal of clinical chemistry and clinical
9 biochemistry* 1987; 25: 91-94.
- 10 12. Hajibemani A, Mirzaei A, Sharifiyazdi H, Ghasrodashti AR. Vaginal fornix
11 discharge cellularity and its leukocyte esterase activity for diagnosis of
12 endometritis in dairy cows. *Journal of the Faculty of Veterinary Medicine
13 Istanbul University* 2016; 42 (1): 57-64.
- 14 13. Singh J, Honparkhe M, Chandra M, Kumar A, Ghuman SPS et al. Diagnostic
15 efficacy of uterine cytobrush technique for subclinical endometritis in crossbred
16 dairy cattle. *The Indian Veterinary Journal* 2016; 93 (02): 11-13.
- 17 14. Bhattacharyya HK, Makhdoomi DM, Hafiz A, Fazili MR. Clinico-therapeutic
18 management of sub-clinical metritis in cows. *Intas polivet* 2011; 12 (1): 26-27.
- 19 15. Bhat FA, Bhattacharyya HK, Hussain SA. White side test: A simple and rapid test
20 for evaluation of nonspecific bacterial genital infections of repeat breeding cattle.
21 *Veterinary Research Forum* 2014; 5 (3): 177-180.
- 22 16. Kaufmann TB, Drillich M, Tenhagen BA, Forderung D, Heuwieser W.
23 Prevalence of bovine subclinical endometritis 4h after insemination and its effects
24 on first service conception rate. *Theriogenology* 2009; 71: 385-391.

- 1 17. Wagener K, Gabler C, Drillich M. A review of the ongoing discussion about
2 definition, diagnosis and pathomechanism of subclinical endometritis in dairy
3 cows. *Theriogenology* 2017; 94: 21-30.
- 4 18. Santos NR, Roman HB, Gilbert RO. The use of leukocyte esterase reagent strips
5 for diagnosis of subclinical endometritis in dairy cows [Abstract]. *Theriogenology*
6 2006; 66: 666-667.
- 7 19. Cheong S, Nydam D, Galvao K, Crosier B, Ricci A et al. Use of reagent test strips
8 for diagnosis of endometritis in dairy cows. *Theriogenology* 2012; 77: 858-864.
- 9 20. Adnane M, Chapwanya A, Kaidi R, Meade KG, O'Farrelly C. Profiling
10 inflammatory biomarkers in cervico-vaginal mucus (CVM) postpartum: Potential
11 early indicators of bovine clinical endometritis?. *Theriogenology* 2017; 103:
12 117-122.
- 13 21. Sacks D. Clinical diagnosis and management by laboratory methods. 20th ed.
14 Philadelphia: WB Saunders; 2001. ISBN 0-7216-8864-0.
- 15 22. Bhat FA, Bhattacharyya HK, Fazili MR, Hussain SA, Khan MZ. Studies on
16 estrual cervical mucus of repeat breeding cows with special reference to ovulatory
17 disturbances and genital infection. *Theriogenology Insight: An International*
18 *Journal of Reproduction in all Animals* 2015; 5 (2): 113.
- 19 23. Kumar A, Srivastava S, Yadav SK, Yadav AK, Saurabh et al. Physico-chemical
20 characteristics of cervical discharge in endometritic repeat breeder cow.
21 *International Journal of Pure and Applied Bioscience* 2017; 5 (3): 821-831.
- 22 24. Gupta JP, Shyma KP, Mohandas AC, Sneharaj RK. Use of White side test to study
23 subclinical endometritis in cross-bred cattles. *Bhartiya Krishi Anushandhan*
24 *Patrika* 2011; 26 (3&4): 123-125.

25

1
2

Cut off's for number of Leukocytes/μl in estrual mucus	Sensitivity (%)	Specificity (%)
≥ 15 Leu/ μ l (\pm)	100.00	0.00
> 15 Leu/ μ l (\pm)	86.67	66.67
> 70 Leu/ μ l (+)	50.00	88.89
> 125 Leu/ μ l (++)	16.67	100.00
> 500 Leu/ μ l	0.00	100.00

3
4
5

Table-1: Sensitivity and Specificity of Leukocyte Esterase strip test at different cut-off values based on Receiver Operator Characteristic analysis.


Cut off's for amount of estrual mucus protein	Sensitivity (%)	Specificity (%)
≥ 15 mg/dL (\pm)	100.00	0.00
> 15 mg/dL (\pm)	83.33	33.33
> 30 mg/dL (+)	53.33	66.67
> 100 mg/dL (++)	6.67	100.00
> 300 mg/dL (+++)	0.00	100.00

6
7
8
9
10
11
12


Table-2: Sensitivity and Specificity of protein at different cut-off values based on Receiver Operator Characteristic analysis.

Cut off's for pH levels of estrual mucus	Sensitivity (%)	Specificity (%)
≥ 6.5	100.00	0.00
> 6.5	93.33	11.11
> 7	90.00	11.11
> 7.5	73.33	44.44
> 8	66.67	77.78
> 8.5	40.00	88.89
> 9	0.00	100.00


13
14
15
16
17
18
19
20
21
22



1 **Table-3: Sensitivity and Specificity of pH at different cut-off values based on Receiver Operator**



2 **Characteristic analysis.**



3

4 **Figure-1: Area under curve (AUC) plot graph of Leukocyte Esterase strip test.**

1

Figure-2: Area under curve (AUC) plot graph of protein.

2

Figure-3: Area under curve (AUC) plot graph of pH.

Parameter (cut-off)	Se and Sp ¹	%
LE (> ±) ²	Se	86.67
	Sp	66.67
WST ³	Se	63.33
	Sp	55.56
Protein (> +)	Se	53.33
	Sp	66.67
pH (> 8)	Se	66.67
	Sp	77.78
LE and pH in combination	Se	60.00
	Sp	77.78
LE and Protein in combination	Se	46.67
	Sp	77.78
LE, protein and pH in combination	Se	33.33
	Sp	77.78

3 Endometrial Cytology: Positive = 60, negative = 18.

¹ Se: Sensitivity; Sp: Specificity.

² LE: Leukocyte Esterase.

³ WST: White Side Test.




Figure-4: Sensitivity and Specificity of diagnostic tests in comparison to Endometrial Cytology.

(Note: Se: Sensitivity; Sp: Specificity; LE: Leukocyte Esterase; WST: White Side Test)

- 1 **Table-4: Sensitivity and Specificity of Leukocyte Esterase strip test, White Side Test, protein, pH,**
- 2 **combination of Leukocyte Esterase and pH, Leukocyte Esterase and protein, and Leukocyte**
- 3 **Esterase, protein and pH in comparison to Endometrial Cytology.**