

1 **Effects of intracerebroventricularly administered opioid peptide antagonists**

2 **on tissue glycogen levels in rats after exercise**

3
4 **Ayşe Şebnem İLHAN^{1,*}, Şevin GÜNEY², Sibel DİNÇER²**

5
6 ¹Department of Department of Basic Medical Sciences Gülhane Faculty of Dentistry,
7 University of Health Sciences, Ankara, Turkey

8 ²Department of Physiology, Faculty of Medicine, Gazi University, Ankara, Turkey

9
10 ***Correspondence:** aysesebnem@yahoo.com

11
12 **ORCID:**

13 Ayşe Şebnem İLHAN: <https://orcid.org/0000-0001-8746-9378>

14 Şevin GÜNEY: <https://orcid.org/0000-0001-6968-5472>

15 Sibel DİNÇER: <https://orcid.org/0000-0002-0946-2899>

16
17 **Acknowledgement/Disclaimers/Conflict of interest**

18 Our study was approved by Local Ethics Committee for Animal Experiments (G.U.ET-05.043) of Gazi
19 University, and supported by Gazi University, Scientific Research Projects Unit and by The Scientific
20 and Technological Research Council Of Turkey (TUBITAK-BAYG). The experimental procedures on the
21 animals in this study were performed at Gazi University, Laboratory Animals Raising And Experimental
22 Researches Center.

23 Ayşe Şebnem İLHAN and Şevin GÜNEY have received research grants from TUBITAK-BAYG.

24 Ayşe Şebnem İLHAN, Şevin GÜNEY and Sibel DİNÇER declare that they have no conflict of interest.

1 **Effects of intracerebroventricularly administered opioid peptide antagonists on tissue glycogen**
2 **levels in rats after exercise**

3 **Abstract**

4 **Background/aim:** Physical exercise is a state of physiological stress that requires adaptation of the
5 organism to physical activity. Glycogen is an important and essential energy source for muscle
6 contraction. Skeletal muscle and liver are two important glycogen stores and the energy required to
7 maintain exercise in rodents are provided by destruction of this glycogen depot. In this study, the
8 effects of endogenous opioid peptide antagonism at the central nervous system level on tissue
9 glycogen content after exhaustive exercise were investigated.

10 **Materials and methods:** Rats had intracerebroventricularly (icv) received nonspecific opioid peptide
11 receptor antagonist, naloxone (50 µg/10 µL in saline) and δ-opioid receptor-selective antagonist
12 naltrindole (50 µg/10 µL in saline) and then exercised till exhaustion. After exhaustion, skeletal muscle,
13 heart, and liver were excised immediately.

14 **Results:** Both opioid peptide antagonists decreased glycogen levels in skeletal muscle. Although in
15 soleus muscle, this decrease was not statistically significant ($p>0.05$) but in gastrocnemius muscle, it
16 was significant in the icv naloxone administered group compared with control ($p<0.05$). Heart glycogen
17 levels increased significantly in both naloxone and naltrindole groups compared to control and sham-
18 operated groups ($p<0.05$). Heart glycogen levels were higher in the naloxone group than naltrindole
19 ($p<0.05$). Liver glycogen levels were elevated significantly with icv naloxone administration compared
20 with the control group ($p<0.05$). Glycogen levels in the naloxone group was also significantly higher
21 than the naltrindole group ($p<0.05$).

22 **Conclusion:** Our findings indicate that icv administered opioid peptide antagonists may play a role in
23 glycogen metabolism in peripheral tissues such as skeletal muscle, heart, and liver.

24 **Keywords:** Exercise, glycogen, endogenous opioid peptide, naloxone, naltrindole

1 1. Introduction

2 Physical exercise is a state of physiological stress that requires adaptation of the organism to physical
3 activity. Exercise causes an increase in heart rate, blood pressure, tidal volume, and respiratory
4 capacity, but also affects the levels of various hormones and metabolites [1].

5 Glycogen is an important and essential energy source for muscle contraction. Skeletal muscle and liver
6 are two important glycogen stores [2] and the energy required to maintain exercise in rodents is
7 provided by destruction of this glycogen storage. Although muscle glycogen is the biggest source of
8 carbohydrates during exercise, the ratio of destruction depends on the severity of exercise [3]. When
9 the duration of an exercise at a certain severity is prolonged, a decrease in glycogen level occurs [4].
10 Various hormonal and central factors can affect the depletion of muscle glycogen depots during
11 exercise [5,6,7]. Exercise is perceived by the organism as a stress factor [8]. It is known to increase the
12 secretion of endogenous opioid peptides (EOPs) such as β -endorphins, as well as many stress
13 hormones [9].

14 Endorphins are released from the anterior pituitary gland with adrenocorticotrophic hormone (ACTH)
15 in response to stress and in direct proportion to the severity of exercise [10,11]. EOPs are thought to
16 play an important role in the regulation of hormonal and metabolic responses to exercise. It has been
17 shown that increased serum β -endorphin concentration due to exercise can be associated with various
18 psychological and physiological changes in such as mood, perception of pain, hormone levels like ACTH,
19 growth hormone, prolactin, catecholamines, and cortisol; and menstrual irregularities in female
20 athletes [12,13].

21 Endogenous opioid peptides, especially β -endorphins, have been shown to facilitate muscle glucose
22 uptake during exercise [14]. Although there is no evidence that EOPs have a direct effect on glycogen
23 metabolism, they can be expected to be effective on glycogen metabolism by seriously affecting
24 glucose uptake. Endogenous opioid peptide receptors are known to be widely distributed in areas
25 related to glucose homeostases, such as the pancreas and pituitary gland [15]. Although there are

1 studies which show that centrally administered morphine and similar compounds may affect glycogen
2 levels in peripheral tissues such as liver [16], there is no study showing how central opioid peptide
3 receptor antagonism affects glycogen levels in peripheral tissues in relation to exercise.

4 As is known, physiological events and diseases that disrupt the regulation of central opioid release
5 could also affect glycogen metabolism. Therefore, it is necessary to clarify the effects of central OP on
6 glycogen levels in peripheral tissues. Thus, in this study, we aimed to investigate the effects of
7 intracerebroventricularly administered opioid peptide receptor antagonists on glycogen levels in
8 peripheral tissues such as skeletal muscle, heart, and liver after exhausting exercise in rats.

9

10 **2. Materials and methods**

11 In this study, forty Wistar albino male rats, with an average weight of 210 g were used. The rats were
12 fed with standard chow and tap water ad libitum and lived at 12:12 hr light-dark cycle with room
13 temperature at 21 °C as standard animal care condition. According to administrations, rats were
14 divided into five groups as Control (C, n=8), Sham (Sh, n=8), Saline (S, n=8), Naloxone (NX, n=8), and
15 Naltrindole (NT, n=8). In all groups, rats had an exhausting exercise. Among the subjects, those who
16 were injured or refused to run and those who couldn't tolerate isoflurane sedation were excluded.
17 This study was approved by Gazi University Ethical Committee for Experimental Research on Animals.

18 Preliminary studies were carried out for the subjects to familiarize the treadmill. All subjects were
19 exercised on the treadmill for a total of 10 minutes/day for five consecutive days before performing
20 stereotaxic surgery. In this period the speed was 7 m/min and 10 m/min in the first two days then
21 progressively increased to 20 m/min, the angle was always 15° till the familiarization period.

22 Except for the control group, guide cannulas were placed in the left lateral cerebral ventricles of all
23 animals for icv administrations. For this purpose, the subjects were anesthetized with alphazyn (10
24 mg/kg) and ketamine (90 mg/kg) mixture and then placed in a stereotaxic apparatus. The coordinates
25 for icv application were determined as Anteroposterior (A): -0.8 mm, Lateral (L): -1.5 mm and,

1 Dorsoventral (DV): -3.8 mm using the rat brain atlas [17] and referring to the junction point of the
2 lambda and bregma point on the skull. The coordinates were also determined with preliminary
3 experiments and confirmed by methylene blue (Figure) and histological sections.

4 Rats in all groups were sedated 90 minutes before the exhaustive exercise with isoflurane inhalation.
5 Using a Hamilton injector, 10 μ L saline, 50 μ g / 10 μ L naloxone (Sigma Aldrich, Germany) and 50 μ g /10
6 μ L naltrindole (Sigma Aldrich, Germany) were intracerebroventricularly administered to the subjects
7 in saline, naloxone, and naltrindole groups, respectively.

8 Rats were subjected to exercise until they were exhausted on a treadmill at 20 m/ min speed and 15
9 degrees incline. When the subjects refused to run and remained in the supine position for more than
10 10 seconds, they were determined as "exhausted". Blood samples were collected from the heart of
11 rats that were lightly sedated with isoflurane immediately after exhaustion. Rats were decapitated
12 after cervical dislocation and rapidly all tissues were excised and frozen in liquid nitrogen; stored in a
13 -80 °C freezer for glycogen determination.

14 Tissue glycogen levels were measured according to the method of Lo S et al. [18]. Tissue samples
15 weighing 0.3 gr. were placed in test tubes and then 1.5 mL 30% KOH (Merck) saturated with Na_2SO_4
16 (Merck) was added. The tubes were boiled in a boiling water bath for 30 minutes until a homogeneous
17 solution was obtained, then cooled on ice. An equal volume (1.5 mL) of 96% ethanol with the KOH was
18 added. Tissues were kept on ice for 30 minutes and then centrifuged at 2400 rpm for 30 minutes.
19 Supernatants were discarded; the remaining pellets were dissolved in each of 3 mL distilled water. The
20 sample in each tube was mixed with a glass rod for an equal time to dissolve the glycogen in the pellets.
21 1 mL of this mixture was taken and 1 mL of phenol (Fluka, Germany) and 5 mL of 96-98% H_2SO_4 (Merck)
22 were added to them. 1 mL phenol and 5 mL 96-98% H_2SO_4 were added on standard tubes containing
23 1 mL of distilled water and 5 mg/mL glycogen standard (Sigma Aldrich, Germany). Samples in all tubes
24 were vortexed and incubated for 10 minutes at room temperature. After the incubation, the tubes
25 were kept in a water bath set at 30 °C for 20 minutes. 300 μ L samples taken from each tube were

1 placed in a 96-well plate and were read spectrophotometrically at 490 nm against standard and blank.
2 The absorbance of the samples was expressed as mg glycogen / g tissue.

3

4 **2.1. Statistical Analysis**

5 Data analysis was performed using the Statistical Package for Social Sciences (SPSS) v.23.0.
6 Kruskal-Wallis test was used whether there was any difference between the treatments in terms
7 of glycogen level in soleus, gastrocnemius, cardiac, and liver tissues. Bonferroni correction was
8 used for the determination of where the difference originated. Whether there is a difference
9 between administrations in each group, it was compared with Mann Whitney U Test. Values
10 were expressed as mean \pm standard deviation (SD), standard error of the mean (SEM), and
11 median (minimum, maximum) and $p < 0.05$ values were considered as statistically significant.
12 The effect size in calculating the sample size is 0.60, the type I error was taken as 0.05 and the
13 power value as 0.80.

14

15 **3. Results**

16 3.1. Soleus Muscle Glycogen Level

17 The soleus muscle glycogen levels of rats after exhausting exercise are shown in Table 1 as Mean \pm SD
18 and SEM; also in Table 2 as Median (min, max). When the soleus muscle glycogen levels were
19 compared in terms of administrations, no statistically significant difference was observed between the
20 groups ($p > 0.05$).

21

22 3.2. Gastrocnemius Muscle Glycogen Level

1 The gastrocnemius muscle glycogen levels of the rats after exhausting exercise are shown in Table 1
2 as Mean \pm SD and SEM; also in Table 2 as Median (min, max). When the gastrocnemius muscle glycogen
3 levels were compared, in naloxone group there was a statistically significant decrease compared to the
4 control ($p=0.045$).

5

6 3.3. Cardiac Tissue Glycogen Level

7 The heart tissue glycogen levels of the rats after exhausting exercise are shown in Table 1 as Mean \pm
8 SD and SEM; also in Table 2 as Median (min, max). When the groups were compared in terms of cardiac
9 glycogen levels, there were significant increases in naloxone and naltrindole groups compared to
10 control and sham groups ($p=0.006$ and $p=0.003$, respectively). When naloxone and naltrindole treated
11 groups were compared, cardiac tissue glycogen levels in naloxone treated group was found to be
12 statistically significantly higher than naltrindole treated group ($p=0.002$).

13

14 3.4. Liver Glycogen Level

15 The liver tissue glycogen levels of the rats after exhausting exercise are shown in Table 1 as Mean \pm SD
16 and SEM; also in Table 2 as Median (min, max). The glycogen level was found to be statistically
17 significantly higher in the naloxone group compared to the control group ($p=0.045$). When the groups
18 treated with naloxone and naltrindole were compared, liver tissue glycogen levels were found to be
19 statistically significantly higher in the group treated with naloxone compared to the group treated with
20 naltrindole ($p=0.028$).

21

22 4. Discussion

1 In our study, we aimed to investigate the effects of central opioid peptide receptor antagonism on
2 skeletal muscles such as soleus and gastrocnemius as well as heart and liver tissues after exhaustive
3 exercise.

4 Opioid peptides are known to play a role in controlling blood glucose levels during rest and exercise.
5 Circulating levels of β -endorphins, an important component of the opioid system, have been shown to
6 increase with sufficient intensity and time of exercise [19]. There are many studies in the literature
7 showing that β -endorphins and opioid peptide antagonists can be applied by different methods to
8 investigate the role of endogenous opioid peptides in glucose homeostasis during exercise. Hashiguchi
9 et al. administered intracerebroventricularly to demonstrate the effect of central opioids on glycogen
10 levels in peripheral tissues as we did in our study [16]. Radosevich et al. practiced on dogs to
11 demonstrate icv and iv effects of β -endorphins on blood glucose level; they reported that central
12 administration had increased blood glucose levels whereas peripheral application did not affect
13 glucose homeostasis [20]. Various studies also report that these effects of endogenous opioid peptides
14 can be antagonized with naloxone in experimental conditions [21]. Angelopoulos et al. demonstrated
15 that administration of iv naloxone antagonizes the glucagon response increasing the effect of
16 endogenous opioid peptides during exercise in humans [22]. In literature, studies conducted to
17 investigate the central effects of opioid peptides report that applications are generally performed as
18 icv, and for this purpose, guide cannulas are placed with the stereotaxic method after the selection of
19 suitable coordinates. In our study, a guide cannula was placed in the left lateral cerebral ventricle with
20 a similar method. We waited for a week in accordance with the literature [16, 23] for the recovery
21 after stereotaxic surgery. At the end of this period, glycogen levels in skeletal muscles, cardiac and liver
22 tissues of the exhausted rats were investigated. Glycogen is the storage form of glucose found in most
23 cell types and is an important and essential energy source for muscle contraction. Skeletal muscle and
24 liver are the main important glycogen stores in the body as is known [2].

1 In rodents, both liver and muscle glycogen is largely used to maintain exercise. Skeletal muscles
2 primarily use intramuscular glycogen depots during exercise [24] as known.

3 Although muscle glycogen is the biggest source of carbohydrates during exercise, the rate of
4 destruction depends on the severity of exercise [3]. Muscle glucose uptake during exercise is also
5 correlated with the duration and severity of exercise [25-27]. EOPs, especially β -endorphins, were
6 shown to facilitate muscle glucose uptake during exercise [14]. Morphine is also known to be a
7 hormone-like effect that facilitates glucose uptake by muscles. Morphine affects glucose uptake
8 seriously and also affects glycogen metabolism [28]. Physical activity stimulates the release of β -
9 endorphins into the circulation [29]. A strong, nonspecific opioid antagonist naloxone is mostly used
10 in studies to explain the mechanisms related to β -endorphin secretion during exercise.

11 Both of the antagonists we used in our study, reduced glycogen levels in skeletal muscles. Although
12 decreases in glycogen levels are not significant in soleus muscle, in the naloxone group, glycogen levels
13 of the gastrocnemius muscle was found to be significantly decreased compared to the control group.
14 This difference may be due to the different types of muscle fibers.

15 As it is known, soleus muscle is slower and gastrocnemius muscle has more fast-twitch muscle fibers.
16 Evans et al. have shown that the concentration of β -endorphin receptors is higher in fast muscles;
17 hence, stimulation of muscle glucose uptake through β -endorphins in fast muscles is greater [30].
18 Probably, this is the reason of why glycogen levels in the gastrocnemius muscle are generally higher
19 than the soleus muscle in all groups.

20 A significant decrease of glycogen levels in naloxone group of gastrocnemius muscle suggests that the
21 effects of β -endorphins to increase muscle glucose uptake peripherally may be prevented by central
22 antagonism. Therefore, we think that present glycogen stores are spent to supply the energy needs of
23 the muscle. It is known that endorphins-containing nerve cells and opioid peptide receptors are widely
24 distributed in nuclei of the central nervous system, which play a role in the autonomic regulation of
25 the cardiovascular system [31,32]. Otherwise, the presence of opioid receptors on autonomic targets

1 such as the cardiovascular system or lungs proves the presence of direct effects of endorphins in these
2 tissues [33].

3 Myocardial cells are known to have the ability to synthesize, store and secrete opioid peptides [34].
4 Three types of opioid peptides (encephalins, endorphins, and dynorphins) are synthesized in the heart,
5 and the presence of preproenkephalin, prodynorphin, and proopiomelanocortin precursor gene
6 expressions in atrial [35] and ventricular [35,36] cardiac myocytes are reported in the literature.

7 In our study, cardiac glycogen levels in all groups were found higher than skeletal muscle levels. This
8 may be related to the energy metabolism of the heart being different from skeletal muscles. The
9 energy required for myocardial function during rest and exercise is provided by glucose, fatty acids,
10 and lactate. While all three substrates contribute to the reconstruction of ATP at rest, most of the
11 energy comes from free fatty acid breakdown. In the above-moderate exercise, lactate influx to the
12 blood from active skeletal muscle increases a lot and the heart provides most of its energy by oxidation
13 of circulating lactate. In prolonged submaximal exercise, the use of free fatty acids in myocardial
14 metabolism becomes more prominent in supplying the total energy need [37]. This can be the reason
15 why cardiac tissue glycogen level is higher than skeletal muscles in all groups. During exercise, the
16 energy need of the heart is likely to be obtained from other sources, and therefore the use of glycogen
17 is relatively low. In our previous study, we obtained that blood lactate levels after exhausting exercise
18 were higher than pre-exercise levels in all groups [38]. Cardiac glycogen can be higher than skeletal
19 muscle levels because of the heart's using circulating lactate during exhausting exercise as an energy
20 source. When the cardiac tissue is compared in terms of administrations, it has been observed that the
21 opioid peptide antagonism increases the glycogen content of the heart. This increase was found to be
22 significant compared to both control and sham groups; nevertheless, the absence of a significant
23 difference between control and sham groups supports that this effect is caused by antagonist
24 administration.

1 In our study, post-exercise liver glycogen levels were also studied. Glycogen levels in naloxone group
2 were higher among all groups. This increase was statistically significant when compared with the
3 control and naltrindole groups. Hashiguchi et al. showed that intravenously administered morphine
4 and its active metabolites do not alter glycogen levels in peripheral tissues of rats, while
5 intracerebroventricular administered morphine reduces liver glycogen content [16]. Researchers
6 stated that the cannula insertion process for icv administration led to an acute stress response; caused
7 significant increases in plasma corticosterone and catecholamine levels [39] Akil et al. also
8 demonstrated a parallel increase in plasma ACTH and β -endorphine levels in case of stress [15]. Gunion
9 et al. suggested that the roles of opioids in stress response are achieved by stimulating the pituitary-
10 adrenal axis and μ -receptors that activate the sympathetic nervous system [40]. It was shown that
11 endorphins play a role in neurotransmission in sympathetic ganglia in central nervous system and
12 plexuses of the digestive system; in the periphery, it was also shown that they are released into
13 circulation to play a possible hormone role by stimulation of pituitary gland and adrenal medulla [33].
14 Many studies also support that catecholamines and corticosterone can modulate the liver glycogen
15 content. It is reported that epinephrine induces glycogenolysis by modulating phosphorylase activity
16 in the liver and skeletal muscle; corticosterone increases glycogenolytic effects of epinephrine. Thus,
17 hormonal changes caused by icv opiates cause a decrease in liver glycogen levels [16]. Since the
18 organism perceives exercise as a stress factor [8], plasma cortisol and catecholamine levels increase
19 significantly during moderate and high-intensity exercise [41,42].

20 Our findings have shown that intracerebroventricular opioid antagonism causes an increase in liver
21 glycogen levels. This increase is more significant, especially in the naloxone group. Naloxone is known
22 to be a nonspecific antagonist and has a greater affinity for the μ -receptor, although it binds to all
23 three opioid receptors.

24 In our study, although hormone levels such as cortisol and catecholamine were not examined; We
25 think that activation of pituitary-adrenal axis and possible catecholamine release induced by opioid

1 peptides during exercise may be prevented by centrally administered opioid peptide antagonists. Thus,
2 the glycogenolytic effects of endogenous opioid peptides on the liver via the pituitary-adrenal axis and
3 sympathetic nervous system may have decreased relatively, and therefore liver glycogen levels may
4 remain higher in the naloxone-treated group.

5 As a result; findings of our study show that opioid peptide antagonists administered
6 intracerebroventricularly may have an effect on glycogen levels in peripheral tissues such as skeletal
7 muscle, heart, and liver. Although there are many studies in the literature on central opioid peptide
8 antagonism and exhausting exercise no other studies have been found on how central antagonism
9 affects glycogen levels in peripheral tissues after exhausting exercise. On the other hand, since NX and
10 NT are also inverse agonists [43, 44], the basal activity of endogenous OPR has decreased, thus allowing
11 us to see the effects of OPR on glycogen levels more clearly. However, in order to explain the existing
12 and possible mechanisms in more detail, further studies are needed.

13

14 **References**

15 1. Fox EL, Bowers RW, Foss ML. Energy Sources In: The physiological basis of physical education and
16 athletics. 4th ed. Dubuque (Iowa): wcb Publishers, 1989.

17 2. Roach PJ, Skurat AV, Haris RA. Regulation of glycogen metabolism. In: Cherrington AD, Jefferson LS
18 (editors). The endocrine pancreas and regulation of metabolism, (NY), Oxford University Press; 2001.

19 3. Vøllestad NK, Blom PCS. Effect of varying exercise intensity on glycogen depletion in human muscle
20 fibers. *Acta Physiologica Scandinavica* 1985; 125 (3): 395-405. doi: 10.1111/j.1748-
21 1716.1985.tb07735.x

22 4. Hargreaves M, McConell G, Proietto J. Influence of muscle glycogen on glycogenolysis and glucose
23 uptake during exercise. *Journal of Applied Physiology* 1985; 78: 288-292. doi:
24 10.1152/jappl.1995.78.1.288

- 1 5. Kayser B. Exercise starts and ends in the brain. *European Journal of Applied Physiology* 2003; 90 (3-
2 4): 411-419. doi: 10.1007/s00421-003-0902-7
- 3 6. Noakes TD, Peltonen JE, Rusko HK. Evidence that a central governor regulates exercise performance
4 during acute hypoxia and hyperoxia. *The Journal of Experimental Biology* 2001; 204: 3225-3234.
5 doi: 10.1242/jeb.204.18.3225
- 6 7. Molina PE, Abumrad NN. Metabolic effects of opiates and opioid peptides. *Advances in*
7 *Neuroimmunology* 1994; 4 (2): 105-116. doi: 10.1016/s0960-5428(05)80005-1
- 8 8. Rosa Neto JC, Lira FS, Oyama LM, Zanchi NE, Yamashita AS et al. Exhaustive exercise causes an anti-
9 inflammatory effect in skeletal muscle and a pro-inflammatory effect in adipose tissue in rats.
10 *European Journal of Applied Physiology* 2009; 106: 697–704. doi:10.1007/s00421-009-1070-1
- 11 9. Guillemin R, Vargo T, Rossier J, Minick S, Ling N et al. β -endorphin and adrenocorticotropin are
12 selected concomitantly by the pituitary gland. *Science* 1977; 197 (4311): 1367-1369.
13 doi: 10.1126/science.197601
- 14 10. Oleshansky MA, Zoltick JM, Herman RH, Mougey EH, Meyerhoff JL. The influence of fitness on
15 neuroendocrine responses to exhaustive treadmill exercise. *European Journal of Applied Physiology*
16 *and Occupational Physiology* 1990; 59 (6): 405-410. doi:10.1007/BF02388620
- 17 11. Sforzo GA. Opioids and exercise. An update. *Sports Medicine* 1989; 7 (2): 109-124. doi:
18 10.2165/00007256-198907020-00003
- 19 12. Harber VJ, Sutton JR. Endorphins and exercise. *Sports Medicine* 1984; 1: 154-171. doi:
20 10.2165/00007256-198401020-00004
- 21 13. Vaccarino AL, Kastin AJ. Endogenous opiates: 2000. *Peptides* 2001; 22: 2257-2328. doi:
22 10.1016/s0196-9781(01)00566-62

- 1 14. Angelopoulos TJ, Robertson RJ, Goss FL, Utter A. Insulin and glucagon immunoreactivity during
2 high-intensity exercise under opiate blockade. *European Journal of Applied Physiology and*
3 *Occupational Physiology* 1997; 75: 132-135. doi: 10.1007/s004210050137
- 4 15. Akil H, Watson SJ, Young E, Lewis ME, Khachaturian H et al. Endogenous opioids: biology and
5 function. *Annual Review of Neuroscience* 1984; 7: 223-255. doi:
6 10.1146/annurev.ne.07.030184.001255
- 7 16. Hashiguchi Y, Molina PE, Boxer R, Naukam R, Abumrad NN. Differential responses of brain, liver,
8 and muscle glycogen to opiates and surgical stress. *Surgery Today* 1998; 28 (4): 471-474. doi:
9 10.1007/s005950050168
- 10 17. Paxinos G, Watson C. *The rat brain: in stereotaxic coordinates*. 4th ed. San Diego, (CA): Academic
11 Press; 1998
- 12 18. Lo S, Russell JC, Taylor AW. Determination of glycogen in small tissue samples. *Journal of Applied*
13 *Physiology* 1970; 28 (2): 234-236. doi: 10.1152/jappl.1970.28.2.234
- 14 19. Goldfarb AH, Jamurtas AZ. Beta-endorphin response to exercise. An update. *Sports Medicine* 1997;
15 24 (1): 8-16. doi: 10.2165/00007256-199724010-00002
- 16 20. Radosevich PM, Lacy DB, Brown LL, Williams PE, Abumrad NN. Central effects of beta-endorphins
17 on glucose homeostasis in the conscious dog. *The American Journal of Physiology* 1989; 256 (2 Pt 1):
18 E322-330. doi: 10.1152/ajpendo.1989.256.2.E322
- 19 21. Stagg NJ, Mata HP, Ibrahim MM, Henriksen EJ, Porreca F et al. Regular exercise reverses sensory
20 hypersensitivity in a rat neuropathic pain model: role of endogenous opioids. *Anesthesiology* 2011;
21 114 (4): 940-948. doi:10.1097/ALN.0b013e318210f880
- 22 22. Angelopoulos TJ, Robertson RJ, Goss FL, Utter A. Insulin and glucagon immunoreactivity during
23 high-intensity exercise under opiate blockade. *European Journal of Applied Physiology and*
24 *Occupational Physiology* 1997; 75 (2): 132-135. doi: 10.1007/s004210050137

- 1 23. Soares DD, Lima NR, Coimbra CC, Marubayashi U. Evidence that tryptophan reduces mechanical
2 efficiency and running performance in rats. *Pharmacology, Biochemistry, and Behavior* 2003; 74 (2):
3 357-362. doi: 10.1016/s0091-3057(02)01003-1
- 4 24. Ryder JW, Kawano Y, Galuska D, Fahlman R, Wallberg-Henriksson H et al. Postexercise glucose
5 uptake and glycogen synthesis in skeletal muscle from GLUT4-deficient mice. *FASEB journal: official
6 publication of the Federation of American Societies for Experimental Biology* 1999; 13 (15): 2246-2256.
7 doi: 10.1096/fasebj.13.15.2246
- 8 25. Romijn JA, Coyle EF, Sidossis LS, Gastaldelli A, Horowitz JF et al. Regulation of endogenous fat and
9 carbohydrate metabolism in relation to exercise intensity and duration *The American Journal of
10 Physiology*,1993; 265: E380–391. doi: 10.1152/ajpendo.1993.265.3.E380
- 11 26. Katz A, Broberg S, Sahlin K, Wahren J. Leg glucose uptake during maximal dynamic exercise in
12 humans. *The American Journal of Physiology* 1986; 251: E65–70. doi: 10.1152/ajpendo.1986.251.1.E65
- 13 27. Katz A, Sahlin K, Broberg S. Regulation of glucose utilization in human skeletal muscle during
14 moderate dynamic exercise. *The American Journal of Physiology* 1991; 260: E411–415. doi:
15 10.1152/ajpendo.1991.260.3.E411
- 16 28. Peng CH, Walsh O. Effects of morphine on the hormonal control of metabolism. I. In vitro effects
17 of morphine and hydrocortisone on utilization of glucose by muscle of normal and chronically
18 morphinised rats. *Biochemical Pharmacology* 1963; 12: 921-930. doi: 10.1016/0006-2952(63)90014-5
- 19 29. Schwarz L, Kindermann W. Changes in beta-endorphin levels in response to aerobic and anaerobic
20 exercise. *Sports Medicine* 1992; 13 (1): 25-36. doi: 10.2165/00007256-199213010-00003
- 21 30. Evans AA, Tunnicliffe G, Knights P, Bailey CJ, Smith ME. Delta opioid receptors mediate glucose
22 uptake in skeletal muscles of lean and obese-diabetic (ob/ob) mice. *Metabolism* 2001; 50 (12): 1402-
23 1408. doi: 10.1053/meta.2001.28158

- 1 31. Elde R, Hökfelt T, Johansson O, Terenius L. Immunohistochemical studies using antibodies to
2 leucine-enkephalin: initial observations on the nervous system of the rat. *Neuroscience* 1976; 1 (4):
3 349-351. doi: 10.1016/0306-4522(76)90063-4
- 4 32. Goodman RR, Snyder SH, Kuhar MJ, Young WS 3rd. Differentiation of delta and mu opiate receptor
5 localizations by light microscopic autoradiography. *Proceedings of the National Academy of Sciences*
6 of the United States of America 1980; 77 (10): 6239-6243. doi: 10.1073/pnas.77.10.6239
- 7 33. Holaday JW. Cardiovascular consequences of endogenous opiate antagonism. *Biochemical*
8 *Pharmacology* 1983; 32 (4): 573-585. doi: 10.1016/0006-2952(83)90479-3
- 9 34. Barron BA, Jones CE, Caffrey JL. Pericardial repair depresses canine cardiac catecholamines and
10 met-enkephalin. *Regulatory Peptides* 1995; 59 (3): 313-320. doi: 10.1016/0167-0115(95)00086-q
- 11 35. McLaughlin PJ, Wu Y. Opioid gene expression in the developing and adult rat heart. *Developmental*
12 *Dynamics* 1998; 211 (2): 153-163. doi: 10.1002/(SICI)1097-0177(199802)211:2<153::AID-
13 AJA4>3.0.CO;2-G
- 14 36. Springhorn JP, Claycomb WC. Preproenkephalin mRNA expression in developing rat heart and in
15 cultured ventricular cardiac muscle cells. *The Biochemical Journal* 1989; 258 (1): 73-78. doi:
16 10.1042/bj2580073
- 17 37. McArdle WD, Katch FI, Katch VL. *Exercise physiology; energy, nutrition and human performance.*
18 6th ed. Philadelphia, Pennsylvania: Lippincott Williams & Wilkins; 2007.
- 19 38. Güney Ş, İlhan AŞ, Çetin F, Dinçer S. The effects of intracerebroventrically administered opioid
20 peptide receptor antagonists on exercise performance. *Isokinetics and Exercise Science* 2012; 20: 205-
21 209. doi: 10.3233/IES-2012-0460
- 22 39. Hashiguchi Y, Molina PE, Abumrad NN. Morphine-3-glucuronide: hyperglycemic and
23 neuroendocrine potentiating effects. *Brain Research* 1995; 694 (1-2): 13-20. doi: 10.1016/0006-
24 8993(95)00697-o

1 40. Gunion MW, Rosenthal MJ, Morley JE, Miller S, Zib B et al. μ -receptor mediates elevated glucose
2 and corticosteron after third ventricle injection of opioid peptides. The American Journal of
3 Physiology 1991; 261: R70-R81. doi: 10.1152/ajpregu.1991.261.1.R70

4 41. Hill EE, Zack E, Battaglini C, Viru M, Viru A, Hackney AC. Exercise and circulating cortisol levels: the
5 intensity threshold effect. Journal of Endocrinological Investigation 2008; 31 (7): 587-591.
6 doi:10.1007/BF03345606

7 42. Sgherza AL, Axen K, Fain R, Hoffman RS, Dunbar CC et al Effect of naloxone on perceived exertion
8 and exercise capacity during maximal cycle ergometry. Journal of Applied Physiology 2002; 93: 2023-
9 2028. doi:10.1152/jappphysiol.00521.2002

10 43. Sullivan LC, Chavera TS, Jamshidi RJ, Berg KA, Clarke WP. Constitutive desensitization of opioid
11 receptors in peripheral sensory neurons. The Journal of Pharmacology and Experimental
12 Therapeutics. 2016; 359 (3): 411-419. doi: 10.1124/jpet.116.232835

13 44. Higashi E, Hirayama S, Nikaido J, Shibasaki M, Kono T et al. Development of novel δ opioid
14 receptor inverse agonists without a basic nitrogen atom and their antitussive effects in mice. ACS
15 Chemical Neuroscience. 2019; 10 (9): 3939-3945. doi: 10.1021/acscchemneuro.9b00368

16

17

18

19

20

21



1

2 Figure. Verification of intracerebroventricular coordinates with methylene blue application.

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

1 Table 1. Mean, standard deviation, and standard error of the mean values of glycogen levels in soleus,
 2 gastrocnemius, cardiac, and liver tissue samples of rats after exhaustive exercise

Gr	Soleus		Gastrocnemius		Cardiac		Liver	
	Mean±SD	SEM	Mean±SD	SEM	Mean±SD	SEM	Mean±SD	SEM
C	0.70±0.52	0.23	1.16±0.53	0.24	1.36±0.11	0.05	1.88±0.51	0.23
SH	1.43±2.29	1.02	1.00±1.07	0.48	1.36±0.20	0.09	2.21±0.25	0.11
S	0.88±1.92	0.68	0.70±0.38	0.14	2.47±1.00	0.35	2.27±0.44	0.16
NX	0.4±0.44	0.18	0.60±0.62	0.25	3.21±0.83	0.34	2.45±0.38	0.15
NT	0.91±1.11	0.39	0.83±0.47	0.17	2.36±0.16	0.05	1.96±0.28	0.10

3

4 Gr: Groups; C: Control; SH: Sham; S: Saline; NX: Naloxone; NT: Naltrindole

5 SD: Standard deviation; SEM: Standard error of the mean

6

7

8

9

10

11

12

13

14

15

16

1 Table 2. Median (minimum, maximum) values of glycogen levels in soleus, gastrocnemius, cardiac and
 2 liver tissue samples of rats after exhaustive exercise

Groups	Soleus Median (min, max)	Gastrocnemius Median (min, max)	Cardiac Median (min, max)	Liver Median (min, max)
C	0.90 (0.02, 1.29)	1.06 (0.60, 2.03) ^a	1.42 (1.23, 1.47) ^{a,b}	1.73 (1.31, 2.68) ^a
SH	0.82 (0.01, 5.46)	0.70 (0.27, 2.88)	1.29 (1.11, 1.61) ^{c,d}	2.14 (1.94, 2.56)
S	0.11 (0.01, 5.60)	0.62 (0.28, 1.38)	2.88 (1.17, 3.62)	2.21 (1.59, 2.90)
NX	0.34 (0.02, 1.15)	0.44 (0.14, 1.81) ^a	2.90 (2.72, 4.88) ^{a,c,e}	2.41 (2.04, 2.93) ^{a,b}
NT	0.44 (0.03, 3.27)	0.90 (0.26, 1.51)	2.32 (2.17, 2.59) ^{b,d,e}	1.88 (1.57, 2.54) ^b
p, Kruskal Wallis Test	> 0.05	< 0.05	< 0.05	< 0.05
p, Bonferroni correction		^a C-NX= 0.045	^a C-NX=0.006 ^b C-NT=0.003 ^c SH-NX=0.006 ^d SH-NT=0.003 ^e NX-NT=0.002	^a C-NX=0.045 ^b NX-NT=0.028

3

4 C: Control; SH: Sham; S: Saline; NX: Naloxone; NT: Naltrindole

5 min: Minimum; max: Maximum

6

7