

High heat generating granites of Kestanbol: future enhanced geothermal system (EGS) province in western Anatolia

Dornadula CHANDRASEKHARAM*^{ORCID}, Alper BABA^{ORCID}

İzmir Institute of Technology, İzmir, Turkey

Received: 17.06.2021 • Accepted/Published Online: 17.08.2021 • Final Version: 01.12.2021

Abstract: Although the western Anatolian region is a foci for hydrothermal systems, this region has several high heat-generating granitic intrusive bodies that qualify to be candidates for enhanced geothermal systems (EGS). Considering the future energy requirement, carbon dioxide emissions reduction strategies, food, and water security issues, these granites appear to be the future clean energy source for the country. One such granite intrusive is located in the Kestanbol area in the western Anatolian region. The radioactive heat generation of this 28 Ma old granite varies from 5.25 to 10.38 $\mu\text{W}/\text{m}^3$ with a heat flow of 92.47 to 128.61 mW/m^2 . These values concur with the measured geothermal gradients and heat flow values measured from exploratory bore wells. High radon content in the thermal waters in these areas indicates interaction between the circulating fluids and the Kestanbol granite. This is for the first time evaluation of the EGS potential of granite intrusive in Turkey has been made. The Kestanbol intrusive is placed under a compressive stress regime within the Anatolian-Aegean regional tectonic framework.

Key words: Geothermal energy, EGS, radionuclide, granite, Turkey

1. Introduction

Within the Alpine-Himalayan orogenic regime (Tethys regime), the Anatolian fault zone and associated tectonic structures and geothermal provinces occupy an important segment. The Paleotethyan and Neotethyan ocean basins outcrop between the E-W trending tectonic belts, namely, the Pontides, Anatolides, and Taurides. The E-W trending Neotethyan Subduction zones hosts, besides obducted Cretaceous ophiolites, several granitoid intrusives (Bingol et al., 1982; Örgün et al., 2007; Dilek et al., 2009; Şahin et al., 2010; Black, 2012; Angı et al., 2016). These granitoids outcrop at several places in the western, central, northeast, southeast Anatolia. The granitoids in west Anatolia is of Eocene-Oligo-Miocene in age, while the rest belong to the Late Cretaceous age. These granitoids show high natural radioactivity levels due to high concentrations of uranium, thorium, and potassium. As a result, these rocks generate abnormal heat greater than the heat generated by normal granites discussed in the later sections. The heat can be extracted through circulating fluids, and the heat can be utilized for power generation and other direct applications. In this paper, our focus is on the Kestanbol granitoids of the Biga Peninsula in western Anatolia. This 28 Ma granitoid is intruded into the Rhodope-Serbo-Macedonian Massif and outcrops over an area of 16 sq. km. The Upper

Miocene-Pliocene coarse-grained clastic and shallow marine carbonates overly intrusive. The concentration of uranium, thorium, and potassium in these granites is the highest of all the granites of Turkey. The distribution of granites in Anatolia is shown in Figure 1.

2. Geology of western Anatolia

During the Cenozoic Era, western Anatolia experienced intensive magmatic activity represented by volcanic and plutonic rocks (Figure 2). Several authors have reported the geology, geochemistry and tectonic configuration of these rocks (Şengör and Yılmaz, 1981; Yılmaz, 1989; Güleç, 1991; Harris et al., 1994; Altunkaynak and Yılmaz, 1998; Aldanmaz et al., 2000; Okay and Satır, 2000, 2006; Köprübaşı and Aldanmaz, 2004; Altunkaynak and Dilek, 2006, 2013; Dilek and Altunkaynak, 2007, 2010; Altunkaynak and Genç, 2008; Boztuğ et al., 2009; Ersoy et al., 2009; Erkül, 2010, 2012; Hasözbeke et al., 2010; Altunkaynak et al., 2010, 2012a, 2012b; Erkül and Erkül, 2012; Erkül et al., 2013; Papadopoulos et al., 2016). The plutonic rocks are represented by I type granitoids and medium to high potassium calc-alkaline rocks (Harris et al., 1994; Köprübaşı and Aldanmaz, 2004; Altunkaynak, 2007; Altunkaynak et al., 2012a). All the granitic intrusive occur along fault zones (Figure 2). The older Eocene

* Correspondence: dchandra50@gmail.com




Figure 1. Occurrence of granites in Turkey (modified after Akbaş et al., 2011).

granite plutons outcrop along the İzmir-Ankara Suture (IAS) zone. Greater than 12 granite and granitic plutons outcrop in the western Anatolia zone (Figure 2).

The Eocene granitoids include granite, quartz diorite, granodiorite, syenite, and monzogranite. They are intruded into the Cretaceous blue-schist and ophiolites. The quartz diorite, granodiorite, and syenite occur around Orhaneli, Topuk, and Gurgenyala. In contrast, monzogranite, granodiorite, and granite occur around Yalova, Fıstıklı (Armutlu), Karabiga, and Kapıdağ, south of Marmara Sea and north area (IAS). The quartz diorite, granodiorite, and syenite are intruded into the Cretaceous blueschists and ophiolites. At the same time, the 35 million-year-old monzogranite and granite are emplaced into the metamorphic rocks of western Pontides (Delaloye and Bingöl, 2000; Altunkaynak et al., 2012 a,b). The older Miocene granitoids (younger) exposed towards the west of Anatolia contain higher radioactive elements such as U, Th and K compared to those granitoids of Eocene age. Thus, the heat-generating capacity of these granites is higher relative to the Eocene (older) granites. This paper focuses on the Miocene granites of western Anatolia, particularly those occurring in the Kestanbol Region.

3. Geothermal gradient and heat flow over western Anatolia

The Aegean Sea, adjacent to western Anatolia, are loci of intense tectonic activity. This region is subjected to intense crustal extension and subduction accompanied by

intense andesitic volcanism during the Oligocene (Fytikas et al., 1984; Dilek et al., 2009; Jolivet et al., 2015). These tectonic and volcanic activities have resulted in two major crustal extension regimes: an early E-W extension during Miocene to Early Pliocene and N-S extension during Pliocene to Quaternary. The younger extensional regime resulted in the formation of horst-graben structures in the Menderes Massif (Kocyigit et al., 1999). In addition, these active tectonic regime has resulted in high heat flow and high geothermal gradient in this region (Erickson et al., 1976; Eckstein, 1978).

Several deep exploratory geothermal gradient wells in and around the Tuzla geothermal field located south of Çanakkale in the western Anatolian region registered very high geothermal gradients and high heat flow values (Figure 3). The temperature of 145 °C was recorded at 50 m depth, and well-blow outs occurred due to high steam and boiling environment at such depths. Deep exploratory wells drilled to 800–1020 m into the pyroclastics recorded bottom hole temperatures of 173 °C (Karamandereci and Ongur, 1974; Baba et al., 2005).

High-resolution equilibrium temperatures from 113 boreholes with a depth of 100 m were analyzed to determine the conductive heat flow in the western Anatolian region (Erkan, 2015). The coastal region extending from İzmir to Çanakkale showed elevated heat flow values varying from 85 to 95 mWm², while the region over the Menderes Massif recorded values above 100 mWm². The high heat flow values are associated with deep-seated normal or




Figure 2. Simplified geological map of western Anatolia and the eastern Aegean region (modified after Dilek and Altunkaynak, 2009).




Figure 3. Geothermal sources of Turkey and their surface temperatures with favorability analysis (geothermal source data taken from Akkus et al., 2005; Basemap Imagery from Earthstar Geographics, Esri, HERE, Garmin, FAO, NOAA, USGS).

strike-slip faults and volcanic centers. These values are typical of regions related to orogenic (Mesozoic-Cenozoic) and volcanic activity (Cenozoic). In addition to borehole data, aeromagnetic data was also utilized to understand the subsurface structures responsible for high heat flow values (Eckstein, 1978). The Curie point temperature is essential to substantiate the anomalous heat flow in western Anatolia. The Curie point temperature (CPT) deduced from aeromagnetic was published for the western Anatolian region (Karat and Aydin, 2004). The high heat flow values lie over the regions where CPT is shallow.

In addition to the borehole exploration, airborne magnetic data was also employed to estimate the geothermal gradient and heat flow values using Curie point depth (CPD), obtained airborne magnetic maps, was utilized to estimate the geothermal gradient. Heat flow values were obtained from the conductivity values and geothermal gradient. The heat flow value obtained varies from 100 to 160 mW/m² along the coastal region, extending from İzmir to Çanakkale (Akin et al., 2014). The Curie point of depth in these sites is also shallow varying from 10 to 6 km.

The sites that recorded high heat flow values and shallow CPD include Balıkesir, İzmir, Manisa, Aydın, Denizli, and Çanakkale.

Although conventional heat flow measurements along western Anatolia are limited, heat flow measurements based on bottom hole temperatures established reasonable heat flow maps for the entire region (Tezcan and Turgay, 1991). The same data have been utilized to establish the geothermal gradient in this region.

The heat flow values vary from 50 to 133 mW/m², and the corresponding geothermal gradient varies from 39 to 57 °C/km; the higher values are recorded along the Aegean Sea coast of İzmir and Çanakkale, i.e. Çanakkale and the peninsular part of İzmir. The Curie point depth calculated based on the aeromagnetic anomaly map along western Anatolia varies from 12 km (near İzmir) to 19 km over Çanakkale. In addition to the active tectonic regime of the Çanakkale region, resulting in high heat flow values, the presence of fertile granites in this region (granite with high content of uranium, thorium, and potassium) is making this region most suitable for initiating projects related to EGS (enhanced geothermal systems).

4. Geothermal provinces of western Anatolia

Due to high heat flow and geothermal gradients associated with active and intense tectonic and volcanic activities, this region has developed high enthalpy geothermal systems (with recorded reservoir temperatures approximately 240 °C) along the western Anatolia, represented by numerous thermal springs, with fluids circulating along the deep faults associated with the horst and graben structures (Serpen et al., 2009; Ugur et al., 2014). The surface manifestations of the geothermal systems are represented by thermal springs (Figure 4) with temperatures varying from 34 to 80 °C.

Exploratory bore-wells drilled near Tuzla (south of Kestanbol) indicate high-temperature geothermal systems in this province with a recorded bottom hole temperature of 145 °C from a 50 m deep bore well (Baba et al., 2005). Similarly, two exploratory bore wells drilled to a depth of about 1000 m reveal high-temperature systems at 333 m depth with a recorded temperature of 175 °C. Well blow-outs in this region indicate the presence of high-pressure geothermal systems. The geothermal systems are of two-phase with 13% steam and a fluid flow rate of 130 t/h (Baba et al., 2005). The presence of high-temperature hydrothermal alteration assemblages indicate reservoir temperature located in the pyroclastics of the order of 220 °C (Sener and Gevrek, 2000; Baba et al., 2005).

The Kestanbol thermal springs (47–68 °C) are historically famous for their healing properties. There are two groups of thermal springs, one with high sulfur content and the other with high radon content due to high radioactivity (Demirsoy et al., 2018). The presence of radionuclides has been established, and the source of the radionuclides is the high radiogenic Kestanbol granites (Baba et al., 2008).

A detailed account of the geothermal manifestation of Kestanbol was given by Baba and Ertekin (2007). The issuing temperature of the thermal springs varies from 66 to 76 °C with a flow rate of 6 L/s. Located near the seashore, the thermal water show mixing of seawater represented by high Na-Cl content. In addition, tritium content varies from 0.22 to 0.25 TU indicating deep circulation of the thermal fluids (Baba and Ertekin, 2007).

5. The Kestanbol granites

Western Anatolia experienced extensive magmatic activity during Eocene to Miocene period, represented by plutonic and volcanic activities (Yilmaz, 1997; 1998; Delaloye and Bingol, 2000; Yilmaz et al., 2001; Arik and Aydin, 2011). During this period, this region was under lithospheric spreading and crustal thinning (Aldanmaz, 2006). The earlier magmatic activity was represented by granitic pluton, and basaltic lava flows represented the late phase.

Kestanbol, located in northwestern Turkey, hosts young granitic and volcanic rocks. The younger granites were intruded into the metamorphic basement giving rise to a

contact metamorphic aureole (Figure 5). The Kestanbol granitoid intruded into the metasedimentary rocks is a quartz monzonite related to the collision tectonic between Anatolian-Tauride and Pontides that occurred during the Late Cretaceous period. This N-S convergence continued until the Neogene period giving rise to magmatic activity in the Early Miocene (Karacik and Yilmaz, 1998; Sahin et al., 2010). The magmatic activity was represented by both intrusive and extrusive phases.

The volcanic rocks associated with the Kestanbol granites include lava flows, ignimbrites, and lahar deposits. The radiometric age ($^{40}\text{Ar}/^{39}\text{Ar}$) of the Kestanbol granites varies from 22.21 to 21.22 Ma (Early Miocene) (Akal, 2013). The Kestanbol granites are characterized by high uranium, thorium, and potassium content compared to other younger Eocene and Miocene granites of western Anatolia (e.g., Kozak pluton, Eybek pluton, Eğrigöz pluton, Koyunoba pluton, Karaburun granodiorite).

The Kestanbol quartz monzonite, emplaced into the regionally metamorphosed basement rocks, encloses several enclaves and is traversed by several dykes of aplite, pegmatite, mafic lamprophyre, and latite. The granitoid mass is widely exposed around the Kocali and Alada villages of Kestanbol (Arik and Aydin, 2011). The Kestanbol pluton was derived from crustal melts contaminated with mantle-derived mafic magma during its formation (Yilmaz et al., 2010). The Kestanbol quartz monzonites are holocrystalline with porphyritic texture with large potash feldspar megacrysts. Besides K-feldspars, these rocks contain plagioclase, quartz, biotite, hornblende and pyroxene in the groundmass (Arik and Aydin, 2011). The presence of thorite, uranothorite, allanite, and zircon either as inclusions in biotite and hornblende or as individual minerals, in considerable amounts, makes these rocks highly radiogenic (Örgün et al., 2007). Several dikes (approximately 2 m) of aplite, pegmatite, granophyre, and lamprophyre are found traversing the Kestanbol granitoid. These intrusive, together with hydrothermal alterations, created zones with a high concentration of radioactive minerals making these granitoids highly radiogenic (Orgun et al., 2007).

5.1. Radioactive characteristics of Kestanbol granites

The Kestanbol granitoid, due to the presence of significant content of radioactive minerals described above, is characterized by high radioactivity. Even the air around the area has registered very high gamma radiation levels varying from 46 to 9200 nGy/h (nanoGray/hour). Even the site near Kestanbol thermal springs reported a value of 880 nGy/h (Orgun et al., 2007). These values are considered very high for this region. The measured ^{137}Cs activities in the rock samples vary from 0.9 to 6.57 Bq/kg, which is regarded as very high. The concentration of U, Th, and K in the Kestanbol granitoid is shown in Table 1.




Figure 4. Geothermal provinces and thermal springs of Western Anatolia (modified after Akkus et al., 2005; Baba and Sozbilir, 2012; tectonic structures digitized from Emre et al., 2013).




Figure 5. Granite exposure around Kestanol (modified after Mützenberg, 1997).

The heat flow values (Table 1) calculated based on the RHP are similar to those reported based on field measurements, and CPD estimation reported (Eckstein, 1978; Karat and Aydin, 2004; Akin et al., 2014).

The radioactive heat production (RHP in $\mu\text{W}/\text{m}^3$) by granites has been calculated using the heat generation

constant and the uranium, thorium, and potassium concentrations C_U , C_{Th} , C_K using equation suggested by Rybach (1976) and Cermak et al. (1982):

$$RPH = \rho(9.52 C_U + 2.56 C_{Th} + 3.48 C_K) \times 10^{-5}$$

where ρ is the density of rock in kg/m^3 ; C_U and C_{Th} are the concentration of U and Th in mg/kg , respectively,

Table 1. The heat generation of Kestanol granites and the heat flow value over the region are based on U, Th, and K content (U, Th, and K contents are from Orgun et al. 2007).

Sample no.	U (ppm)	Th (ppm)	K (wt %)	RHP ($\mu\text{W}/\text{m}^3$)	HF (mW/m^3)
1	11.90	50.00	3.74	6.87	108.67
2	8.20	54.00	3.95	6.21	102.11
3	8.30	62.00	4.14	6.81	108.08
4	17.40	80.00	3.98	10.38	143.76
5	16.10	59.00	3.83	8.58	125.76
6	14.30	62.00	3.92	8.33	123.29
7	15.70	61.00	3.76	8.61	126.05
8	16.30	62.00	4.11	8.86	128.61
9	15.90	59.00	3.92	8.53	125.33
10	14.00	62.00	3.91	8.25	122.51
11	10.70	47.00	3.82	6.36	103.58
12	11.80	58.00	3.88	7.41	114.07
13	10.40	42.00	3.76	5.93	99.30
14	12.60	53.00	3.69	7.25	112.49
15	17.00	47.00	3.49	7.95	119.47
16	9.70	47.00	3.38	6.06	100.59
17	9.60	40.00	3.81	5.59	95.90
18	7.50	43.00	3.69	5.25	92.47
19	12.30	65.00	3.70	8.00	120.02
20	14.10	54.00	3.72	7.71	117.06
21	7.30	36.00	3.56	4.70	86.99
22	11.10	47.00	4.03	6.48	104.80
29	15.40	59.00	3.77	8.39	123.91
32	14.30	65.00	3.88	8.53	125.33
27	9.70	50.00	4.57	6.38	103.78
61	9.90	40.00	3.76	5.66	96.63
62	10.80	63.00	3.67	7.48	114.75

and C_K is the concentration of K in weight percentage in the granites. The surface heat flow values were calculated using the proposed equation by (Lachenbruch, 1968)

$$Q = Q_0 + D \times A$$

where Q is the heat flow at the surface, Q_0 is an initial value of heat flow unrelated to the specific decay of radioactive element at a certain time, D is the thickness of rock over which the distribution of radioactive element is more or less homogeneous, and A is the radioactive heat production. Since the thin crustal thickness (approximately 25km) is observed in the coastal region of the western part of Turkey (Tezel et al., 2013), therefore, the background heat flow value 40 mW/m^2 is considered in the west part of Turkey. Based on the heat flow value, the subsurface temperature has been calculated using the following relation (Vernekar, 1975)

$$Q = \left(\frac{dT}{dz} \right)$$

where k is the thermal conductivity of the rock and dT/dz is the geothermal gradient. The surface temperature has been calculated by taking the average surface temperature of about 25 °C (Vernekar, 1975) and thermal conductivity of the granitic rock as 3.8 $\text{Wm}^{-1}\text{C}^{-1}$.

6. Stress field status of western Anatolia

The western Anatolian region was under compression due to several collision events from Mesozoic to Early Tertiary, resulting in structural fabric folds and faults. The initial structural fabric was trending NW-SE in the eastern Aegean Sea, changing to E-W and ENE to WNW across the western Anatolian region. The major regional forces act on the western Anatolia northward movement

of the African plate, northwest movement of the Arabian plate, and west and SW movement of the Anatolian plate culminating into the Aegean arc in the Aegean Sea west of Turkey (Figure 6).

Western Turkey is an active crustal extension zone. This zone is located south of the North Anatolian Fault Zone (NAFZ) and north of the Aegean subduction zone (Figure 6). The extension due to westward motion of Turkey (strike-slip fault associated with the North Anatolian Fault system, moving at the rate of 36 mm/year) relative to Eurasian is accommodated by the shortening in Aegean subduction zone (McKenzie, 1972, Taymaz et al., 1991, Jackson, 1994)

Detailed stress field analyses were carried out by Rabai et al. (1992) using earthquake focal mechanism, in situ stress measurements (nearly 284 measurements) based on hydraulic fracturing, well blow-outs, over coring, and flat-jack procedure (Rabai et al., 1992) for regions covering the western and eastern Mediterranean region, northern Africa and NW Arabia and the Russian plates. These regions exert forces on the North Anatolian Fault Zone in

the northern part of Turkey. Based on the above data Rabi et al. (1992) evolved a regional stress field map for the entire regions. Based on a simple numerical approach to calculate the S_{hmax} and S_{hmin} directions was developed by Rabi et al. (1992). In the Anatolian region, the S_{hmax} is perpendicular to the N-S convergence between the Arabian and Russian plate (Figure 6) and changes progressively from NW-SE (in the east Anatolia) to NE-SE (in the western Anatolia). The stress state changes from compressional in the east to extensional in the west. The Anatolian lateral movement is absorbed by the Aegean trench; a part of this lateral stress is resulting in deformation of the continental blocks present between the Anatolian fault zone and the Aegean trench. This implies that all the rock formations along the western part of the Bagan peninsula are under a compressive stress regime.

7. Discussion

The western part of Turkey is loci of several geothermal provinces represented by hundreds of thermal springs with temperatures varying from 40 to 86 °C. The province




Figure 6. Major regional tectonic regimes over western Anatolia (modified after Sengor and Dyer, 1979); Sengor (1980), Barka (1992), Bozkurt (2001), Kocycigit and Ozacar (2003); modified after from Baba et al. (2021).

that falls within this region includes Kestanol and Tuzla (Çanakkale). The geothermal manifestations are associated with deep-seated faults, large sedimentary basins, and volcanic sites. This region is also represented by several plutonic rocks of the Miocene age, such as the Kestanol granitoid. These granitoid, due to their crustal origin, contain a high concentration of radioactive elements (U, Th, and K) due to the presence of minerals such as thorite, zircon, and allanite. The heat generated by the Kestanol granitoid is 5 to 8 mW/m², which is greater than the average heat generated by the granites of 5 mW/m². The gamma-ray values in the soils and the air surrounding the Kestanol granitoid plutons are anomalously high of 9200 nGy/h, and over the Kestanol granitoid, the value is 880 nGy/h. Besides the natural heat flow conveyed to the surface from the mantle and Aegean subduction zone, this granite also contributes considerable heat to the region. The heat flow values contribution by the Kestanol granites vary from 99 to 143 mW/m² that is similar to the heat flow values measured from the exploration boreholes drilled near Tuzla and estimated from the CPD deduced from aeromagnetic traverses over the western region of Turkey. Such high heat generating granites are the target for initiating enhanced geothermal systems, like the ones operating in Slutz in France. The Kestanol granitoid is covered by a sequence of Late Miocene volcanic rocks overlain by Pliocene sedimentary sequence. The estimated temperature of the granite at 2 km is about 90 °C and at 3 km depth, it is 120 °C. The Kestanol granitoid is under a

convenient NE-SW S_{hmax} , making it a suitable candidate to initiate the EGS project. A schematic section across NE-SW traverse (from NAF to Aegean trench) is presented in Figure 7.

The Aegean extensional tectonic fabric encloses Anatolide-Tauride and Sakarya continental plates, which collided in the Paleocene. The ophiolites and the blueschists of the Cretaceous were derived from the collision of the above two plates. The plutonic activity resulting from the post-Eocene–Oligocene collision event north of the suture zone marks the oldest magmatic event in this region. This magmatic activity migrated southwards, changing the composition from calc-alkalic to alkalic. The Quaternary volcanism appears to have resulted due to the lithospheric extension and decompressional melting associated with upwelling of the asthenosphere, which has resulted in Quaternary alkaline volcanism in the south central part of the Aegean extensional province (Dilek and Altunkaynak, 2009).

10. Conclusion

The Miocene Kestanol granitoid, a quartz monzonite intrusion, has an anomalous concentration of U, Th, and K and is one of the high heat generating granites located south of Çanakkale in the western Anatolian region. The Kestanol granitoid is a product of crustal melting and intruded into the older metamorphics and younger volcano-sedimentary sequence of post Miocene-Pliocene sequence. The presence of high-temperature geothermal




Figure 7. Interpretative tectonic cross section along a NNE-SSW-trending profile through the Africa-Eurasia convergence zone and the Aegean extensional province (modified after Dilek and Altunkaynak, 2009).

systems in this province, together with high-temperature bottom hole temperatures recorded from the exploratory drill hole and suitable temperature of the granite at 3 km depth and convenient stress fields, makes this granite a suitable candidate for initiating enhanced geothermal systems projects.

References

- Akal C (2013). Coeval Shoshonitic-ultrapotassic dyke emplacements within the Kestanol Pluton, Ezine – Biga Peninsula (NW Anatolia). *Turkish Journal of Earth Sciences* 22: 220-238.
- Akbaş B, Akdeniz N, Aksay A, Altun İE, Balcı V et al. (2011). Turkey Geology Map, General Directorate of Mineral Reserach and Exploration Publications. Ankara-Turkey. Erişim adresi: <http://yerbilimleri.mta.gov.tr>
- Akin U, Ulugergerli E, Kutlu S (2014). The assessment of the geothermal potential of Turkey by means of heat flow estimation. *Bulletin Mineral Resources Exploration* 149: 201-210.
- Akkuş İ, Akıllı H, Ceyhan S, Dilemre A, Tekin Z (2001). Türkiye Jeotermal Kaynaklar Envanteri, MTA Genel Müdürlüğü Envanter Serisi, Ankara, 201.
- Aknin U, Ulugergerli EU, Kutlu S (2014). The assessment of the geothermal potential of Turkey by means of heat flow estimation. *Bulletin Mineral Resources Exploration* 149: 201-210.
- Aldanmaz E, Pearce J, Thirlwall MF, Mitchell J (2000). Petrogenetic evolution of late Cenozoic, post-collision volcanism in western Anatolia, Turkey. *Journal of Volcanology and Geothermal Research* 102: 67-95.
- Aldanmaz E (2006). Mineral-chemical constraints on the Miocene calc-alkaline and shoshonitic volcanic rocks of Western Turkey: disequilibrium phenocryst assemblages as indicators of magma storage and mixing conditions. *Turkish Journal of Earth Science* 15: 47-73.
- Altunkaynak Ş, Dilek Y (2006). Timing and nature of post-collisional volcanism in Western Anatolia and geodynamic implications. In: Dilek Y, Pavlides S, editors. *Post-Collisional Tectonics and Magmatism of the Eastern Mediterranean Region*. Boulder, CO, USA: Geological Society of America Special Papers, pp. 321-351.
- Altunkaynak Ş, Dilek Y (2013). Eocene mafic volcanism in northern Anatolia: its causes and mantle sources in the absence of active subduction. *International Geological Reviews* 55: 1641-1659.
- Altunkaynak S, Dilek Y, Genc SC, Sunal G, Gertisser R et al. (2012a). Spatial, temporal and geochemical evolution of Oligo–Miocene granitoid magmatism in western Anatolia, Turkey. *Gondwana Research* 21: 961-986.
- Altunkaynak Ş, Genc ŞC (2008). Petrogenesis and time-progressive evolution of the Cenozoic continental volcanism in the Biga Peninsula, NW Anatolia (Turkey). *Lithos* 102: 316-340.
- Altunkaynak S, Rogerw NM, Kelley SP (2010). Causes and effects of geochemical variations in Late Cenozoic volcanism of the Foca volcanic centre, NW Anatolia, Turkey. *International Geological Reviews* 55: 579-607.
- Altunkaynak S, Sunal G, Aldanmaz E, Genc SC, Dilek Y et al. (2012b). Eocene granitic magmatism in NW Anatolia (Turkey) revisited: new implications from comparative zircon SHRIMP U-Pb and ⁴⁰Ar-³⁹Ar geochronology and isotope geochemistry on magma genesis and emplacement. *Lithos* 155: 289-309.
- Altunkaynak Ş, Yılmaz Y (1998). The Kozak magmatic complex; western Anatolia. *Journal of Volcanology and Geothermal Research* 85: 211-231.
- Altunkaynak S (2007). Collision-driven slab breakoff magmatism in northwestern Anatolia, Turkey. *The Journal of Geology* 115: 63-82.
- Angı OS, Yavuz O, Yalçın T, Çiftçi E (2016). Mineralogy-induced radiological aspects with characterization of commercial granites exploited in Turkey. *Bulletin of Engineering Geology and the Environment* 76: 507-522. doi: 10.1007/s10064-016-0894-2
- Arik F, Aydın U (2011). Mineralogical and petrographical characteristics of the Aladag skarn. *Scientific Research and Essays* 6: 592-606.
- Baba A, Uzelli T, Sozbilir H (2021). Distribution of geothermal arsenic in relation to geothermal play types: A global review and case study from the Anatolian plate (Turkey). *Journal of Hazardous Materials* 414: 125510. doi: 10.1016/j.jhazmat.2021.125510
- Baba A, Ertekin C (2007). Determination of the source and age of the geothermal fluid and its effects on groundwater resources in Kestanol (Çanakkale-Turkey). *GQ07: Securing Groundwater Quality in Urban and Industrial Environments* (Proc. 6th International Groundwater Quality Conference held in Fremantle, Western Australia, 2-7 December 2007).
- Baba A, Özcan H, Deniz O (2005). Environmental Impact by Spill of Geothermal fluids at the Geothermal Field of Tuzla, Çanakkale-Turkey. *Proceed. World Geothermal Congress-2005, Turkey*.
- Baba A, Deniz O, Özcan H, Erees S, Cetiner SZ (2008). Geochemical and radionuclide profile of Tuzla geothermal field, Turkey. *Environmental Monitoring Assessment* 145: 361-374.
- Barka AA (1992). The North Anatolian Fault zone. *Annals of Tectonics* 6: 164-195.
- Bingol E, Delaloye M, Ataman G (1982). Granitic intrusions in western Anatolia: a contribution to the geodynamic study of this area. *Eclogae Geologicae Helvetiae* 75: 437-446. doi: [10.5169/seals-165237](https://doi.org/10.5169/seals-165237)

- Black KN (2012). Geochemical and Geochronological Relationships between Granitoid Plutons of the Biga Peninsula, NW Turkey. M.Sc. Thesis, University of Texas at Austin.
- Bozkurt E (2001). Neotectonics of Turkey - a synthesis. *Geodinamica Acta* 14: 3-30.
- Boztuğ D, Harlavan Y, Jonckheere R, Can I, Sarı R (2009). Geochemistry and K-Ar cooling ages of the Ilica, Cataldağ (Balıkesir) and Kozak (İzmir) granitoids, west Anatolia, Turkey. *Geological Journal* 44: 79-103.
- Canbaz B, Fu sun Cam N, Yapraka G, Candan O (2010). Natural radioactivity (^{226}Ra , ^{232}Th and ^{40}K) and assessment of radiological hazards in the Kestanbol granitoids, Turkey. *Radiation Protection Dosimetry* 141 (2): 192-198. doi:10.1093/rpd/ncq165
- Cermak V, Huckenholz HG, Rybach L, Schmid R (1982). Radioactive heat generation in rocks. In: K. Hellwege (Editor), *Landolt-Bornstein numerical data and functional relationships in science and technology. New Series, Group V. Geophysics and Space Research*. Springer, Berlin, Heidelberg, New York, pp. 433-481.
- Delaloye M, Bingöl E (2000). Granitoids from western and Northwestern Anatolia: geochemistry and modeling of geodynamic evolution. *International Geology Review* 42: 241-268.
- Demirsoy N, Basaran CH, Sandalcı S (2018). Historical Kestanbol Hot Springs: "The water that resurrects" *Lokman Hekim Dergisi* 8 (1): 23-32.
- Dilek Y, Altunkaynak Ş (2007). Cenozoic crustal evolution and mantle dynamics of post-collisional magmatism in western Anatolia. *International Geological Reviews* 49: 431-453.
- Dilek Y, Altunkaynak Ş (2010). Geochemistry of Neogene-Quaternary alkaline volcanism in western Anatolia, Turkey, and implications for the Aegean mantle. *International Geological Reviews* 52: 631-655.
- Dilek Y, Altunkaynak Ş (2009). Geochemical and temporal evolution of Cenozoic magmatism in western Turkey: Mantle response to collision, slab breakoff, and lithospheric tearing in an orogenic belt., In *Geodynamics of collision and collapse at the Africa-Arabia-Eurasia Subduction Zone.*, Londra: The Geological Society of London, Special Publication, 311, pp. 213-233.
- Dilek Y, Altunkaynak Ş, Öner Z (2009). Syn-extensional granitoids in the Menderes core complex and the late Cenozoic extensional tectonics of the Aegean province. *Geological Society, London. Special Publications* 321: 197-223.
- Eckstein Y (1978). Review of heat flow data from the eastern Mediterranean region. *Pure and Applied Geophysics* 117: 150-159.
- Emre Ö, Duman TY, Özalp S, Elmacı H, Olgun Ş et al. (2013). Açıklamalı Türkiye Diri Fay Haritası. Ölçek 1:1.250.000, Maden Tetkik ve Arama Genel Müdürlüğü, Özel Yayın Serisi-30, Ankara-Türkiye. ISBN: 978-605-5310-56-1
- Erickson AJ, Simmons MG, Ryan WBF (1976). Review of Heat Flow from the Mediterranean and Aegean Seas, in *Int. Symp. on the Struc. Hist. of the Medit. Basin, Split, Yugoslavia, Oct. 1976, Proc.*, pp. 263-280 (ed. by B. Biju-Duval and L. Montadert), (Editions Technip, Paris, 1977).
- Erkan K (2015). Geothermal investigations in western Anatolia using equilibrium temperatures from shallow boreholes. *Solid Earth* 6: 103-113.
- Erkul F (2010). Tectonic significance of synextensional ductile shear zones within the Early Miocene Alacamdag granites, northwestern Turkey. *Geological Magazine* 147: 611-637.
- Erkul F, Tatar Erkul S, Ersoy Y, Uysal İ, Klotzli U (2013). Petrology, mineral chemistry and Sr-Nd-Pb isotopic compositions of granitoids in the central Menderes Core Complex: constraints on the evolution of Aegean lithosphere slab. *Lithos* 180-181: 74-91.
- Ersoy YE, Helvacı C, Palmer MR (2009). Petrogenesis of the Neogene volcanic units in the NE-SW-trending basins in western Anatolia, Turkey. *Contributions to Mineralogy and Petrology* 163: 379-401.
- Fytikas M, Innocenti E, Manetti P, Mazzuoli R, Peccerillo A et al. (1984). Tertiary to Quaternary evolution of volcanism in the Aegean region. *Geological Society London* 17: 687-699.
- Gulec N (1991). Crust-mantle interaction in western Turkey: implications from Sr and Nd isotope geochemistry of Tertiary and Quaternary volcanics. *Geological Magazine* 23: 417-435.
- Harris NBW, Kelley S, Okay AI (1994). Post-collisional magmatism and tectonics in northwest Anatolia. *Contributions to Mineralogy and Petrology* 117: 241-252.
- Hasozbek A, Satır M, Erdoğan B, Akay E, Siebel W (2010). Early Miocene post-collisional magmatism in NW Turkey: geochemical and geochronological constraints. *International Geological Reviews* 53: 1098-1119.
- Jackson J (1994). Active tectonics of the Aegean region. *Annual Reviews, Earth and Planetary Sciences* 22: 239-71.
- Jolivet L, Menant A, Sternai P, Rabillard A, Arbaret L et al. (2015). The geological signature of a slab tear below the Aegean. *Tectonophysics* 659: 166-182. doi: 110.1016/j.tecto.2015.1008.100
- Karacık Z, Yılmaz Y (1998). Geology of ignimbrites and the associated volcano-plutonic complex of the Ezine area, northwestern Anatolia. *Journal of Volcanology and Geothermal Research* 85: 251-264.
- Karamandereci IH, Öngür T (1974). The report of gradient wells finished of Tuzla Çanakkale geothermal field. MTA report, no: 5524, Ankara.
- Karat HI, Aydın I (2004). Report on Preparation of the Curie Isotherm Depth Map of Turkey, Unpublished Report no. 10638, MTA, Ankara (in Turkish).
- Koçyiğit A, Özacar A (2003). Extensional neotectonic regime through the NE edge of the outer İsparta angle, SW Turkey: New field and seismic data. *Turkish Journal of Earth Sciences* 12: 67-90.
- Koçyiğit A, Yusufoglu H, Bozkurt E (1999). Evidence from the Gediz Graben for episodic two-stage extension in western Turkey. *Journal of the Geological Society, London* 156: 605-616.

- Koprubaşı N, Aldanmaz E (2004). Geochemical constraints on the petrogenesis of Cenozoic I-type granitoids in Northwest Anatolia, Turkey: evidence for magma generation by lithospheric delamination in a post-collisional setting. *International Geological Reviews* 46: 705-729.
- Lachenbruch AH (1968). Preliminary geothermal model of the Sierra Nevada. *Journal of Geophysical Research* 73: 6977-6989.
- McKenzie D (1972). Active tectonics of the Mediterranean Region. *Geophysical Journal of the Royal Astronomical Society of London* 30: 109-185.
- Mützenbergs S (1997). Nature and origin of the thermal springs in the Tuzla area, Western Anatolia, Turkey. In: *The Marmara Poly-Project* (ed. by Schindler, C. & Pfister, M.), vdf hochschulverlag AG an der ETH, Zurich, 301-317.
- Ocakoglu N, Demirbag E, Kuscı I (2004). Neotectonic structures in the area offshore of Alacat, Döğanbey and Kusadas (western Turkey): evidence of strike-slip faulting in the Aegean extensional province. *Tectonophysics* 391: 67- 83.
- Okay AI, Satır M (2000). Coeval plutonism and metamorphism in the latest Oligocene metamorphic core complex in northwest Turkey. *Geological Magazine* 137: 495-516.
- Okay AI, Satır M (2006). Geochronology of Eocene plutonism and metamorphism in northwest Turkey: evidence for a possible magmatic arc. *Geodinamica Acta* 19: 251-266.
- Örgün Y, Altınsoy N, Şahin SY, Güngör Y, Gültekin AH et al. (2007). Natural and anthropogenic radionuclides in rocks and beach sands from Ezine region (Çanakkale), Western Anatolia, Turkey. *Applied Radiation and Isotopes* 65: 739-747.
- Papadopoulos A, Altunkaynak S, Koroneos A, Ünal A, Kamacı O (2016). Distribution of natural radioactivity and assessment of radioactive dose of the western Anatolian pluton, Turkey. *Turkish Journal Earth Sciences* 25: 434-455.
- Rabai S, Philip H, Taboada A (1992). Modern tectonic stress field in the Mediterranean region: evidence for variation in stress directions at different scales. *Geophysical Journal International* 10: 106-140.
- Rybach L (1976). Radioactive Heat Production: A Physical Property Determined by the Chemistry. In: R.G.I. Strens (Editor), *The Physical and Chemistry of Minerals and Rocks*. Wiley-Interscience Publication, New York, USA, pp. 245-276.
- Şahin SY, Örgün Y, Güngör Y, Göker AF, Gültekin AH et al. (2010). Mineral and Whole-rock Geochemistry of the Kestanol Granitoid (Ezine-Çanakkale) and its Mafic Microgranular Enclaves in Northwestern Anatolia: Evidence of Felsic and Mafic Magma Interaction. *Turkish Journal of Earth Sciences* 19: 101-122.
- Sener M, Gevrek AI (2000). Distribution and significance of hydrothermal alteration minerals in the Tuzla hydrothermal system, Çanakkale, Turkey. *Journal of Volcanology and Geothermal Research* 96: 215-228.
- Şengör AMC, Yılmaz Y (1981). Tethyan evolution of Turkey: a plate tectonic approach. *Tectonophysics* 75: 181-241.
- Şengör AMC, Dyer JMN (1979). Neotectonic provinces of the Tethyan orogenic belt of the eastern Mediterranean; variations in tectonic style and magmatism in a collision zone, *Eos, Transactions, American Geophysical Union*, 60 (18), p. 390, 1979. Meeting: American Geophysical Union, 1979 spring annual meeting, Washington, D.C., United States.
- Şengör AMC (1980). Türkiye'nin Neotektoniğinin Esasları: TJK Konferanslar Dizisi, s. 40. Ankara.
- Serpen U, Aksoy N, Ongur T, Korkmaz ED (2009). Geothermal energy in Turkey: 2008 update. *Geothermics* 38: 227-237.
- Tatar Erkul S, (2012). Petrogenetic evolution of the Early Miocene Alacamdag volcano-plutonic complex, northwestern Turkey: implications for the geodynamic framework of the Aegean region. *International Journal of Earth Science* 10: 197-219.
- Tatar Erkul S, Erkul F (2012). Magma interaction processes in synextensional granitoids: the Tertiary Menderes Metamorphic Core Complex, western Turkey. *Lithos* 142-143: 16-33.
- Taymaz T, Jackson J, McKenzie D (1991). Active tectonics of the north and central Aegean Sea. *Geophysical Journal International* 106: 433-490.
- Tezcan AK, Turgay MI (1991). Heat flow and temperature distribution in Turkey, edited by: Cermak V, Haenal R, and Zui V *Geothermal atlas of Europe*, Herman Haack Verlag, Gotha, Germany, pp 84-85.
- Tezel T, Shibutani T, Kaypak B (2013). Crustal thickness of Turkey determined by receiver function. *Journal of Asian Earth Sciences* 75: 36-45.
- Uğur A, Emin UU, Semih K (2014). The Assessment of Geothermal Potential of Turkey by means of heat flow estimation. *Bulletin Mineral Resources Exploration* 149: 201-210.
- Uzel B, Sozbilir H, Ozkaymak C (2012). Neotectonic Evolution of an Actively Growing Superimposed Basin in Western Anatolia: The Inner Bay of İzmir, Turkey. *Turkish Journal Earth Sciences* 4: 439 - 471.
- Vernekar AD (1975). A calculation of normal temperature at the earth's surface. *Journal of the Atmospheric Sciences* 32: 2067-2081.
- Yılmaz S, Örgün Y, Güngör Y, Göker AF, Gültekin AH et. al. (2010). Mineral and whole-rock geochemistry of the Kestanol granitoid (Ezine-Çanakkale) and its mafic microgranular enclaves in northwestern anatolia: evidence of felsic and mafic magma interaction. *Turkish Journal of Earth Sciences* 19: 101-122.
- Yılmaz Y (1989). An approach to the origin of young volcanic rocks of western Turkey. In: Şengör AMC, editor. *Tectonic Evolution of the Tethyan Region*. The Hague, the Netherlands: Kluwer Academic, pp. 159-189.
- Yılmaz Y, Genc SC, Karacık Z, Altunkaynak S (2001). Two contrasting magmatic associations of NW Anatolia and their tectonic significance. *Journal of Geodynamics* 31: 243-271.
- Yılmaz Y (1997). *Geology of Western Anatolia: active tectonics of northwestern Anatolia*. The Marmara Poly Project. VDF, Hochschulverlag Ag An Der ETH, Zurich 1-20.