

**Anti-inflammatory photodynamic therapy potential of polyoxyethylene-substituted
perylene diimide, nitrocatechol and azo dye**

Özgül HAKLI¹, Kasım OCAKOĞLU², Furkan AYAZ^{3,*}

¹ Department of Chemistry, Faculty of Science, Muğla Sıtkı Koçman University,
Muğla, Turkey

²Department of Engineering Fundamental Sciences, Faculty of Engineering, Tarsus
University, Mersin, Turkey

³Department of Biotechnology, Faculty of Arts and Science, Mersin University, Mersin,
Turkey

*Correspondence: furkanayaz@mersin.edu.tr

ORCIDs

Özgül HAKLI: <https://orcid.org/0000-0001-9316-2444>

Kasım OCAKOĞLU: <https://orcid.org/0000-0003-2807-0425>

Furkan AYAZ: <https://orcid.org/0000-0003-0271-0594>

Cite as: Haklı Ö, Ocakoğlu K, Ayaz F. Anti-inflammatory photodynamic therapy
potential of polyoxyethylene-substituted perylene diimide, nitrocatechol and azo dye.
Turkish Journal of Chemistry. doi: 10.3906/kim-2104-8

Abstract: Photodynamic therapy (PDT) applications enable light-controlled activation of drug candidates instead of their constitutive activities to prevent undesired side effects associated with their constant activities. A specific wavelength of light is utilized to enable electron mobility in the chemical structure which results in differential activities that may alter cell viability and cellular functions. Canonical photodynamic therapy applications mostly focus on cytotoxicity-based anti-microbial and anti-cancer properties of the PDT agents. In this study, we focused on subtoxic concentrations of three different molecules containing polyoxyethylene group and examined their anti-inflammatory activities on stimulated mammalian macrophages. Stimulated macrophages produce pro-inflammatory cytokines TNF and IL6. In the presence of a light source, our PDT agents were activated for 5 and 10 min during their application to the macrophages. Based on the ELISA results, the compounds had anti-inflammatory PDT activities. Trypan blue staining results suggest that these derivatives exerted their activities without leading to cytotoxicity. Our results suggest non-canonical PDT applications of these derivatives that can alter cellular activities without leading to cell death.

Keywords: Inflammation, perylene, photodynamic therapy, macrophages, anti-inflammatory compounds

1. Introduction

Polyoxyethylene side chains are preferred in molecules for various reasons due to their greater conformational flexibility, increased polarity and increased solubility [1]. The molecules described here possess polyoxyethylene substituent groups (Figure 1). Perylene diimide derivative **1** was synthesized by using the triethylene glycol

monomethyl ether derivative polyoxy group substituted on a benzene ring. The polyoxyethylene side chain increases the polarity and solubility of the molecule. Perylene-3,4,9,10-tetracarboxylic diimide derivatives containing a π -conjugated system with a perylene aromatic core have gathered a substantial interest in the field as fluorophores. Both symmetrical and unsymmetrical perylene diimides (PDIs), have been chemically and photodynamically stable, are easy to synthesize and characterize, have high solubility in organic solvents, have convenient excitation and emission spectra in the visible region (400-450 nm B band, 500-700 nm Q band). PDIs have been widely used in broad areas. They have versatile electrical and optical utilizations. These are not limited but can be listed as semiconductors in photovoltaic systems, electrophotographic technologies, dye lasers, transistors, light-emitting diodes, and photorefractive thin films [2-5]. They have been also used in cancer theranostics due to their high thermostability, large π - π conjugated structure, superior photochemical properties, and high fluorescence quantum yields [6]. These molecules have been used as ligands for photodynamic therapy applications. They are known to inhibit the telomerase activity in cancer cells. These molecules' mechanism of action on cancer cells is through the production of reactive oxygen species upon receiving the visible light irradiation. These molecules' binding ability to G-quadruplex structures of the DNA has been an active area of investigation since this DNA structure is well known for telomerase activity. By targeting this region telomerase activity and therefore cancer cell proliferation will be eliminated [7]. Although active oxygen species-based functional properties of PDIs as a PDT agent have been studied extensively, their subtoxic and non-canonical PDT potentials have not been investigated yet [8-12]. Azo compound (molecule **2**) containing polyoxy group was obtained by using a chlorine

derivative of triethylene glycol and aniline derivative. The polyoxyethylene group as a substituent was bonded to the oxygen atom in the hydroxyl group in the benzene ring. Azo dyes contain the (-N = N-) part in their structure. This structure is conjugated with two, different or identical, mono- or polycyclic aromatic systems. They have found wide application in the pharmaceutical, cosmetic, food, dyeing/textile industry and analytical chemistry due to their specific physicochemical properties and biological activities. There are numerous biological activities that make them medically attractive compounds [13-14]. Molecule **3** containing polyoxyethylene units was obtained by using chlorine derivative of triethylene glycol and 4-nitro catechol. Polyoxyethylene units are attached to the oxygen atom in both hydroxyl groups as substituents. 4-Nitrocatechol and other catechol derivatives are known to be active against the intestinal bacterial panel and nitrophenol derivatives have antimicrobial activity [15].

In this study, we focused on subtoxic concentrations of PDI derivatives as potential PDT agents to suppress inflammatory responses created by mammalian macrophages. These PDI derivatives were chosen due to their photodynamic activity potentials and in this study, we aimed to investigate the differences in their efficiencies for their possible utilization in the future. Macrophages are chosen for this study due to their pro-inflammatory TNF, IL6, IL1, IL12, and GMCSF production potentials. They can regulate the immune system by these signaling molecules. Therefore, to decipher the activity of a possible immunomodulatory compound initial screening on these cells gives crucial information. Moreover, these cells can phagocytose cell debris and infectious agents as well as their exo and endotoxins to further present them to the T cells of the adaptive immunity. Together with cytokine production, this presentation-based activation of the T cells makes macrophages an important player in the

determination of the type and strength of the immune response against a certain danger to the body [16, 17].

Hence to investigate the anti-inflammatory PDT potentials of PDI derivatives they were tested on activated mammalian macrophages. Based on TNF and IL6 ELISA results, these derivatives had anti-inflammatory activities and their PDT potential was differential based on structural differences.

2. Materials and Methods

2.1. Synthesis of N,N'-Bis (4-{2-[2-(2-methoxyethoxy ethoxy)ethoxy} phenyl)-

3,4:9,10-perylene tetracarboxydiimide (1)

The synthesis procedure was illustrated in our previous work [18, 19]. Perylene-3,4:9,10- tetracarboxylic acid bisanhydride (0.69 mmol, 270 mg), 4-{2-[2-(2-methoxyethoxy) ethoxy] ethoxy} aniline (1.5 mmol, 382.5 mg) and imidazole (5 g) were heated at 140 °C for 4.5 hours under inert atmosphere. Then 2N HCl (200 mL) was added to the reaction solution and the resulting mixture was stirred for 1 h at room temperature. It was extracted with chloroform (600 mL). The organic phase evaporated under vacuum and crude product was purified by column chromatography (neutral alumina; CH₂Cl₂-MeOH, 10:1). Yield: 62%. FT-IR (cm⁻¹): 2922-2867, 1704-1663, 1595-1512, 1455-1404, 1361, 1299- 1255, 1178, 1124. ¹H NMR (CDCl₃, 400 MHz): δ (ppm) = 8.68-8.59 (q, 8H, ArH), 7.20-7.02 (q, 8H, ArH), 4.15 (t, 4H, ArOCH₂-), 3.84 (t, 4H, ArOCH₂CH₂), 3.7 (t, 4H, -OCH₂-CH₂OCH₃), 3.63 3.61 (m, 8H, - OCH₂CH₂O-), 3.5 (t, 4H, -CH₂OCH₃), 3.33 (s, 6H, -OCH₃), C₅₀H₄₆N₂O₁₂.

2.2. Synthesis of 2{2[2(2-methoxyethoxy)ethoxy]ethoxy}-5-[(E)-(4

nitrophenyl)diazanyl]benzaldehyde (2)

1-(3-formyl-4-hydroxyphenylazo)-4-nitrobenzene (2) was synthesized before according to literature [20,21]. Chloro-2-[2-(2-methoxyethoxy)ethoxy]ethane (2.8 mmol) was added to a mixture of 1-(3-formyl-4-hydroxy- phenylazo)-4 nitrobenzene (6.4mmol)

and K_2CO_3 (12.8 mmol) in DMF (20mL) under inert atmosphere. The mixture was heated at 140 °C for 19h and then refluxed for 2h. After concentrating, purification was done by column chromatography (Silica, $CHCl_3$:MeOH, 10:0.5). Yield: 85%. 1H NMR ($CDCl_3$, 400 MHz): δ (ppm) = 10.05 (s, 1H), 8.39(d, 2H), 8.27(t, 1H), 8.22(t, 1H), 8.01 (d, 2H), 7.14(d, 1H), 4.3(t, 2H), 3.9(t, 2H),3.7(t, 2H),3.6(m, 4H), 3.5(t, 2H),3.33(s, 3H). $C_{20}H_{23}N_3O_7$.

2.3. Synthesis of 1,2-di{2-[2-(2-methoxyethoxy)ethoxy]ethoxy}-4-nitrobenzene(3)

The synthesis procedure was illustrated in our previous work [22]. Chloro-2-[2-(2-methoxyethoxy)ethoxy]ethane (2.34 g, 2.8 mmol) was added to a mixture of 4-nitrocatechol (1 g, 6.4 mmol) and K_2CO_3 (1.77 g, 12.8 mmol) in DMF (20 mL) under an inert atmosphere. The mixture was heated at 140 °C for 19 hours by stirring. And then it was refluxed for 2 hours. It was stopped after 2 hours. After DMF was removed by rotary evaporation, the resulting solution was purified through a column. The eluent was 4% methanol in $CHCl_3$. Yield: 85%. H^1 NMR ($CDCl_3$, 400 MHz): δ (ppm) = 7.8-6.9 (m, 3H, Ar-H), 4.2-4.1 (q, 4H, $ArOCH_2-$), 3.88-3.85 (m, 4H, $ArOCH_2CH_2-$), 3.72-3.69 (m, 4H, $-OCH_2CH_2OCH_3$), 3.64 3.58 (m, 8H, $-OCH_2CH_2O-$), 3.51-3.48 (m, 4H, $CH_2CH_2OCH_3$), 3.33 (s, 6H, $-OCH_3$); $C_{20}H_{33}NO_{10}$.

For the photodynamic activation Xenon Light 300 Watt/m² was used.

2.4. Stimulation of mammalian macrophages to test the anti-inflammatory PDT activities of compounds

These protocols were explained in detail in our previous studies [23-30]. Mammalian macrophages were RAW 264.7 mouse macrophage cells from ATCC. These cells were grown in RPMI 1640 media together with 10% fetal bovine serum, 1% antibiotics (100 μ g/ml penicillin and 100 μ g/ml streptomycin), and sodium pyruvate. Cell incubations were done in 37 °C 5% CO_2 incubator [23,24,26-30,31]. 10^6 macrophages were plated into each well of 24 well plates in 1mL final volume. They were let adhere to the bottom of the plates after overnight incubation before their activation. For stimulations, sterile DMSO was added into the negative control wells, 1 μ g/mL of LPS (1mg/mL,

Enzo Life Sciences, Salmonella Minnesota R595), and sterile DMSO was added into the positive control wells and 1ug/mL and 10ug/mL of the compounds molecules (Figure 1) was put into the appropriate wells with or without 1 ug/mL LPS. These conditions were created as triplicate setups: Dark conditions, 5 min Xenon light exposure, and 10 min Xenon light exposure. Before 24 hours of incubation, light exposure procedures were conducted right after the addition of the stimulants and compounds. Afterward, the plates were put into the incubator for 24 hours. After the incubation period, the supernatants were collected and kept at -80°C for ELISA. Trypan blue staining with a hemocytometer was done to assess the cell viability. TNF α and IL6 BD ELISA kits were purchased and ELISA protocols were followed by using the manufacturer's instructions to determine the changes in the production of these pro-inflammatory cytokines [23,24,26-30,32].

Statistical Analysis: 3 biologically independent data sets were combined to do a student t-test by using GraphPad Prism Software 5 [23, 24, 26-30].

3. Results

3.1. Compounds lacked immunostimulatory activities on mammalian macrophages

Compounds were tested on unactivated mammalian macrophages to determine their possible intrinsic stimulatory activities on unstimulated macrophages. They did not exert the production of pro-inflammatory TNF and IL6 cytokines by the macrophages in the absence of LPS as an activator. Moreover, having dark and light-induced conditions did not change the results and these compounds lacked immunostimulatory activity under all conditions (Figure 2 and Figure3).

3.2. Compounds had differential anti-inflammatory PDT potentials on activated mammalian macrophages based on TNF and IL6 production levels

In order to test the anti-inflammatory potentials of compounds, they were also tested on mammalian macrophages in the presence of LPS as a stimulant. Under dark conditions other than compound **1**, all of the derivatives had anti-inflammatory activities on macrophages in terms of TNF production. TNF was not detected in the supernatants of the groups that were treated with compounds **2** and **3**, whereas compound **1** did not affect TNF production levels in dark conditions. Upon light-induced activation, compound **1** gained anti-inflammatory activity and completely suppressed the production of TNF both after 5 min and 10 min of light exposure. Compound **2** and **3** retained their anti-inflammatory activity similar to dark conditions based on TNF production. These results suggest that all of the compounds have anti-inflammatory activity on the activated macrophages by reducing the TNF production but compound **1** had PDT potential. Compounds **2** and **3** were constitutively active since they were able to suppress TNF production independent of light-induced activation (Figure 3).

IL6 was another cytokine that we measured from activated macrophages. Based on our results, compound **1** lacked the ability to alter IL6 production both in dark and light conditions. Whereas, compounds **2** and **3** gained a strong anti-inflammatory activity only after light exposure. These compounds (**2** and **3**) could not change IL6 production by activated macrophages in dark conditions. Only their (**2** and **3**) higher concentrations showed PDT potential. Compound **3**'s higher concentrations lead to complete knockout of IL6 production by activated macrophages after 5 and 10 min of Xenon light exposure. Whereas compound **2** gained this activity on IL6 production after 10 min of Xenon light exposure only with its high concentrations.

These results imply that compound **1** would not be useful in anti-inflammatory PDT applications where IL6 production is a target but it will be useful for targeting TNF while sparing IL6 production by macrophages. Compounds **2** and **3** can constitutively block TNF production independent of light exposure but they exerted an effective anti-inflammatory PDT potential to suppress IL6 production with their higher concentrations only after light exposure (Figure 2 and Figure3).

3.3. Compounds did not have cytotoxic effects on mammalian macrophages

In this study, we focused on non-toxic concentrations of these agents and we aimed to determine their biological activities at subtoxic concentrations both at dark and light-induced conditions. Trypan blue staining was conducted to confirm that these compounds were non-toxic at the concentrations that they were used. Our results suggest that they did not have a substantial effect on the cell viability both in dark and light conditions.

4. Conclusion

Compounds have been studied for their photovoltaic applications in solar cells. These molecules also possess biological activities. Compounds exerted anti-cancer activity and some studies also suggest their possible utilization for tumor cell imaging due to their ability to interact with DNA [9-12, 32]. Moreover, polyoxyethylene groups have been used as prodrug and drug delivery molecules in various applications. They increase the uptake of the drug molecules by the cells and tissues while enabling more controlled delivery of the main active ingredients of the drug formulations. Due to their biocompatible nature as well as the ability to increase the bioavailability and solubility of drug molecules, it was designed a perylene diimide derivative substituted with polyoxyethylene groups [33-37]. During the production process, we obtained two benzene derivatives with polyoxyethylene substitutions.

In this study, we examined these molecules' anti-inflammatory PDT potential on mammalian macrophages. Studies mostly focus on a cytotoxic aspect of PDT and present data on anti-tumor, anti-fungal, or anti-bacterial properties of PDT agents [38-44].

In this study, we utilized subtoxic concentrations of the agents which were confirmed by Trypan Blue staining (Figure 4). Intermediate compounds had similar activities on TNF production levels of activated macrophages and they were strong anti-inflammatory agents even in dark conditions. They kept this property and knocked TNF production out by activated macrophages after 5 min and 10 min of light exposure as well. On the other hand, having perylene diimide in the structure enabled a PDT potential since PDI derivative was inert on stimulated macrophages for the production of TNF in the dark and gained strong anti-inflammatory property by completely knocking out TNF production after 5 min and 10 min of Xenon light exposure. These results suggest that having perylene diimide in the molecular formula enabled PDT application potential.

IL6 was another pro-inflammatory cytokine that we examined in this study. Our results suggest that these compounds were not able to alter IL6 production by stimulated macrophages which could be an advantage in situations where TNF production should be suppressed while sparing IL6 production by the macrophages. Moreover, the intermediate products suppressed IL6 production at their higher concentrations especially with light exposure. Photodynamic therapy presents opportunities for more controlled and localized activation of drug molecules to increase their efficiency and decrease their possible side effects. So far, most of the studies utilize photosensitizers and different dye molecules as PDT agents they mostly focus on their cytotoxic capacities [38-44].

The uV spectrums of the studied compounds are given in Figure 5. Compound 1 has characteristic absorption peaks at 455 nm, 492 nm and 527 nm corresponding to $\pi-\pi^*$

singlet transition. Compound 2 shows absorption peaks at 277 nm and 369 nm. Compound 3 gives absorption peaks at 302 nm and 348 nm.

Our results suggest that compound 1 was not as effective as compound 2 and 3 in terms of its anti-inflammatory activities. The reason behind this difference might be due to the differences in their structures and how it might affect their interactions with the signaling pathways that play important role in the inflammatory response. In our future studies, we are planning to focus on their possible effects on inflammatory pathways such as JNK, ERK and PI3K [45].

In this study, we aimed to focus on their subtoxic concentrations and determine possible PDT activities. The compounds that we are presenting in this study had anti-inflammatory PDT potential on activated mammalian macrophages. Further studies on their in vivo efficacy will bring out their potential utilization in the clinic to cope with inflammatory and autoimmune diseases.

Acknowledgment

We would like to thank Prof. Dr. Juan Anguita from CIC Biogune for his guidance and support.

References

1. Cormier RA, Gregg BA. Synthesis and characterization of liquid crystalline perylene diimides. *Chemistry of Materials* 1998; 10: 1309-1319.
2. Cheng H, Zhao Y, Xu H, Hu Y, Zhang L et al. Rapid and visual detection of protamine based on ionic self-assembly of a water soluble perylene diimide derivative. *Dyes and Pigments* 2020; 180: 108456. doi: 10.1016/j.dyepig.2020.108456
3. Dincalp H, Kızılok S, Haklı Birel O, Icli S. Synthesis and G-quadruplex binding study of a novel full visible absorbing perylene diimide dye. *Journal of Photochemistry and Photobiology A: Chemistry* 2012; 235: 40–48. doi:10.1016/j.jphotochem.2012.03.004
4. Yu Y-Q, Zhang HY, Chai YQ, Yuan R, Ying Zhuo Y. Sensitive electrochemiluminescent aptasensor based on perylene derivative sasanovel co-reaction accelerator for signal amplification. *Biosensors and Bioelectronics* 2016; 85: 8–15. doi: 10.1016/j.bios.2016.04.088
5. Georgiev NI, Saida AI, Toshkova RA, Tzoneva RD, Bojinov VB. A novel water-soluble perylenetetra-carboxylic diimide as a fluorescent pH probe: Chemosensing, biocompatibility and cell imaging. *Dyes and Pigments* 2019; 160: 28–36. doi: 10.1016/j.dyepig.2018.07.048.
6. Li H, Yue L, Wu M, Wu F. Self-assembly of methylene violet-conjugated perylene diimide with photodynamic/photothermal properties for DNA photocleavage and cancer treatment. *Colloids and Surfaces B: Biointerfaces* 2020; 196: 111351. doi: 10.1016/j.colsurfb.2020.111351

7. Dincalp H, Avcıbası N, İcli S. Spectral properties and G-quadruplex DNA binding selectivities of a series of unsymmetrical perylene diimides. *Journal of Photochemistry and Photobiology A: Chemistry* 2007; 185: 1–12. doi: 10.1016/j.jphotochem.2006.04.035
8. Wang J, Liu D, Zhu Y, Zhou S, Guan S. Supramolecular packing dominant photocatalytic oxidation and anticancer performance of PDI. *Applied Catalysis B: Environmental* 2018; 231: 251–261.
9. Hemalatha CN, Muthkumar VA. Application of 3D QSAR and docking studies in optimization of perylene diimides as anti-cancer agent. *Indian Journal of Pharmaceutical Education and Research* 2018; 52 (4): 666-675.
10. Sun P, Yuan P, Wang G, Deng W, Tian S et al. High density glycopolymers functionalized perylene diimide nanoparticles for tumor-targeted photoacoustic imaging and enhanced photothermal therapy. *Biomacromolecules* 2017; 18: 3375–3386. doi: 10.1021/acs.biomac.7b01029
11. Mazzitelli CL, Brodbelt JS, Kern JT, Rodriguez M, Kerwin SM. Evaluation of binding of perylene diimide and benzannulated perylene diimide ligands to DNA by electrospray ionization mass spectrometry. *Journal of the American Society for Mass Spectrometry* 2006; 17: 593–604. doi: 10.1016/j.jasms.2005.12.011
12. Zhao H, Hussain S, Liu X, Li S, Lv F et al. Design of an amphiphilic perylene diimide for optical recognition of anticancer drug through a chirality-Induced helical structure. *Chemistry a European Journal* 2019; 25: 9834 – 9839.
doi: 10.1002/chem.201901948
13. Weglarz-Tomczak E, Gorecki L. Azo dyes – biological activity and synthetic strategy. *CHEMIK* 2012; 66, 12: 1298-1307.

14. Gup R, Giziroglu E, Kirkan B. Synthesis and spectroscopic properties of new azo-dyes and azo-metal complexes derived from barbituric acid and aminoquinoline. *Dyes and Pigments* 2007; 73: 40-46. doi: 10.1016/j.dyepig.2005.10.005
15. Rachakonda S, Naaz F, Ali I, Krs P, Rao M B et al. Synthesis and antimicrobial activity of 1,2,3-triazole-tethered nitroguaiacol ethers. *Asian Journal of Pharmaceutical and Clinical Research* 2019; 12 (5): 329-334. doi: 10.22159/ajpcr.2019.v12i5.29603
16. Snyder RJ, Lantis J, Kirsner RS, Shah V, Molyneaux M et al. Macrophages: A review of their role in wound healing and their therapeutic use. *Wound Repair and Regeneration* 2016; 24: 613–629. doi: 10.1111/wrr.12444
17. He W, Kapate N, Shields CW, Mitragot S. Drug delivery to macrophages: a review of targeting drugs and drug carriers to macrophages for inflammatory diseases. *Advanced Drug Delivery Reviews* 2020; 165–166: 15–40. doi: 10.1016/j.addr.2019.12.001
18. Haklı Ö. Synthesis of perylene diimide derivatives and studies on photophysical properties of perylene diimides and some vitamin K3 derivatives: stability tests under concentrated sun light. PhD, İzmir, Turkey, 2006.
19. Kus M, Haklı Ö, Zafer C, Varlikli C, Demic S et al. Optical and electrochemical properties of polyether derivatives of perylene diimides adsorbed on nanocrystalline metal oxide films. *Organic Electronics* 2008; 9: 757–766. doi: 10.1016/j.orgel.2008.05.009

20. Birel O, Kavasoglu N, Kavasoglu AS, Dincalp H, Metin B. Fabrication and electrical characterization of Al/diazo compound containing polyoxy chain/p-Si device structure. *Physica B* 2013; 412: 64–69. doi: 10.1016/j.physb.2012.11.030
21. Dincalp H, Yavuz S, Haklı Ö, Zafer C, Özsoy C et al. Optical and photovoltaic properties of salicylaldehyde-based azo ligands. *Journal of Photochemistry and Photobiology A: Chemistry* 2010; 210: 8–16.
doi: 10.1016/j.jphotochem.2009.12.012
22. Haklı Birel O, Zafer C, Dincalp H, Aydın B, Can M. Highly soluble polyoxyethylene-perylene diimide: optical, electrochemical and photovoltaic studies. *Journal of Chemical Society of Pakistan* 2010; 33 (4): 2011.
23. Ayaz F, Demirbag B, Ocakoglu K. Immunoactive photosensitizers had photodynamic immunostimulatory and immunomodulatory effects on mammalian macrophages. *Photodiagnosis and Photodynamic Therapy* 2020; 32: 102034. doi: 10.1016/j.pdpdt.2020.102034
24. Gonul I, Demirbag B, Ocakoglu K, Ayaz F. Unique photodynamic antimicrobial Schiff bases and their copper complexes exert immunomodulatory activity on mammalian macrophages. *Journal of Coordination Chemistry* 2020; 73 (20-22): 2878-2888. doi: 10.1080/00958972.2020.1836362
25. Ayaz F, Colak SG, Ocakoglu K. Investigating the immunostimulatory and immunomodulatory effects of cis and trans isomers of ruthenium polypyridyl complexes on the mammalian macrophage-like cells. *ChemistrySelect* 2020; 5 (37): 11648-11653.

26. Önal HT, Yuzer A, Ince M, Ayaz F. Photo induced anti-inflammatory activities of a thiophene substituted subphthalocyanine derivative. *Photodiagnosis and Photodynamic Therapy* 2020; 30: 101701. doi: 10.1016/j.pdpdt.2020.101701
27. Ayaz F, Gonul I, Demirbag B, Ocakoglu K. Novel copper bearing schiff bases with photodynamic anti-inflammatory and anti-microbial activities. *Applied Biochemistry and Biotechnology* 2020; 191 (2): 716-727. doi: 10.1007/s12010-019-03223-7
28. Yüzer A, Ayaz F, Ince M. Immunomodulatory activities of zinc(II)phthalocyanine on the mammalian macrophages through p38 pathway: Potential ex vivo immunomodulatory PDT reagents. *Bioorganic Chemistry* 2019; 92: 103249. doi: 10.1016/j.bioorg.2019.103249
29. Ayaz F, Gonul İ, Demirbag B, Ocakoglu, K. Differential immunomodulatory activities of schiff base complexes depending on their metal conjugation. *Inflammation* 2019; 42 (5): 1878-1885. doi: 10.1007/s10753-019-01050-w
30. Ayaz F, Ugur N, Ocakoglu K, Ince M. Photo-induced anti-inflammatory activities of chloro substituted subphthalocyanines on the mammalian macrophage in vitro. *Photodiagnosis and Photodynamic Therapy* 2019; 25: 499-503. doi: 10.1016/j.pdpdt.2019.02.002
31. Ayaz F, Colak SG, Ocakoglu K. Investigating the immunostimulatory and immunomodulatory effects of cis and trans isomers of ruthenium polypyridyl complexes on the mammalian macrophage-like cells. *ChemistrySelect* 2020; 5 (37): 11648-11653.
32. Wang J, Liu D, Zhu Y, Zhou S, Guan S. Supramolecular packing dominant photocatalytic oxidation and anticancer performance of PDI. *Applied Catalysis*

- B: Environmental 2018; 231: 251-261.
33. Pardakhty A, Varshosaz J, Rouholamini A. In vitro study of polyoxyethylene alkyl ether niosomes for delivery of insulin. *International Journal of Pharmaceutics* 2007; 328 (2), 130-141. doi: 10.1016/j.ijpharm.2006.08.002
 34. Cornforth JW, Hart PD, Nicholls GA, Rees RJW, Stock JA. Antituberculous effects of certain surface-active polyoxyethylene ethers. *British Journal of Pharmacology and Chemotherapy* 1955; 10 (1): 73-86. doi: 10.1111/j.1476-5381.1955.tb00063.x
 35. Pizzo SV. Preparation in vivo properties and proposed clinical use of polyoxyethylene-modified tissue plasminogen activator and streptokinase. *Advanced Drug Delivery Reviews* 1991; 6 (2): 153-166.
 36. Andrews G P, Zhai H, Tipping S, Jones D S. Characterisation of the thermal, spectroscopic and drug dissolution properties of mefenamic acid and polyoxyethylene–polyoxypropylene solid dispersions. *Journal of Pharmaceutical Sciences* 2009; 98 (12): 4545-4556. doi: 10.1002/jps.21752
 37. Bonina FP, Puglia C, Barbuzzi T, de Caprariis P, Palagiano F et al. In vitro and in vivo evaluation of polyoxyethylene esters as dermal prodrugs of ketoprofen, naproxen and diclofenac. *European Journal of Pharmaceutical Sciences* 2001; 14 (2): 123-134. doi: 10.1016/s0928-0987(01)00163-4
 38. Zhang J, Jiang C, Longo JPF, Azevedo RB, Zhang H et al. An updated overview on the development of new photosensitizers for anticancer photodynamic therapy. *Acta Pharmaceutica Sinica B* 2018; 8 (2): 137-146. doi: 10.1016/j.apsb.2017.09.003
 39. Allison RR, Moghissi K. Oncologic photodynamic therapy: Clinical strategies

- that modulate mechanisms of action. *Photodiagnosis and Photodynamic Therapy* 2013; 10 (4): 331-341. doi: 10.1016/j.pdpdt.2013.03.011
40. Allison RR, Bagnato VS, Sibata CH. Future of oncologic photodynamic therapy. *Future Oncology* 2010; 6 (6): 929-940. doi: 10.2217/fon.10.51
41. Rahimi S, Janani M, Lotfi M, Shahi S, Aghbali A et al. A review of antibacterial agents in endodontic treatment. *Iranian Endodontic Journal* 2014; 9 (3): 161-168.
42. Maisch T, Szeimies RM, Jori G, Abels C. Antibacterial photodynamic therapy in dermatology. *Photochemical & Photobiological Sciences* 2004; 3 (10): 907-917. doi: 10.1039/b407622b
43. Donnelly RF, McCarron PA, Tunney MM. Antifungal photodynamic therapy. *Microbiological Research* 2008; 163 (1): 1-12. doi: 10.1016/j.micres.2007.08.001
44. Javed F, Samaranayake LP, Romanos GE. Treatment of oral fungal infections using antimicrobial photodynamic therapy: A systematic review of currently available evidence. *Photochemical & Photobiological Sciences* 2014; 13 (5): 726-734. doi: 10.1039/c3pp50426c
45. Newton K, Dixit VM. Signaling in innate immunity and inflammation. *Cold Spring Harbor perspectives in biology* 2012; 4 (3): 1-19. doi: 10.1101/cshperspect.a006049




Figure 1. Molecular structures of the compounds containing polyoxyethylene group.




Figure 2. TNF α ELISA results for 1 ug/mL and 10 ug/mL compound (1), compound (2), and compound (3) application to either untreated or 1 ug/mL LPS activated macrophages. Supernatants were analyzed after 24 hours of incubation, N = 3 (*P < 0.001, **P < 0.0005, ***P < 0.0001). The DMSO, as the solvent of the compounds, was added into the negative control and positive control wells in 10uL volume, 1 ug/mL LPS was used in positive control wells. This setup was used for dark conditions (A) as well as for 5 min (B) and 10 min (C) Xenon light exposure. The light exposure was done right after the addition of the compounds and afterward, the 24 hours incubation was started.




Figure 3. IL6 ELISA results for 1 ug/mL and 10 ug/mL compound (1), compound (2), and compound (3) application to either untreated or 1 ug/mL LPS activated macrophages. Supernatants were analyzed after 24 hours of incubation, N = 3 (*P < 0.001, ** P < 0.0005, ***P < 0.0001). The DMSO, as the solvent of the compounds, was added into the negative control and positive control wells in 10uL volume, 1 ug/mL LPS was used in positive control wells. This setup was used for dark conditions (A) as well as for 5 min (B) and 10 min (C) Xenon light exposure. The light exposure was done right after the addition of the compounds and afterward, the 24 hours incubation was started.




Figure 4. Cell viability results for 1 ug/mL and 10 ug/mL compound (1), compound (2), and compound (3) application to either untreated or 1 ug/mL LPS activated macrophages. The cell counting was done with Trypan blue after 24 hours of incubation, N = 3 (*P < 0.001, **P < 0.0005, ***P < 0.0001). The DMSO, as the solvent of the compounds, was added into the negative control and positive control wells in 10uL volume, 1 ug/mL LPS was used in positive control wells. This setup was used for dark conditions (A) as well as for 5 min (B) and 10 min (C) Xenon light exposure. The light exposure was done right after the addition of the compounds and afterward, the 24 hours incubation was started.




Figure 5. UV-Vis absorption spectra of compound (1), compound (2), and compound (3) in DMSO