

## **Determination of plant-specific retrotransposons in chicken**

**Levent Mercan<sup>1</sup>, Cihat Erdem Bulbul<sup>1</sup>, Fatih Bilgi<sup>1</sup>, Sevgi Marakli<sup>2\*</sup>**

<sup>1</sup>Department of Agricultural Biotechnology Department, Faculty of Agriculture,  
Ondokuz Mayıs University, Turkey.

<sup>2</sup>Department of Medical Services and Techniques, Sabuncuoglu Serefeddin Health  
Services Vocational School, Amasya University, Turkey.

Levent Mercan: 0000-0002-6790-1458

Cihat Erdem Bulbul: 0000-0002-8609-6239

Fatih Bilgi: 0000-0002-8239-2217

Sevgi Marakli: 0000-0001-5796-7819

## **Abstract**

Mobile genetic elements are also known as transposons, comprising a significant proportion of plant and animal genomes. Retrotransposons, a class of transposons, move copy and paste mechanism, and therefore cause genome obesity. Moreover, the transferability of retrotransposons within and between different kingdoms has provided valuable information for evolutionary relationships in addition to detailed knowledge about genomes. We identified barley-specific retrotransposons (*Nikita* and *Sukkula*) and determined polymorphism ratios of these retrotransposons in a chicken genome for the first time to gain new insights for retrotransposons found in different genomes. Transposon sequences belonging to chicken and barley genomes were also investigated to understand the evolutionary relationships. For this purpose, the samples from Turkish native chicken Gerze were analysed using a retrotransposon-based molecular marker, IRAP (Inter-Retrotransposon Amplified Polymorphism). As a result of the analyses, there was no polymorphism among the samples in *Sukkula* analysis while *Nikita* showed 0-60% polymorphism. Moreover, *in silico* analyses also presented that these two retrotransposons are closely related to other chicken retrotransposons. Obtaining results could offer a practical approach for chicken genome analyses, especially for the identification of genes to obtain desired traits.

**Keywords:** Gerze chicken, IRAP, Mobile genetic elements, *Nikita*, *Sukkula*

## **1. Introduction**

Transposable elements (TEs) constitute a major part of the genome in vertebrate genomes [1]. They are classified as Class I (RNA transposon or retrotransposon) and Class II (DNA transposon), moving through the “copy and paste” or “cut and paste” mechanisms, respectively [2]. TEs have been gained more importance due to role in genetic variation, evolution and contribution to speciation [3]. Therefore, there are many studies related to the identification of TEs and their roles in gene expression and gene expression regulations [3–5] because of the effect of retrotransposons by changing the dynamic functions of the host DNA [6].

Sequencing genomes belonging to bacteria, plants and animals have increased due to improving NGS technologies [7–9]. Chicken (*Gallus gallus*) is one of them with ~1.2 billion base pairs. The genome size of chicken is approximately one-third of most mammalian including the human genome. On the other hand, TEs content is remarkably low (~10%) in the chicken genome [10–12].

The chicken, the first genome sequenced domestic animal, is also one of the primary model organisms that bridge the evolutionary gap between mammals and other vertebrates [13]. During domestication, the chicken’s genetic structure has been changed and differentiated by natural and artificial selections [14]. The domestic chicken (*Gallus gallus domesticus*) originated from the red junglefowl (*Gallus gallus*) in Southeast Asia, the gray junglefowl (*Gallus sonneratii*) in Southwest India, and the Sri Lankan junglefowl (*Gallus lafayetii*) in Sri Lanka. The domestic chicken is transported from China and India to Europe via Russia and Anatolia via migration and trade routes [15]. Since Anatolia is a gateway of domestic chicken spreading to the rest of the world, it is crucial to study Turkish chicken breeds for retrotransposon analyses

on the chicken genome. Gerze chicken as one of the critical native genetic resources for Turkey is very suitable for such research.

The studied chicken-specific transposons in this presented study were CR1, ALVE and ART-CH groups belonging to ERV. The CR1 (chicken repeat 1) element, a short interspersed repetitive DNA element belonging to the non-LTR class [10]. Another retrotransposon ALVE (avian leukosis virus subgroup E- ALV-E) is characterised in *ev* loci. Furthermore, ART-CH (avian retrotransposon from the chicken genome) is found in 50 genomic copies in the chicken genome, showing homology to the avian leucosis and sarcoma virus (ALSV) [16].

Many studies determine specific retrotransposons in different genomes, reporting evolutionary relationships among species [17,18]. In this study, we aimed to identify barley-specific retrotransposons (*Nikita* and *Sukkula*) via IRAP (Inter-Retrotransposon Amplified Polymorphism) molecular marker technique in the chicken genome for the first time. Moreover, *in silico* analyses were also performed to figure out the evolutionary relationships between *Nikita-Sukkula* and chicken-specific retrotransposons.

## **2. Material and methods**

### **2.1. Obtaining samples and genomic DNA isolation**

In this study, the samples from Gerze chicken population under conservation of the Republic of Turkey Ministry of Agriculture and Forestry Gerze Directorate were used for *Nikita* and *Sukkula* retrotransposon movements. Ondokuz Mayıs University Animal Experiments Local Ethics Committee permitted this study with 06.11.2012/1 number. For this purpose, blood samples from 9 Gerze chickens were collected and genomic

DNAs (gDNAs) from samples were isolated by using salting-out method [19]. Qualitative and quantitative of gDNAs were evaluated on 1% agarose gel and a NanoDrop® spectrophotometer (Thermo, USA), respectively. The experiment was replicated three times.

## **2.2. *Nikita* and *Sukkula* IRAP-PCR analyses**

IRAP analysis was performed according to Kalendar and Schulman [20]. PCR assays were performed in the T100 Thermal Cycler (BIO-RAD, USA). Amplification of the reactions was optimised in a final volume of 25  $\mu$ L containing 6.5  $\mu$ L ultrapure water, 12.5  $\mu$ L PCR master mix (DreamTaq Green PCR Master Mix, Thermo Scientific), 2  $\mu$ L of primer (0.8  $\mu$ M) and 4  $\mu$ L of template genomic DNA (3.2  $\mu$ M). Final concentrations were given in parenthesis. Primer sequences are 5'ACCCCTCTAGGCGACATCC3' for *Nikita* and 3'GGAACGTCGGCATCGGGCTG5' for *Sukkula* [21]. PCR conditions were as follows: one initial denaturation at 95°C for 3 min followed by 35 cycles of denaturation at 95°C for 30 sec, annealing at 54°C for 30 sec and extension at 72°C for 30 sec. The final elongation step was performed at 72°C for 5 min. The amplification products and molecular weight marker (100 bp DNA Ladder, Solis BioDyne) were resolved on 1% agarose gel in 1X Tris–Acetic Acid–EDTA at 90 V for 90 min and photographed on a Gel Documentation System (BIO-RAD). IRAP-PCR band profiles were evaluated visually and polymorphism ratios were calculated by Jaccard's coefficient in all samples [22].

### 2.3. Multiple alignment and phylogenetic tree analyses

Different transposons (*ALVE*, *CRI*, *ART-CH-O*, *ART-CH-L*, *ART-CH-D*, *ART-CH-B*, *ART-CH-R*, *ART-CH-H*, *ART*) belonging to the chicken genome as well as barley-specific *Nikita* and *Sukkula* were retrieved from NCBI (The National Center for Biotechnology Information – [www.ncbi.nlm.nih.gov](http://www.ncbi.nlm.nih.gov)) (Table 1). Evolutionary relationships among transposons were evaluated by multiple alignment analyses and phylogenetic tree construction. Clustal Omega (<http://www.ebi.ac.uk/Tools/msa/clustalo>) and MEGA X programs were used for this purpose.

**Table 1.** Transposons' accession numbers.

Multiple alignment results were used to construct a phylogenetic tree via MEGA X [23]. The program was adjusted as neighbour-joining (NJ) method [24] with p-distance model [25], and even bootstrap test (1000 replicates) for this analysis [26].

## 3. Results and discussion

### 3.1. Barley-specific *Nikita* and *Sukkula* retrotransposons identified in chicken genome

Due to the interesting nature and their abundance in the genome, retrotransposons are an excellent source for investigations of genome variations by using molecular markers. Therefore, different retrotransposon-based molecular markers such as REMAP (REtrotransposon-Microsatellite-Amplified Polymorphism), RBIP (Retrotransposon-Based Insertion Polymorphism), SSAP (Sequence-Specific Amplified Polymorphism) and iPBS (inter-Primer Binding Site) etc. have been developed not only genome variation but also explanation of disease and pathways related to biotic/abiotic stress in

different genomes [27,28]. In this study, *Nikita* and *Sukkula* retrotransposons were identified by using IRAP method relying on amplifying the DNA segments between two nearby retrotransposons using outward-facing primers [20]. Due to low cost, reproducibility, high polymorphism and informativity, this technique has been used for genetic assessment for several plant species because retrotransposons constitute up to 80% of plant genomes [29–31].

We studied 9 different chicken gDNAs to analyse the retrotransposons' movements using the IRAP technique. The experiment was replicated three times, and consistent results were obtained. Therefore, findings were given in one replicate. For the first time, these retrotransposons were identified in the chicken genome (Figure 1). Both monomorphic and polymorphic bands were observed in *Nikita* while polymorphism was not detected in *Sukkula*.

**Figure 1.** IRAP-PCR results of *Nikita* and *Sukkula*. M for markers and numbers from 1 to 9 indicated chicken samples

Therefore, *Nikita* IRAP-PCR analyses resulted in 45 scorable bands including 40 monomorphic and 5 polymorphic bands ranged from 100 to 800 bp. Monomorphic bands were indicated as positive (+) and polymorphic bands as negative (-) in Table 2.

**Table 2.** Band profiles of *Nikita* retrotransposon

According to these band profiles, we calculated polymorphism rates in all samples when compared to each other by using Jaccard's coefficient. Polymorphism ratios were between 0-60% in samples (Table 3).

**Table 3.** *Nikita* polymorphism percentages (%)

The transferability of TEs among species has provided valuable results to figure out evolutionary relationships. In our previous studies, different retrotransposons were identified in different plant genomes [17,18,32]. Many studies showed that the polymorphism rates of the same retrotransposons could be variable in different organs [33], among different individuals in the same species [31], and also different species [34] in normal and or stress conditions. Supporting these reports, we observed that there was no polymorphic band in *Sukkula* retrotransposon while *Nikita* polymorphism ratios were 0-60% among chicken samples. Similar to our study, Saraswathi et al. [35] also used *Nikita* and *Sukkula* retrotransposon in the germplasm collection of *Musa*. They reported that different primer pairs together with *Nikita* and *Sukkula* showed polymorphic band profiles among samples. This is an expected result because retrotransposon movements could be affected by many different situations including epigenetic and environmental conditions.

Although transposons' movements across families and even phyla within the same kingdoms have been identified in different studies, there is still little information about transposon transfer between kingdoms Animalia and Plantae [36]. The sequenced genomes have gained significant importance to identify the horizontal transfer of transposon (HTT) between animal and plant genomes. Gao et al. [37] performed one of



these investigations, reporting one of the first potential examples of HTT between animals and flowering plants by performing sequencing, phylogenetic and evolutionary analyses.

### **3.2. Sequence similarities were detected among transposons' sequences**

We also surveyed the evolutionary relationships among *Nikita*, *Sukkula* and chicken-specific retrotransposons using *in silico* screening. For this purpose, a total of 12 transposons consisting of chicken- and barley-specific retrotransposons were analysed to identify relationships among them. Figure 2 indicated part of the alignment analysis. We revealed that *Sukkula* and other sequences shared more similarity in clustal analysis.

**Figure 2.** Multiple sequence alignment of chicken- and barley-specific transposons. Every four nucleotides presented in different colours.

The phylogenetic tree was revealed by analysing 12 nucleotide sequences with a total of 502 positions (Figure 3). *Sukkula* and *CRI* sequences were found in the first clade. Moreover, two groups consisting of *ART* in one group and *Nikita*, *ALVE*, *ART-CH-R*, *ART-CH-H*, *ART-CH-B*, *ART-CH-O*, *ART-CH-D* and *ART-CH-L* in another group were observed in the second clade.

**Figure 3.** Phylogenetic analysis of chicken- and barley-specific transposons' sequences. Bootstrap percentages are indicated at the branch points.

Meyerowitz [38] concluded that approximately 1,600 Mya plants and animals diverged from a common ancestor. Sequencing results have provided valuable proof related to horizontal transfer of retrotransposons between two kingdoms [36,37]. Concordant with this, we observed that CR1 element resembles *Sukkula* retrotransposon whereas *Nikita* retrotransposon is similar to other chicken retrotransposons.

In conclusion, domestic animals have been commonly studied to identify roles in gene(s) especially related to phenotypic traits [39]. Transposons are one of the main evolution drivers affecting genes' expression. Therefore, increasing knowledge about genomes in these animals makes it possible to apply different methods to get desired traits. Here, we performed IRAP-PCR analyses to investigate the existence of barley-specific retrotransposons and *in silico* analyses to understand evolutionary relationships but no sequencing experiment was performed. Therefore, sequencing of native chickens' genomes will also give us detailed information between plants and chickens.

### **Role of funding source**

This study does not receive any grant.

### **Ethical Standards**

All procedures were approved and regularly controlled by the Animal Ethics Committee of Ondokuz Mayıs University (OMU HADYEK 06.11.2012/1).

### **Declaration of Competing Interest**

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

## **Acknowledgements**

The authors thank to the staff of Republic of Turkey Ministry of Agriculture and Forestry's Gerze Directorate for sampling.

## **Ethical clearance**

Ondokuz Mayıs University Animal Experiments Local Ethics Committee gave permission for this study with 06.11.2012/1 number.

## **References**

1. Mandal PK, Kazazian HH. SnapShot: vertebrate transposons. *Cell* 2008; 135 (1): 192-192. doi: 10.1016/j.cell.2008.09.028
2. Cordaux R, Batzer MA. The impact of retrotransposons on human genome evolution. *Nature Reviews Genetics* 2009; 10 (10): 691-703. doi: 10.1038/nrg2640
3. Zhang HH, Peccoud J, Zhang XG, Gilbert C. Horizontal transfer and evolution of transposable elements in vertebrates. *Nature Communications* 2020; 11 (1): 1-10. doi: 10.1038/s41467-020-15149-4
4. Ghonaim M, Kalendar R, Barakat H, Elsherif N, Ashry N et al. High-throughput retrotransposon-based genetic diversity of maize germplasm assessment and analysis. *Molecular Biology Reports* 2020; 47 (3): 1589-1603. doi: 10.1007/s11033-020-05246-4

5. Ramos KS, Moore S, Runge I, Tavera-Garcia MA, Cascone I et al. The nucleolin antagonist *N6L* inhibits *LINE1* retrotransposon activity in non-small cell lung carcinoma cells. *Journal of Cancer* 2020; 11 (3): 733-740. doi: 10.7150/jca.37776
6. Pettersson ME, Jern P. Whole-genome analysis of domestic chicken selection lines suggests segregating variation in *ERV* makeups. *Genes* 2019; 10 (2): 162. doi: 10.3390/genes10020162
7. Wick RR, Judd LM, Gorrie CL, Holt KE. Completing bacterial genome assemblies with multiplex MinION sequencing. *Microbial Genomics* 2017; 3 (10): e000132. doi: 10.1099/mgen.0.000132
8. Chen F, Dong W, Zhang J, Guo X, Chen J et al. The sequenced angiosperm genomes and genome databases. *Frontiers in Plant Science* 2018; 9: 418. doi: 10.3389/fpls.2018.00418
9. Arumugam R, Uli JE, Annavi G. A review of the application of next generation sequencing (NGS) in wild terrestrial vertebrate research. *Annual Research & Review in Biology* 2019; 31 (5): 1-9. doi: 10.9734/arrb/2019/v31i530061
10. International Chicken Genome Sequencing Consortium. Sequence and comparative analysis of the chicken genome provide unique perspectives on vertebrate evolution. *Nature* 2004; 432 (7018): 695-716. doi: 10.1038/nature03154
11. International Chicken Genome Sequencing Consortium. A genetic variation map for chicken with 2.8 million single nucleotide polymorphisms. *Nature* 2004; 432 (7018): 717-722. doi: 10.1038/nature03156
12. Hughes AL, Piontkivska H. DNA repeat arrays in chicken and human genomes and the adaptive evolution of avian genome size. *BMC Ecology and Evolution* 2005; 5 (1): 12. doi: 10.1186/1471-2148-5-12

13. Cogburn LA, Porter TE, Duclos M, Simon J, Burgess SC et al. Functional genomics of the chicken—a model organism. *Poultry Science* 2007; 86 (10): 2059-2094. doi: 10.1093/ps/86.10.2059
14. Lyimo CM, Weigend A, Msoffe PL, Eding H, Simianer H et al. Global diversity and genetic contributions of chicken populations from African, Asian and European regions. *Animal Genetics* 2014; 45: 836-848. doi: 10.1111/age.12230
15. Weigend S, Romanov M. Current strategies for the assessment and evaluation of genetic diversity in chicken resources. *Worlds Poultry Science Journal* 2001; 57: 282-828. doi: 10.1079/WPS20010020.
16. Sacco MA, Flannery DMJ, Howes K, Venugopal K. Avian endogenous retrovirus *EAV-HP* shares regions of identity with avian leukosis virus subgroup J and the avian retrotransposon *ART-CH*. *Journal of Virology* 2000; 74 (3): 1296-1306. doi: 10.1128/jvi.74.3.1296-1306.2000
17. Cakmak B, Marakli S, Gozukirmizi N. *SIRE1* retrotransposons in barley (*Hordeum vulgare* L.). *Russian Journal of Genetics* 2015; 51 (7): 661-672. doi: 10.1134/S1022795415070029
18. Marakli S, Calis A, Gozukirmizi N. Determination of barley-specific retrotransposons' movements in *Pinus nigra ssp. pallasiana* varieties: *pyramidata* and *Seneriana*. *Russian Journal of Genetics* 2019; 55 (1): 71-78. doi: 10.1134/S1022795419010101
19. Miller SA, Dykes DD, Polesky A. A simple salting out procedure for extracting DNA from human nucleated cells. *Nucleic Acids Research* 1988; 16 (3): 1215. doi: 10.1093/nar/16.3.1215

20. Kalendar R, Schulman AH. IRAP and REMAP for retrotransposon-based genotyping and fingerprinting. *Nature Protocols* 2006; 1 (5): 2478-2484. doi: 10.1038/nprot.2006.377
21. Leigh F, Kalendar R, Lea V, Lee D, Donini P et al. Comparison of the utility of barley retrotransposon families for genetic analysis by molecular marker techniques. *Molecular Genetics and Genomics* 2003; 269 (4): 464-474. doi: 10.1007/s00438-003-0850-2
22. Jaccard P. Nouvelles recherches sur la distribution florale. *Bulletin de la Société Vaudoise des Sciences Naturelles* 1908; 44: 223-270. doi: 10.5169/seals-268384
23. Kumar S, Stecher G, Li M, Knyaz C, Tamura K. MEGA X: Molecular evolutionary genetics analysis across computing platforms. *Molecular Biology and Evolution* 2018; 35 (6): 1547-1549. doi: 10.1093/molbev/msy096
24. Saitou N, Nei M. The neighbor-joining method: A new method for reconstructing phylogenetic trees. *Molecular Biology and Evolution* 1987; 4 (4): 406-425. doi: 10.1093/oxfordjournals.molbev.a040454
25. Nei M, Kumar S. *Molecular evolution and phylogenetics*. Oxford University Press. New York; 2000.
26. Felsenstein J. Confidence limits on phylogenies: An approach using the bootstrap. *Evolution* 1985; 39 (4): 783-791. doi: 10.1111/j.1558-5646.1985.tb00420.x
27. Gozukirmizi N, Yilmaz S, Marakli S, Temel A. Retrotransposon-based molecular markers; tools for variation analysis in plants. In: Taški-Ajduković, K. (Ed.), *Applications of Molecular Markers in Plant Genome Analysis and Breeding*. Research Signpost Kerala; 2015:19-45.

28. Thomas EO, Zuniga G, Sun W, Frost B. Awakening the dark side: retrotransposon activation in neurodegenerative disorders. *Current Opinion in Neurobiology* 2020; 61: 65-72. doi: 10.1016/j.conb.2020.01.012
29. Alzohairy A, Yousef M, Edris S, Kerti B, Gyulai G et al. Detection of LTR retrotransposons reactivation induced by *in vitro* environmental stresses in barley (*Hordeum vulgare*) via RT-qPCR. *Life Science Journal* 2012; 9 (4): 5019-5026.
30. Tufan AF, Ibisoglu MS, Yilmaz S, Gozukirmizi N. Investigation of rice (*Oryza sativa* L.) retrotransposons in different taxa. *Genetic Resources and Crop Evolution* 2020: 1-7. doi: 10.1007/s10722-020-00919-2
31. Yigider E, Taspinar MS, Aydin M, Agar G. Cobalt-induced retrotransposon polymorphism and humic acid protection on maize genome. *Biologia Futura* 2020: 1-8. doi: 10.1007/s42977-020-00001-z
32. Marakli S. Retrotransposon analyses in *Cucurbitaceae* family. *International Journal of Science Letters* 2019; 1 (1): 68-76. doi: 10.38058/ijsl.592537
33. Marakli S, Yilmaz S, Gozukirmizi N. *BARE1* and *BAGY2* retrotransposon movements and expression analyses in developing barley seedlings. *Biotechnology & Biotechnological Equipment* 2012; 26 (6): 3451-3456. doi: 10.5504/BBEQ.2012.0112
34. Cakmak B, Marakli S, Gozukirmizi N. *Sukkula* retrotransposon movements in the human genome. *Biotechnology & Biotechnological Equipment* 2017; 31 (4): 756-760. doi: 10.1080/13102818.2017.1316684
35. Saraswathi MS, Uma S, Ramaraj S, Durai P, Mustaffa MM et al. Inter retrotransposon based genetic diversity and phylogenetic analysis among the *Musa* germplasm accessions. *Journal of Plant Biochemistry and Biotechnology* 2020; 29 (1): 114-124. doi: 10.1007/s13562-019-00519-x

36. Lin X, Faridi N, Casola C. An ancient transkingdom horizontal transfer of *penelope-like* retroelements from arthropods to conifers. *Genome Biology and Evolution* 2016; 8 (4): 1252-1266. doi: 10.1093/gbe/evw076
37. Gao D, Chu Y, Xia H, Xu C, Heyduk K et al. Horizontal transfer of non-LTR retrotransposons from arthropods to flowering plants. *Molecular Biology and Evolution* 2018; 35 (2): 354-364. doi: 10.1093/molbev/msx275
38. Meyerowitz EM. Plants compared to animals: the broadest comparative study of development. *Science*. 2002; 295 (5559): 1482-1485. doi: 10.1126/science.1066609
39. Rubin CJ, Zody MC, Eriksson J, Meadows JRS, Sherwood E et al. Whole-genome resequencing reveals loci under selection during chicken domestication. *Nature* 2010; 464 (7288): 587-591. doi: 10.1038/nature08832



**Table 1.** Transposons' accession numbers.


<b>Name</b>	<b>Accession number</b>	<b>Name</b>	<b>Accession number</b>
<i>Nikita</i>	AF474072.1:c22229-21152	<i>ART-CH-L</i>	DQ500090.1
<i>Sukkula</i>	AF453665.1	<i>ART-CH-H</i>	DQ500091.1
<i>Sukkula</i>	AH014393.2:c86082-81112	<i>ART-CH-D</i>	DQ500089.1
<i>ALVE</i>	KJ908690.1	<i>ART-CH-B</i>	DQ500088.1
<i>CRI</i>	U88211.1	<i>ART-CH-R</i>	DQ500092.1
<i>ART-CH-O</i>	DQ500087.1	<i>ART</i>	L25262.1

**Table 2.** Band profiles of *Nikita* retrotransposon

	<b>1</b>	<b>2</b>	<b>3</b>	<b>4</b>	<b>5</b>	<b>6</b>	<b>7</b>	<b>8</b>	<b>9</b>
+	4	4	5	3	5	5	4	5	5
-	1	1	0	2	0	0	1	0	0

**Table 3.** *Nikita* polymorphism percentages (%)

<b>%</b>	<b>1</b>	<b>2</b>	<b>3</b>	<b>4</b>	<b>5</b>	<b>6</b>	<b>7</b>	<b>8</b>	<b>9</b>
<b>1</b>	-								
<b>2</b>	40	-							
<b>3</b>	20	20	-						
<b>4</b>	<b>60</b>	25	40	-					
<b>5</b>	20	20	0	40	-				
<b>6</b>	20	20	0	40	0	-			
<b>7</b>	40	0	20	25	20	20	-		
<b>8</b>	20	20	0	40	0	0	20	-	
<b>9</b>	20	20	0	40	0	0	20	0	-



```


Nikita
ALVE 1 -----
Sukkula 99 TTTGATTGTTGTAATTGTCCTGC---AGTGACTTGTTTGCAATTCAGCCGTGCCATGCAT
Sukkula 100 TTTGATTGTTGTTGATTGCCATGC---CGTGCAT-----TAGACCGTTCATGTAT
CR1 121 GAAAGTGGGCAGACGAGATAGGA-----AACCAGCATGGCTCGGGCAAGGACCTGCTG
ART-CH-O 61 --CAGTATGGCCATGTTAGAGCACAGTGCCCCGTGGGGGAGGCCAGTCAGGGCCCTCCTG
ART-CH-L 61 --CAGTATGGCCATGTTAGAGCACAGTGCCCCGTGGGGGAGGCCAGTCAGGGCCCTCCTG
ART-CH-D 61 --CAGTATGGCCATGTTAGAGCACAGTGCCCCGTGGGGGAGGCCAGTCAGGGCCCTCCTG
ART-CH-B 61 --CAGTATGGCCATGTTAGAGCACAGTGCCCCGTGGGGGAGGCCAGTCAGGGCCCTCCTG
ART-CH-R 61 --CAGTATGGCCATGTTAGAGCACAGTGCCCCGTGGGGGAGGCCAGTCAGGGCCCTCCTG
ART-CH-H 61 --CAGTATGGCCATGTTAGAGCACAGTGCCCCGTGGGGGAGGCCAGTCAGGGCCCTCCTG
ART 118 TACGCCAGTATGCATGTTAGAGCACAAATGCCCTGTGGGGGAGGCCAGTCAGGGCCCTCCTG

Nikita
ALVE 1 -----
Sukkula 155 CATATATGTTGTGCTTTGTGTTGGTGAATTCGTGTGTTGATTTGTGTTCCGGTTTTCTT
Sukkula 146 CATGTGTGTTGTCATCGTGTGGTGTATATCGTGTGTTGATGCTTGTTCCGGTTTCTCTC
CR1 173 GGCACACTGAGGGTGAAGAAAGGTGCGTACAAGCTC-----TGGAAACAAGGGCGTGT
ART-CH-O 119 ATCACAGGAAGGGCTTGC TAAAGGGTATCCGTGCTC-----GGGTATGCG-----
ART-CH-L 119 ATCACAGGAAGGGCTTGC TAAAGGGTATCCGTGCTC-----GGGTATGCG-----
ART-CH-D 119 ATCACAGGAAGGGCTTGC TAAAGGGTATCCGTGCTC-----GGGTATGCG-----
ART-CH-B 119 ATCACAGGAAGGGCTTGC TAAAGGGTATCCGTGCTC-----GGGTATGCG-----
ART-CH-R 119 ATCACAGGAAGGGCTTGC TAAAGGGTATCCGTGCTC-----GGGTATGCG-----
ART-CH-H 119 ATCACAGGAAGGGCTTGC TAAAGGGTATCCGTGCTC-----GGGTATGCG-----
ART 178 ATCACAGGAAGGGCTTGC TCAAGGGTATCCGTGCTC-----GGGTATGCA-----

Nikita
ALVE 1 -----
Sukkula 215 CGTCTCGTTAGAGTTCCG---CAGCGTGCCGGATTGTGAGGACCCGTTGACTACGTCCGG
Sukkula 206 CGTCTCGATAGAGTTCCGC--AAGCGTGCCGGATTGTGAGGACTCGTTCGACTACGTCCGG
CR1 226 CACC TGGGAAGAGTACAGGGATGCTGTCCGGACTTGACAGCCTAGGATCAGGAAAGCCA-
ART-CH-O 164 -GCAGT-----TAAACATCATGAATCTGGGAAATTCGTGGGTACCATCGCGAAAAGTA-
ART-CH-L 164 -GCAGT-----TAAACATCATGAATCTGGGAAATTCGTGGGTACCATCGCGAAAAGTA-
ART-CH-D 164 -GCAGT-----TAAACATCATGAATCTGGGAAATTCGTGGGTACCATCGCGAAAAGTA-
ART-CH-B 164 -GCAGT-----TAAACATCATGAATCTGGGAAATTCGTGGGTACCATCGCGAAAAGTA-
ART-CH-R 164 -GCAGT-----TAAACATCATGAATCTGGGAAATTCGTGGGTACCATCGCGAAAAGTA-
ART-CH-H 164 -GCAGT-----TAAACATCATGAATCTGGGAAATTCGTGGGTACCATCGCGAAAAGTA-
ART 223 -GCAGT-----TAAATATCATGAATCTGGGAAATTCGTGGGTACCATCACGAAAAGTA-

Nikita
ALVE 1 -----
Sukkula 451 TTCAACCTAG-----CAAACCCTGATTGGTTA---TGTTACTGCTTGCTTAAACCC
Sukkula 441 TTCAACCTAG-----CAAACCCTGATTGGCTA---TGTTACTGCTTGCTTAAACCC
CR1 446 AATGAGTTCT-----TTGCCTCAGTCTTCACTGGAGGCCAGGATTCAG
ART-CH-O 336 AAGAACCCTCATCAAGAGGAGCAC TACACCTGCAACGCTGACACTTCCGATGTTGCTGA
ART-CH-L 336 AAGAACCCTCATCAAGAGGAGCAC TACACCTGCAACGCTGACACTTCCGATGTTGCTGA
ART-CH-D 336 AAGAACCCTCATCAAGAGGAGCAC TACACCTGCAACGCTGACACTTCCGATGTTGCTGA
ART-CH-B 336 AAGAACCCTCATCAAGAGGAGCAC TACACCTGCAACGCTGACACTTCCGATGTTGCTGA
ART-CH-R 336 AAGAACCCTCATCAAGAGGAGCAC TACACCTGCAACGCTGACACTTCCGATGTTGCTGA
ART-CH-H 336 AAGAACCCTCATCAAGAGGAGCAC TACACCTGCAACGCTGACACTTCCGATGTTGCTGA
ART 395 AAGAACCCTCGTCAAGAGGGCACT-ACACCTGCAACGCTGATTCTGTTGC-----TA

```



## Figure Captions

**Figure 1.** IRAP-PCR results of *Nikita* and *Sukkula*. M, marker; 1-9, different chicken samples.

**Figure 2.** Multiple sequence alignment of chicken- and barley-specific transposons. Every four nucleotides presented in different colours.

**Figure 3.** Phylogenetic analysis of chicken- and barley-specific transposons' sequences. Bootstrap percentages are indicated at the branch points.