

Comparison of volatile chemical components of *Cystoseira crinita* Duby by hydro-distillation (HD) and solid-phase microextraction (SPME) methods and antimicrobial and scolicidal activities of essential oil and extracts

Tayyibe Beyza YÜCEL*

Giresun University, Health Services Vocational School, Giresun, Turkey

Received: 12.04.2021 • Accepted/Published Online: 25.11.2021 • Final Version: 27.04.2022

Abstract: This study aims to derive an essential oil from *Cystoseira crinita* Duby, analyze the chemical composition of the essential oil, discover the antimicrobial activities of the oil and the extracts, and investigate the scolicidal activities of the extracts. The volatile organic compounds of *Cystoseira crinita* Duby were determined by GC/MS-FID using both hydro-distillation (HD) and solid-phase microextraction (SPME) methods. As a result of the essential oil analysis, 97.14% of 17 compounds and 93.13% of 19 compounds were elucidated. The main compounds identified were hexanal (31.802%), n-hexadecanoic acid (12.654%), trans- β -ionone (9.118%), 2E-hexenal (15.955%), heptadecane (15.729%) and tetradecane (13.458%). In addition, hexane, dichloromethane, chloroform, and methanol extracts of the algae sample were prepared. The antimicrobial activities of the essential oil and extracts on gram-positive bacteria, gram-negative bacteria, and fungi microorganisms were studied. The results revealed activity within the zone diameter range of 8-16 mm. It was observed that all extracts and the essential oil itself showed high activity against the *Pseudomonas aeruginosa* ATCC 27853 microorganism. The chloroform and dichloromethane extracts were also found to demonstrate a high level of efficacy against *Bacillus cereus* ATCC 10876. Furthermore, viability detection was performed and the scolicidal effects of the extracts on protozoa were assessed. All extracts showed a strong scolicidal effect at the dose of 15000 $\mu\text{g/mL}$. For each solution, the difference between hours at each dose and the difference between doses at every hour were compared by One-Way ANOVA. The values of lethal concentration doses (LD_{50} and LD_{90}) were calculated using Probit Analysis. This study provides information about the effects of *Cystoseira crinita* Duby algae extracts and suggests that the experimental studies needed for their use in live cells should be performed.

Key words: *Cystoseira crinita* Duby, volatile compound, antimicrobial effect, scolicidal effect, lethal dose, probit analysis

1. Introduction

Microalgae and macroalgae are vegetative organisms. Macroalgae are aquatic organisms and remarkably dominant in marine ecosystems. They are also widely consumed as food in coastal regions, especially in the Far Eastern countries such as China, Korea, and Japan. Microalgae are classified as brown (Phaeophyta), red (Rhodophyta), and green (Chlorophyta) depending on their pigmentation and if they bear a flagellum [1,2]. Approximately 9000 species have been recorded worldwide, while there are approximately 5000 species recorded and 600 bibliography and distribution records in Turkey [3,4].

Since algae are photosynthetic organisms, they can provide seawater with oxygen (O_2) while also constituting a reliable food source for marine organisms such as shrimp and jellyfish with the acids, alkaloids, amines, cellulose, enzymes, glycosides, trace elements, inorganic minerals, lipids, sterols, ceroids, vitamins, amino acids, proteins, pepsins, and volatile compounds [2,5,6]. Algae's bioactive components (secondary metabolites, diterpenes, brominated sesquiterpenes, sesquiterpenes, polyphenols, flavonoids, minerals, polysaccharides, polyunsaturated fatty acids, vitamins, etc.) means they are often used for their sedative, muscle relaxant, and edema relieving properties in both medicine and pharmacology. They are also used as an agricultural fertilizer and in the production of polysaccharides, such as agar and alginate, in the food industry. Additionally, algae can be used to produce biofuels such as ethanol, butanol, and biogas [7-9]. Many studies show that algae form resistance to bacteria, fungi, and viral pathogens [10,11]. In addition, red and green algae appear to have a preventative effect on the skin, breast, and intestinal cancer [2].

There are more than 7000 species of brown algae, yet less than one percent are found in freshwater environments, while the remaining part can be found in the seas. Some species are microscopic, and some can reach up to 30 meters

* Correspondence: beyza.yucel@giresun.edu.tr

[12]. Brown algae are also rich in polyphenols, which have antioxidant properties. They are used in the rubber industry, dyes, ice cream, and plastic freezers due to their alginate content [13,14]. *Fucus vesiculosus* (Bladderwrack), also known as kelp, is part of the brown algae class, and it contains a high amount of iodine. It is known to have a high antioxidant and antiinflammatory effect and can be used to treat hormonal diseases.

Essential oils derived from plants are aromatic mixtures obtained from root, stem, leaf, fruit, shell, and flower. They are liquid at room temperature and can crystallize with ease [15]. The following is a list of academic studies regarding the essential oil analyses of algae and their anticancer [16], antileukemic [17], antioxidant [18,19], antimicrobial [20], antifeedant [21], antifouling [22], antimalarial [23], and cytotoxic [24,25] properties.

Due to the algae's popularity in far eastern countries as a food group for its cosmetic and medicinal uses and its wide range of biological activity, there are many studies worldwide detailing its benefits. But the number of such studies is quite limited in Turkey, and what studies, primarily using algae collected from the Aegean Sea, were conducted. Therefore, this study was carried out in the Black Sea due to the small number of such studies in this region.

There are a few reports on the antibacterial, antiprotozoal, antimycobacterial, and cytotoxic activity of methanol extract of *Cystoseira crinita* Duby grown in different parts of the world including Turkey. In these studies, biological activities were found quite high. Since this species has been found to have wide biological activity in the limited number of studies in the literature, it is aimed to determine the chemical composition and different biological activity studies of this species.

This study aims to detect the volatile components of *Cystoseira crinita* Duby, which is in the brown algae class, situated around the Sinop province, using both hydro-distillation (HD) and solid-phase microextraction (SPME) methods, and to extract Soxhlet using chloroform (Ch), dichloromethane (DCM), hexane (H) and methanol (MeOH), to determine the antimicrobial activities of both essential oil (HD) and extracts and the scolicidal effects of extracts. There are already studies that determine the antimicrobial and antioxidant properties of various extracts of *C. crinita* collected from different seas worldwide, including Turkey [26,27]. However, no studies have been found regarding the antimicrobial analysis of essential oil and solvent extracts except for methanol and chloroform of *C. crinita* Duby. Also, in previous studies, there is no scolicidal activity of solvent extracts. Although there are antimicrobial activity studies, this study differs in terms of the essential oil composition and test-microorganisms used for antimicrobial activity.

2. Materials and Methods

2.1. Algae material

C. crinita was collected in November 2017 from Sinop, Aklıman and Merkez Tersane provinces. The algae were stored in polyethylene bottles and transported to the laboratory using a cold chain. The sample was first rinsed with seawater to remove sand particles, then cleansed of any remaining sea organisms. It was then left to dry in the open air at room temperature under shade. The dried samples were then cut into smaller sections using a mixer in the laboratory and prepared for testing.

2.2. Isolation of essential oil

The essential oil was extracted using a Clevenger apparatus. To achieve this, the dried and crumbled algae sample *C. crinita* (65 g) was weighed and placed into a two-liter flask and then mixed with distilled water. The essential oils were extracted using a Clevenger apparatus connected to the cooling bath (-12 °C) over four hours. The derived essential oils were extracted from the Clevenger apparatus using HPLC purity *n*-hexane (0.5 mL) and transferred into brown vials. The amount of oil obtained as a result of hydro-distillation with a Clevenger apparatus was weighed and found to be 15.5 mg, respectively. It was then weighed and stored at +4 °C while awaiting the antimicrobial activity tests.

2.3. SPME analysis

Solid phase micro-distillation was used for this process. One gram of air-dried algae material was grounded and placed into a vial for SPME analysis. Vials were 10 mL in volume and sealed with a silicone-rubber septum cap. A polydimethylsiloxane/divinylbenzene fiber was used to extract the volatile components. Extractions were achieved with magnetic stirring.

The fiber coating fibers were conditioned for five minutes at 250 °C in GC injector before being placed into the headspace. Temperature, incubation, and extraction times were set according to the experiment. SPME was processed at 50 °C with an incubation time of 5 min and an extraction time of 10 min. Each sample was then analyzed and reported [28].

2.4. Extraction using the Soxhlet apparatus

Thirty grams of dried *C. crinita* portions were weighed out, put into Soxhlet cartridges, and placed in the reflux. The soxhlet extraction took place over 24 h by adding 250 mL of chloroform (Ch), dichloromethane (DCM), hexane (H), and methanol (MeOH) into each flask. The end solution was then passed through the evaporator and dried. The remaining

residue was weighed (3.2%, 2.5%, 2.5%, and 3.8%, respectively) and stored at +4°C awaiting biological activity tests. All solvents used were 95%–98% pure and Sigma–Aldrich branded.

2.5. Gas chromatography-mass spectrometry (GC-MS/FID)

The gas chromatography chromatography-flame ionization detector (GC-FID) analysis was carried out on a Shimadzu QP2010 plus gas chromatography coupled to a Shimadzu QP2010 ultra mass selective detector. The fiber-containing, extracted aroma compounds were injected into the GC-MS injector (split mode). Separation took place with a Restek Rxi-5MS capillary column of 30 m length, 0.25 mm i.d., and 0.25 µm phase thickness. The oven program began at an initial temperature of 60 °C for 2 min, which was then increased to 240 °C for three minutes, and then a final temperature of 250 °C for four minutes. The injector temperature was 280 °C and the split ratio was 1:20. The carrier gas was helium (99.999%), with a constant flow rate of 1 mL/min; sample size was 1 µL. Detection was processed in electronic impact mode (EI); ionization voltage was fixed to 70 eV, and scan mode (40–450 m/z) was used for mass acquisition.

2.6. Analysis of components

The retention indices (RI) of the components were determined with n-alkanes (C₆-C₃₀) using the Koyats method. Comparison retention indices of volatile components (relative to C₆-C₃₀ alkane standards) were identified by comparing the mass spectra of the two libraries (FFNSC1.2 and W9N11).

2.7. Antimicrobial Analysis

Antimicrobial activity was performed using the disc diffusion method as described by Demirel et al. [29]. For antimicrobial activity, a total of 19 bacterial species were used. The gram-positive bacteria were *Bacillus (B.) subtilis* ATCC 6633, *Bacillus cereus* ATCC 10876, *Bacillus megaterium* DSM32, *Staphylococcus (S.) aureus* ATCC 29213, *Staphylococcus aureus* ATCC 25923, *Enterococcus faecalis (E.)* ATCC 29212, Metisillin Resistant *Staphylococcus aureus* (MRSA) ATCC 67101, *Staphylococcus epidermidis* ATCC 12228; and the gram-negative bacteria were *Cedecea (C.) neteri* ATCC 33855, *Escherichia (E.) coli* ATCC 25922, *Escherichia coli* ATCC 36218, *Pseudomonas (P.) aeruginosa* ATCC 27853, *Pseudomonas aeruginosa* ATCC 9027, *Klebsiella (K.) pneumoniae* ATCC 13883, *Acinetobacter (A.) baumannii* ATCC BAA-747, *Enterobacter (En.) aerogenes* ATCC 13048, *Citrobacter (Cb.) freundii* ATCC 43864, *Salmonella (Sa.) typhimurium* ATCC 14028, *Proteus (Pr.) mirabilis* ATCC 43071. The *Candida (Ca.) albicans* ATCC 10231 strain was used for antifungal activity.

To determine the antimicrobial activity of *C. crinata* essential oil, the essential oil was dissolved in DMSO at a concentration of 8.61 mg/mL. Similarly, in antimicrobial studies of Ch, MeOH, H, and DCM extracts of the algae sample, the extracts were dissolved in DMSO with a concentration of 15 mg/mL. Six mm sterile blank discs were soaked in 2 different amounts of extracts each prepared using different solvents, A:15 µL and B:30 µL. To measure the essential oil and extracts' antimicrobial effect, the microorganisms tested were inoculated in a Müller Hinton Broth and the yeasts in a Sabouraud Dextrose Broth, which were incubated for 18 h at 37 °C. Turbidity measurements were then adjusted with a densitometer to McFarland No: 0.5. Each broth was adjusted to McFarland No: 0.5 under sterile conditions. From those, 100 µL was put into all Mueller Hinton Agar Broths and equally distributed to Petri dishes with a sterile swab. The prepared discs were placed into Petri dishes, and the microorganisms were inoculated. After 24 h of incubating at 37°C, the discs' zone diameters and measured and the values were noted in millimeters. Tobramycin (Bioanalyse, 10 µg/disc), Nystatin (Bioanalyse, 30 µg/disc), and Vancomycin (Bioanalyse, 30 µg/disc) standard antibiotic discs were used as controls.

2.8. Scolicidal analysis

The viability of the protoscolices obtained from a cow liver cyst hydatid was determined using a 0.1% eosin solution. Among cyst hydatid treatment methods, percutaneous interventions can be applied according to its localization. Scolicidal agents can be used in these interventions [30]. Scolicidal agents include 20% or 30% hypertonic saline, 95% ethanol, 0.5% cetrimide. Researchers have stated that percutaneous interventions are safe and effective [31,32]. The solution was dissolved in sterile saline. Its final concentration was adjusted to 15000 µg/mL and was homogenized and sterilized by filtering the membrane.

Sterile saline of 100 µL was added to the wells of the sterile microplate. A total of 100 µL of *C. crinita* extracts were then added to the first wells and diluted to a 1/2 ratio. Once diluted, a 3500 piece/mL concentration of 100 µL protoscolices was added to the wells. As a control, only 100 µL protoscolices with a certain dilution were added to wells 9 and 10. Counting was used to determine the viability rates of protoscolices at certain times. Each counting was performed twice, and the results were averaged. A frequently used in the study routine NaCl solution (30%) was included as a scolicidal agent.

2.9. Statistical analysis

The descriptive statistics of the data set are expressed as a mean ± standard deviation (SD). The differences between means were compared by the one-way ANOVA followed by Tukey's post-hoc test. Lethal dose values (LD₅₀ and LD₉₀) were determined using probit analysis at certain times. A p-value was considered statistically significant at ≤0.05. All statistical

analyses were performed using the SPSS v26 (IBM Inc., Chicago, IL, USA) and the Minitab 19 (Minitab Inc., State College, PA, USA) statistical software.

2.10. Findings

The amount of essential oil extracted from the steam distillation of *C. crinita* weighed to be 15.5 mg (0.085% w/w). A total of 17 and 13 compounds, comprising 97.14% and 93.13% of the HD and SPME, respectively, were identified (**Table 1**). Weighted components in the essential oil derived by hydro-distillation were 38.88% aldehyde, 18.68% terpenoid class compounds, while the SPME main component class was determined to be 53.89% aldehyde and 36.73% hydrocarbons compounds (**Table 1**). Compounds identical to the SPME and HD analyses were found to be hexenal (31.80% and 12.19%), (2*E*,4*E*)-heptadienal (6.50% and 5.95%), pentadecane (1.55%-1.35%), and hexadecane (3.02%-4.73%). In this context, a difference in content can be observed. Main components were hexenal (31.80%), *n*-hexadecanoic acid (12.65%), trans- β -ionone (9.11%) in HD analysis and 2*E*-hexenal (15.95%), heptadecane (15.72%) and tetradecane (13.45%) in SPME analysis.

Table 1. Identified components in the essential oils of *C. crinita*.

| | Retention Time | Compounds | HD % Area ^a | SPME %Area ^a | Exp. RI ^b | Lit. RI |
|----|----------------|---------------------------------------|------------------------|-------------------------|----------------------|---------|
| 1 | 8.593 | Hexenal | 31.802 | 12.194 | 801 | 803 |
| 2 | 10.022 | (<i>E</i>)-2-Hexenal | - | 15.955 | 852 | 851 |
| 3 | 10.162 | (<i>Z</i>)-3-Hexenol | 1.558 | - | 858 | 858 |
| 4 | 11.710 | Heptanal | - | 12.481 | - | 903 |
| 5 | 14.724 | 1-octene-3-ol | 0.679 | | 977 | 978 |
| 6 | 15.305 | Furan-2-pentyl | 2.072 | | 991 | 993 |
| 7 | 15.621 | (2 <i>E</i> ,4 <i>E</i>)-Heptadienal | 6.507 | 5.957 | 1009 | 1012 |
| 8 | 17.022 | Limonene | - | 2.518 | 1031 | 1030 |
| 9 | 17.646 | (<i>E</i>)-2-Octenal | | 1.087 | 1047 | 1049 |
| 10 | 18.135 | Benzene acetaldehyde | 0.579 | | 1056 | 1052 |
| 11 | 20.055 | Nonanal | - | 3.880 | 1102 | 1101 |
| 12 | 24.406 | Decanal | | 2.337 | 1202 | 1201 |
| 13 | 28.220 | Tridecane ^c | | 1.459 | 1303 | 1300 |
| 14 | 32.203 | Tetradecane ^c | | 13.458 | 1403 | 1400 |
| 15 | 34.408 | Geranyl acetone | 2.411 | | 1450 | 1453 |
| 16 | 35.766 | 1-Pentadecene | 3.980 | | 1485 | 1489 |
| 17 | 35.898 | Trans- β -ionone | 9.118 | | 1489 | 1487 |
| 18 | 36.040 | Pentadecane | 1.551 | 1.353 | 1502 | 1500 |
| 19 | 40.008 | Hexadecane | 3.024 | 4.731 | 1601 | 1600 |
| 20 | 41.215 | Benzophenone | 7.950 | | 1625 | 1626 |
| 21 | 43.177 | Heptadecane ^c | - | 15.729 | 1701 | 1700 |
| 22 | 43.963 | Pentadecanone | 3.417 | | 1715 | 1710 |
| 23 | 45.383 | Tetradecanoic acid | 2.694 | | 1758 | 1763 |
| 24 | 48.017 | Hexahydrofarnesyl acetone | 4.579 | | 1849 | 1848 |
| 25 | 51.778 | <i>n</i> -Hexadecanoic acid | 12.654 | | 1960 | 1966 |
| 26 | 56.296 | Phytol | 2.572 | | 2114 | 2110 |
| | | Total | %97.14 | %93.13 | | |

^a RI calculated from retention times relative to that of *n*-alkanes (C₆-C₃₀) on the non-polar HP-5 column.

^b % Area obtained by FID peak-area normalisation.

^c Identified by authentic samples.

The antimicrobial activity results of the essential oil obtained by HD were measured on five microorganisms, including *S. aureus* 25923, MRSA, *B. cereus* ATCC 10876, *P. aeruginosa* 27853, *E. coli* 25922, *Ca. albicans* ATCC 10231, diameter values were determined medium 14–13 mm zone on *B. cereus* and *P. aeruginosa* according to Tobramycin standard agent. But no antimicrobial activity was observed against the fungus *Ca. albicans* according to standart agent as Nystatin. The results are shown in **Table 2**. The Petri view of the essential oil zone on *B. cereus* can be seen in **Figure 1**.

The antimicrobial effect of extract' *C. crinita* according to the disc diffusion method is displayed in **Table 3**. The antimicrobial activities of both Ch and DCM extracts of *C. crinita* at 30 μ L concentration weren't observed to have an antimicrobial effect against *Pr. mirabilis*; Ch, H, and MeOH extract against *E. coli* ATCC 25922 and the H and Ch extract against *Sa. typhimurium* ATCC 14028. In the 8-16 mm zone range at both 15 μ L and 30 μ L concentrations, the antimicrobial effect was observed only against *B. cereus* ATCC 10876 and *P. aeruginosa* ATCC 27853. The highest zone values of all extracts were found against *P. aeruginosa* ATCC 27853 according to Tobramycin as standard agents. All extract concentrations except for 15 μ L dose of M extract were determined antifungal activity against *Ca. albicans* ATCC 10231.

The scolicial effect of Ch, DCM, H and MeOH extracts of *C. crinita* prepared in different concentrations on protoscolices at different hours are shown in **Figures 2–6**.

When examining the graphs, the extracts of *C. crinita* prepared in different concentrations were seen to decrease with time except for the control group. When the applied dose effect was examined, the decrease in the viability rate per time was observed to be faster as the dose amount increased.

The scolicial effect of the *C. crinita* extracts in organic solvents at different concentrations on protoscolices, according to the dose applied at different times, is shown in Table 4 to Table 8.

It was observed that the DCM extract prepared at different concentrations of *C. crinita* had a very strong scolicial effect at 15000 μ g/mL. It has been determined that it affects the parasites in a very short time compared to other concentrations, and no live parasites have been encountered, especially at the 9th h (**Table 4**).

When **Table 5** was examined, it was found that the hexane extract of *C. crinita* had a very strong scolicial effect at the concentration prepared at 15000 μ g/mL. No live parasites were found at the 11th ..

As other extracts DCM and Hexane, it has been determined that Chloroform extract has a very strong scolicial effect at a concentration of 15000 μ g/mL. No live parasites were found at the 11th . (**Table 6**).

Table 2. Antimicrobial activity of the essential oil of *C. crinita*.

| Microorganisms | Inhibition Zone(mm) | | Standards | |
|--------------------------------|---------------------|----|-----------|-----|
| | A | B | Tob | Nys |
| <i>S. aureus</i> 25923 | - | - | 22 | - |
| MRSA | - | - | 20 | - |
| <i>B. cereus</i> ATCC 10876 | - | 14 | 18 | - |
| <i>P. aeruginosa</i> 27853 | - | 13 | 20 | - |
| <i>E. coli</i> 25922 | - | - | 22 | - |
| <i>Ca. albicans</i> ATCC 10231 | - | - | - | 24 |



Figure 1. Antimicrobial test view of essential oil on *B. cereus* A:15 μ L, B:30 μ L, Tob: Tobramycin, Nys: Nystatin.

Table 3. Antimicrobial effect of extracts of *C. crinita* on microorganisms.

| Microorganisms | Inhibition Zone (mm) | | | | | | | | | | |
|---------------------------------|----------------------|-----|------------------|-----|----------------|-----|-------------------|-----|-----------------------|-----|----|
| | Ch ^a | | DCM ^a | | H ^a | | MeOH ^a | | Standard ^a | | |
| | A | B | A | B | A | B | A | B | Tob | Nys | Va |
| <i>S. aureus</i> ATCC 25923 | 6,5 | 9 | 6,5 | 9 | - | 6,5 | 7 | 8 | 20 | - | - |
| <i>S. aureus</i> ATCC 29213 | - | 9 | - | 9 | - | 8 | - | 9 | 20 | - | - |
| <i>S.epidermidis</i> ATCC 12228 | - | 7 | - | 7 | - | 10 | - | 8 | - | - | - |
| MRSA ATCC 67101 | 6,5 | 8 | - | 6,5 | 6,5 | 7 | 7 | 9 | - | - | 17 |
| <i>B. cereus</i> ATCC 10876 | 8 | 12 | 8 | 12 | 6,5 | 7 | 6,5 | 7 | 22 | - | - |
| <i>B. megaterium</i> DSM32 | - | 10 | - | 10 | - | 6,5 | - | 8 | 18 | - | - |
| <i>B. subtilis</i> ATCC 6633 | - | 9 | - | 8 | - | - | - | 10 | 18 | - | - |
| <i>E. faecalis</i> ATCC 29212 | - | 7 | - | 8 | - | 9 | - | 10 | 16 | - | - |
| <i>Paeruginosa</i> ATCC 27853 | 10 | 10 | 7 | 9 | 12 | 16 | 12 | 14 | 18 | - | - |
| <i>Paeruginosa</i> ATCC 9027 | - | 10 | - | 10 | - | 7 | - | 7 | 20 | - | - |
| <i>E. coli</i> ATCC 25922 | - | - | - | 6,5 | - | - | - | - | 20 | - | - |
| <i>E. coli</i> ATCC 36218 | - | 9 | - | 8 | - | 7 | - | 9 | 20 | - | - |
| <i>K.pneumoniae</i> ATCC 3883 | - | 8 | - | 8 | - | 9 | - | 9 | 19 | - | - |
| <i>A.baumannii</i> ATCCBAA-47 | - | 6,5 | - | 6,5 | - | 6,5 | - | 6,5 | 22 | - | - |
| <i>En.aerogenes</i> ATCC 13048 | - | 8 | - | 7 | - | 8 | - | 9 | 17 | - | - |
| <i>Cb. freundii</i> ATCC 43864 | - | 7 | - | 7 | 6,5 | 7 | 6,5 | 7 | 20 | - | - |
| <i>C. neteri</i> ATCC 33855 | - | 9 | - | 9 | - | 8 | - | 8 | 18 | - | - |
| <i>Sa.typhimurium</i> ATCC14028 | - | 7 | - | 7 | - | - | - | - | - | - | - |
| <i>Pr. mirabilis</i> ATCC 43071 | - | - | - | - | - | 8 | - | 7 | 19 | - | - |
| <i>Ca. albicans</i> ATCC 10231 | 7 | 8 | 7 | 9 | 6,5 | 7 | - | 6,5 | - | 24 | - |

Ch: Chloroform, DCM: Dichloromethane, H: Hexane, MeOH: Methanol, A: 15 µL, B: 30 µL
 Tob: Tobramycin, Nys: Nystatin, Va: Vancomycin
 a: mg/disk




Figure 2. Ch extract (%) viability rate.




Figure 3. DCM extract (%) viability rate.




Figure 4. H extract (%) viability rate.




Figure 5. MeOH extract (%) viability rate.




Figure 6. NaCl extract (%) viability rates.

Table 4. Dichloromethane extract of *C. crinita* viability rate comparison at different hours.

| Doses | 3h | 4h | 7h | 9h | 11h | 17h | 20h | 24h | 30h | 32h |
|---------|-----------|------------|----------|----------|----------|-----------|-----------|-----------|--------|--------|
| Control | 100.00 a | 100.00 a | 100.0 a | 100.0 a | 100.0 a | 100.00 a | 100.00 a | 100.00 a | 100.0a | 100.00 |
| 117 | 90.89 ab | 90.00 ab | 78.20 ab | 63.79 b | 56.06 ab | 49.00 b | 31.48 b | 19.23 b | 8.17 b | 0.00 |
| 234 | 86.34 abc | 78.3 abc | 63.00 ab | 53.45 bc | 49.30 bc | 41.30 bc | 27.78 bc | 13.46 bc | 0.00 c | 0.00 |
| 468 | 77.50 bcd | 66.67 abcd | 57.90 ab | 50.00 bc | 45.70 bc | 37.50 bc | 18.52 bcd | 11.54 bcd | 0.00 c | 0.00 |
| 937 | 65.54 cde | 58.33 bcd | 57.90 ab | 48.28 bc | 43.80 bc | 33.80 bcd | 14.81 cde | 9.62 bcd | 0.00 c | 0.00 |
| 1875 | 61.12 de | 55.00 cd | 46.10 ab | 36.21 bc | 32.80 bc | 20.73 bcd | 11.11 de | 7.69 bcd | 0.00 c | 0.00 |
| 3750 | 55.27 e | 38.33 de | 22.30 ab | 25.86 cd | 16.40 bc | 7.48 cd | 7.41 de | 3.846 cd | 0.00 c | 0.00 |
| 7500 | 16.79 f | 16.67 e | 6.78 b | 3.45 d | 0.00 c | 0.00 d | 0.00 e | 0.00 d | 0.00 c | 0.00 |
| 15000 | 11.96 f | 10.00 e | 1.72 b | 0.00 d | 0.00 c | 0.00 d | 0.00 e | 0.00 d | 0.00 c | 0.00 |
| p | <0.001 | <0.001 | 0.011** | <0.001 | 0.001 | <0.001 | <0.001 | <0.001 | <0.001 | - |

-: not calculated

Means that do not share a letter are significantly different ($p < 0.05$).

**:<0.01

It was determined that the methanol extract had a very high scolocidal effect on parasite cells. and no living parasites were found after the 7th hour (Table 7). It is seen that the most effective concentration is 15000 µg/mL as other extracts of *C. crinita*.

It was observed that the NaCl solution used as a scolocidal agent had a very strong scolocidal effect at a concentration of 7500 µg/mL, and no live parasites were found at the 9th . (Table 8).

Determining the scolocidal effect in terms of parasite viability, the difference between the exposure times and the doses of plant extracts was found to be statistically significant ($p < 0.05$). Although this varies according to the extract doses, an interpretation can be made as the dose increases the parasite's viability rate decreases. Again, a decrease was observed in the viability of the parasite, depending on the exposure time.

C. crinita's 15000 µg/mL concentration showed a strong scolocidal effect in all of the Ch, DCM, H, and MeOH solvents. No living parasites were observed at the 7th h with the MeOH extract of *C. crinita* and at the 9th h with the DCM extract (Table 9-10).

From DCM and H extracts of *C. crinita*, the LD₅₀ value of DCM observed in the protoscolices at the 9th h was determined to be 498.5 µg/mL, and its LD₉₀ value was 6800.0 µg/mL; the LD₅₀ value of H extract in protoscolices at 11th h was determined to be 368.2 µg/mL, and its LD₉₀ value was 13675.4 µg/mL (Table 9).

Table 5. Hexane extract of *C. crinita* viability rate comparison at different hours.

| Doses | 3h | 4h | 7h | 9h | 11h | 17h | 20h | 24h | 30h | 32h |
|---------|----------|----------|-----------|-----------|-----------|----------|----------|----------|---------|----------|
| Control | 100.00 a | 100.00 a | 100.00 a | 100.00 a | 100.00 a | 100.00 a | 100.00 a | 100.00 a | 100.0a | 100.0 a |
| 117 | 82.05 b | 78.33 ab | 59.43 b | 55.17 b | 54.80 b | 50.93 b | 48.15 b | 48.08b | 48.92b | 42.92 b |
| 234 | 77.63 bc | 78.33 ab | 57.70 bc | 48.28 bc | 49.30 bc | 47.15 b | 44.44b | 44.23b | 42.83bc | 40.75 b |
| 468 | 68.79 c | 75.00 b | 55.98 bc | 44.83 bcd | 45.63 bc | 41.45 b | 40.74 b | 42.31b | 40.83c | 38.75 b |
| 937 | 58.26 d | 63.33 bc | 42.47 bc | 43.10 bcd | 45.57 bc | 39.67 b | 38.89b | 38.46bc | 38.75c | 32.75 bc |
| 1875 | 52.14 de | 60.00 bc | 39.02 bcd | 36.21 cd | 40.01 bcd | 35.90 b | 25.93 c | 26.92c | 30.58d | 22.50 c |
| 3750 | 46.30 e | 46.67 c | 33.97 cd | 27.59 de | 31.00 bcd | 15.24 c | 5.56 d | 3.65d | 3.67e | 0.41 d |
| 7500 | 25.22 f | 43.33 c | 17.07 de | 17.24 ef | 9.13 cd | 2.81 c | 0.00 d | 0.00d | 0.00e | 0.00 d |
| 15000 | 10.40 g | 4.17 d | 1.53 e | 1.21 f | 0.00 d | 0.00 c | 0.00 d | 0.00 d | 0.00 e | 0.00 d |
| p | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 |

Means that do not share a letter are significantly different ($p < 0.05$)

Table 6. Chloroform extract of *C. crinita* viability rate comparison at different hours.

| Doses | 3h | 4h | 7h | 9h | 11h | 17h | 20h | 24h | 30h | 32h |
|---------|---------|------------|-----------|----------|-----------|-----------|-----------|-------------|----------|----------|
| Control | 100.0 a | 100.00 a | 100.00 a | 100.0 a | 100.00 a | 100.00 a | 100.00 a | 100.00 a | 100.0 a | 100.0 a |
| 117 | 56.83 b | 58.33 b | 59.37 b | 55.17 b | 52.78 b | 45.37 b | 42.59 b | 32.69 b | 30.58 b | 20.50 b |
| 234 | 55.27 b | 53.33 bc | 54.20 bc | 53.45 b | 50.79 bc | 43.45 bc | 35.19 bc | 30.77bc | 26.42 bc | 16.42 b |
| 468 | 52.28 b | 50.00 bcd | 50.92 bc | 50.00 bc | 40.08 bcd | 37.75 bcd | 29.63 cde | 26.92 bc bc | 18.33 cd | 10.25 bc |
| 937 | 47.99 b | 43.33 bcde | 44.14 bc | 43.10 cd | 38.23 bcd | 35.90 bcd | 31.48 cd | 19.23 bcd | 8.167 de | 0.00 c |
| 1875 | 46.43 b | 41.67 bcde | 42.30 bcd | 43.10 cd | 32.74 cde | 32.12 cd | 24.07 de | 15.38 bcd | 6.08 e | 0.00 c |
| 3750 | 39.02 b | 38.33 cde | 38.91 bcd | 36.21 de | 27.18 de | 28.28 d | 20.37 e | 11.54 cd | 0.00 e | 0.00 c |
| 7500 | 34.60 b | 35.00 de | 35.63 cd | 32.76 e | 14.62 ef | 0.00 e | 0.00 f | 0.00 d | 0.00 e | 0.00 c |
| 15000 | 33.17 b | 31.67 e | 20.46 d | 5.17 f | 0.00 f | 0.00 e | 0.00 f | 0.00 d | 0.00 e | 0.00 c |
| p | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 |

Means that do not share a letter are significantly different ($p < 0.05$)

The LD₅₀ value of the *C. crinita* Ch extract observed in protoscolices at the 11th h was determined to be 270.4 µg/mL, and its LD₉₀ value was 10436.6µg/mL; the LD₅₀ value of the M extract observed in the protoscolices at the 7th h was determined to be 32.0 µg/mL, and its LD₉₀ value was 1863.2 µg/mL (**Table 10**).

The LD₅₀ value observed at the 9th hour of the NaCl solution used as a scolicidal agent in protoscolices was determined to be 370.0 µg/mL, and its LD₉₀ value was 4331.7 µg/mL (**Table 11**).

3. Discussion

Essential oils have been used for various purposes over the years. The composition of essential oils is terpenic or non-terpenic volatile compounds. Compounds that make up essential oils are predominately hydrocarbons and oxygenated derivatives of hydrocarbons, but structures such as alcohol, carboxylic acid, ester, aldehyde, ketone, etc. can also be found [33,34].

Terpenes are formed by bonding with isoprene (C₅H₁₀) units. They enter into the essential oil structure in their monoterpene, sesquiterpene, diterpene, and their oxygenated forms. The terpenic compounds in the essential oil are thought to be the reason for their wide range of biological activities. That's why the contents and biological activities of essential oils have become a focus of interest in scientific fields and why relevant studies have grown in importance [35–37].

Table 7. Methanol extract of *C. crinita* viability rate comparison at different hours.

| Doses | 3h | 4h | 7h | 9h | 11h | 17h | 20h | 24h | 30h | 32h |
|----------|------------------|------------------|------------------|------------------|------------------|------------------|------------------|------------------|------------------|--------|
| Control | 100.00 a | 100.0 a | 100.00 a | 100.00 a | 100.00 a | 100.00 s | 100.0 a | 100.0 a | 100.0 a | 100.00 |
| 117 | 47.72 b | 45.00 b | 30.57 b | 27.59 b | 27.31 b | 24.50 b | 24.07 b | 13.46 b | 6.08 b | 0.00 |
| 234 | 46.30 b | 36.67 bc | 27.18 bc | 20.69 bc | 18.19 c | 16.88 b | 11.11 c | 3.85 c | 0.00 c | 0.00 |
| 468 | 37.05 bc | 35.00 bc | 25.40 bc | 15.52 bcd | 12.70 c | 5.63 c | 2.778 d | 0.00 d | 0.00 c | 0.00 |
| 937 | 26.52 cd | 25.00 cd | 18.62 bcd | 12.07 cde | 5.49 d | 2.81 c | 0.00 d | 0.00 d | 0.00 c | 0.00 |
| 1875 | 17.95 cde | 18.33 cde | 11.95 cde | 5.17 de | 0.00 d | 0.00 c | 0.00 d | 0.00 d | 0.00 c | 0.00 |
| 3750 | 16.52 de | 18.33 cde | 6.01 de | 0.00 e | 0.00 d | 0.00 c | 0.00 d | 0.00 d | 0.00 c | 0.00 |
| 7500 | 13.53 de | 6.67 de | 2.53 de | 0.00 e | 0.00 d | 0.00 c | 0.00 d | 0.00 d | 0.00 c | 0.00 |
| 15000 | 4.42 e | 1.50 e | 0.00 e | 0.00 e | 0.00 d | 0.00 c | 0.00 d | 0.00 d | 0.00 c | 0.00 |
| p | <0.001 | - |

-: not calculated

Means that do not share a letter are significantly different ($p < 0.05$).

Table 8. NaCl extract of *C. crinita* viability rate comparison at different hours.

| Doses | 3h | 4h | 7h | 9h | 11h | 17h | 20h | 24h | 30h | 32h |
|----------|------------------|------------------|------------------|------------------|------------------|------------------|------------------|------------------|--------|--------|
| Control | 100.00 a | 100.00 | 100.00 |
| 117 | 71.65 b | 71.67 b | 62.76 b | 60.34 b | 52.84 b | 48.93 b | 37.04 b | 26.92 b | 0.00 | 0.00 |
| 234 | 68.67 b | 63.33 b | 59.37 bc | 53.45 b | 49.21 b | 43.23 b | 31.48 bc | 25.00 bc | 0.00 | 0.00 |
| 468 | 67.10 b | 58.33 bc | 52.59 bc | 46.55 b | 40.15 bc | 35.75 bc | 25.93 bc | 19.23 bcd | 0.00 | 0.00 |
| 937 | 49.15 c | 53.33 bc | 47.47 bc | 37.90 bc | 34.72 bc | 31.98 bcd | 22.22 cd | 17.31 bcd | 0.00 | 0.00 |
| 1875 | 44.73 cd | 40.00 cd | 37.5 cd | 31.0 bc | 23.81 bcd | 16.95 cde | 12.96 de | 9.62 bcd | 0.00 | 0.00 |
| 3750 | 38.75 de | 30.00 de | 23.79 de | 15.52 cd | 12.76 cd | 9.40 de | 7.407 ef | 5.77 cd | 0.00 | 0.00 |
| 7500 | 31.21 e | 25.00 de | 22.07 de | 0.00 d | 0.00 d | 0.00 e | 0.00 f | 0.00 d | 0.00 | 0.00 |
| 15000 | 16.38 f | 11.67 e | 5.11 e | 0.00 d | 0.00 d | 0.00 e | 0.00 f | 0.00 d | 0.00 | 0.00 |
| p | <0.001 | - | - |

-: not calculated

Means that do not share a letter are significantly different ($p < 0.05$).

They have grabbed many scientists' attention due to their wide range of utilization in areas such as cosmetics, medicine, food industry, dentistry, oral care products, perfumery, dyeing, aromatherapy, and phytotherapy. For this reason, the chemical structures and biological activities of essential oils have become a subject of curiosity in recent years [38].

While examining scientific studies, it has been observed that the extraction process has been applied to either fresh or dried algae [39,40]. However, the extracts prepared with fresh samples were found to have higher biological activity than those found in studies with dried samples [41]. The commonly used drying processes used in studies were freeze-drying (lyophilizer) and open-air drying [42–44]. In this study, the algae samples were dried in the open air, under shade, for two to three days after being cleansed of foreign material such as epiphytes and sand.

The compound classes and amounts having the most considerable quantities in the structure of *C. crinita* were found respectively in HD and SPME to be aldehyde (38.88%, 53.89%), terpene and terpenoid compounds (18.684%, 2.52%), and hydrocarbons (10.63%, 36.73%). In the essential oil composition of carboxylic acid compounds, tetradecanoic acid (2.694) and hexadecanoic acid (12.654%) were detected, and the ratio of carboxylic acid compounds in the total essential oil was determined to be 15.35%. Discovered in the essential oil structure derived with HD were the ketone (11.83%) and alcohol (2.24%) class compound structures. The compounds determined in the essential oil composition were collected under a

Table 9. LD₅₀ and LD₉₀ values of *C. crinita* DCM-H extracts per time.

| Time | | Dichloromethane | | | Hexane | | |
|------|------------------|----------------------|----------------------|---------|----------------------|----------------------|---------|
| | | Value | 95% Confidence Limit | | Value | 95% Confidence Limit | |
| | | | Lower | Upper | | Lower | Upper |
| 3 h | LD ₅₀ | 2649.4 ^a | 2592.3 | 2707.4 | 1863.2 ^a | 1813.1 | 1914.3 |
| | LD ₉₀ | 16757.2 ^a | 16187.3 | 17362.8 | 21649.4 ^a | 20631.4 | 22752.9 |
| 4 h | LD ₅₀ | 1755.9 ^a | 1711.5 | 1801.0 | 2401.6 ^a | 2332.1 | 2473.0 |
| | LD ₉₀ | 14567.7 ^a | 14014.0 | 15159.9 | 28833.5 ^a | 27200.5 | 30631.6 |
| 7 h | LD ₅₀ | 874.1 ^a | 849.9 | 898.5 | 544.9 ^a | 521.5 | 568.6 |
| | LD ₉₀ | 7657.7 ^a | 7394.2 | 7938.0 | 13948.2 ^a | 13153.8 | 14825.0 |
| 9 h | LD ₅₀ | 498.5 ^a | 480.4 | 516.8 | 321.0 ^a | 302.9 | 339.4 |
| | LD ₉₀ | 6800.0 ^a | 6525.8 | 7094.9 | 13675.4 ^a | 12786.2 | 14670.6 |
| 11 h | LD ₅₀ | 342.0 ^a | 327.9 | 356.1 | 368.2 ^a | 349.5 | 387.1 |
| | LD ₉₀ | 4811.2 ^a | 4621.3 | 5015.1 | 10679.5 ^a | 10063.5 | 11361.4 |
| 17 h | LD ₅₀ | 185.7 ^a | 175.0 | 196.7 | 289.8 ^a | 274.3 | 305.5 |
| | LD ₉₀ | 5062.6 ^a | 4829.6 | 5315.3 | 8511.8 ^a | 8056.5 | 9012.1 |
| 20 h | LD ₅₀ | 30.3 ^a | 27.0 | 33.8 | 221.9 ^a | 211.5 | 232.5 |
| | LD ₉₀ | 1464.4 ^a | 1399.3 | 1534.3 | 3258.6 ^a | 3133.9 | 3392.3 |
| 24 h | LD ₅₀ | 3.5 ^a | 2.8 | 4.5 | 227.5 ^a | 216.8 | 238.2 |
| | LD ₉₀ | 594.2 ^a | 558.8 | 631.2 | 3150.2 ^a | 3029.4 | 3279.7 |

a: µg/mL

total of 6 classes. Using SPME, volatile components were grouped under three classes: aldehyde, hydrocarbon, and terpene. When examining the effect of the essential oil on a total of five microorganisms, it was observed a high zone formation (14 mm) against gram-positive bacterium *B. cereus* ATCC 10876, and a high zone (13 mm) against the gram-negative bacterium *P. aeruginosa* ATCC 27853. The tobramycin zone value as a standard antibiotic disc was observed to be 18 mm for *B. cereus* and 20 mm for *P. aeruginosa*.

We continued to study the antimicrobial activities of the *C. crinita* extracts in organic solvents. Solutions of 15 and 30 µL for each of the extracts were prepared. Extracts were found to be antimicrobial effective on 8 gram-positive, 11 gram-negative, one fungus, for a total of 20 microorganisms. From the two different concentrations prepared, the extracts with a concentration of 30 µL were observed to form a more effective zone diameter compared to the 15 µL extracts. Extracts with a concentration of 15 µL displayed effects predominately on gram-positive bacteria among 20 microorganisms (*S. aureus* ATCC 29213, MRSA ATCC 67101, *B. cereus* ATCC 10876). Extracts with a concentration of 30 µL generally displayed an effect on all microorganisms in the zone range of 6.5–16 mm except for *Pr. mirabilis* and *E. coli* ATCC 25922. In the extracts with both the 15 and 30 µL concentrations, antimicrobial effects were only observed on *B. cereus* and *P. aeruginosa*.

Plant extracts are highly complex mixtures containing many components. If the disc diffusion method used in this study is indicated by including standard drugs, it can be evaluated. However, the ineffective molecules should be restudied later with dilution methods to eliminate the diffusion problem in the agar method.

Kamenarska et al. (2002) obtained different compounds from the present study in the GC/MS analysis of *C. crinita*, collected from the Aegean Sea at the coast of Kaş. However, the terpene class compounds found by the researchers were similar. This may have been caused by shifting climates and the natural passing of time [26].

Ibtissam et al. (2009), in their study on 32 algae including *C. crinita* in Morocco, the algae samples they examined obtained methanol extracts using the Soxhlet extraction method and examined the antibacterial activity of the extract they obtained. In the study, a zone higher than 20 mm was detected on *S. aureus* ATCC 25923, but it didn't affect other microorganisms [27]. It was observed that the value on *S. aureus* was higher than the results as in our case, which can be explained by the environment, locality, and the subspecies of the algae used.

Table 10. LD₅₀ and LD₉₀ values of *C. crinita* Ch-MeOH extract per time.

| Time | | Chloroform | | | Methanol | | |
|------|------------------|------------------------|----------------------|-----------|---------------------|----------------------|--------|
| | | Value | 95% Confidence Limit | | Value | 95% Confidence Limit | |
| | | | Lower | Upper | | Lower | Upper |
| 3 h | LD ₅₀ | 601.0 ^a | 546.9 | 657.1 | 117.9 ^a | 109.4 | 126.8 |
| | LD ₉₀ | 1542606.0 ^a | 1108426.0 | 2219675.0 | 7693.1 ^a | 7264.5 | 8165.2 |
| 4 h | LD ₅₀ | 441.9 ^a | 396.0 | 489.8 | 90.8 ^a | 83.5 | 98.3 |
| | LD ₉₀ | 1278744.0 ^a | 910975.0 | 1861004.0 | 5093.3 ^a | 4826.0 | 5386.5 |
| 7 h | LD ₅₀ | 543.2 ^a | 502.8 | 584.7 | 32.0 ^a | 28.6 | 35.6 |
| | LD ₉₀ | 187731.0 ^a | 155686.0 | 229673.0 | 1863.2 ^a | 1779.1 | 1953.8 |
| 9 h | LD ₅₀ | 445.6 ^a | 417.1 | 474.8 | 27.6 ^a | 24.8 | 30.6 |
| | LD ₉₀ | 48115.4 ^a | 42865.6 | 54390.6 | 738.1 ^a | 709.8 | 767.9 |
| 11 h | LD ₅₀ | 270.4 ^a | 254.5 | 286.6 | 35.9 ^a | 32.6 | 39.4 |
| | LD ₉₀ | 10436.6 ^a | 9804.9 | 11138.9 | 460.5 ^a | 444.6 | 477.1 |
| 17 h | LD ₅₀ | 203.2 ^a | 189.3 | 217.4 | 2.7 ^a | 2.1 | 3.5 |
| | LD ₉₀ | 10670.7 ^a | 9967.8 | 11459.7 | 856.5 ^a | 804.4 | 911.8 |
| 20 h | LD ₅₀ | 101.1 ^a | 93.2 | 109.3 | 44.1 ^a | 40.4 | 47.7 |
| | LD ₉₀ | 4316.4 ^a | 4094.3 | 4560.0 | 239.1 ^a | 232.2 | 246.2 |
| 24 h | LD ₅₀ | 44.9 ^a | 40.3 | 49.7 | 33.1 ^a | 29.0 | 37.1 |
| | LD ₉₀ | 2299.7 ^a | 2188.5 | 2420.4 | 142.1 ^a | 137.3 | 146.8 |

a: µg/mL

When examining the study, Mhadhebia et al. [45] use various algae extracts prepared with Ch and M to determine the antibacterial activity, including *C. crinita*. The values for the Ch extract of *C. crinita* against *S. aureus* ATCC 25923 and *E. coli* ATCC 35218 in the 6-9 mm zone range are parallel to the values found in our study. However, the antimicrobial values of the M extract were observed to be different.

Selçuk et al. [46] performed different biological activity tests, such as the antiprotozoal, antimycobacterial, and cytotoxic activity for M extracts of algae samples including *C. crinita* and found relatively high values.

In another study, Berber et al. *C. crinita* Duby and *U. intestinalis* seaweed collected from the Sinop coast dried in the open air and coast and prepared methanolic extracts mixing the algae samples in methanol in a water bath for 24 h. They studied the antioxidant and antimicrobial activities of the obtained extracts. Antimicrobial activity was observed against *S. aureus* ATCC 25923 and *S. epidermidis* range of 17–20 mm inhibition zone, while smaller zone diameter was found in our study. The same value antimicrobial activity was found against *B. subtilis* and *E. faecalis* 9–10 mm zone. Although, no antimicrobial activity against *E. coli* and *Ca. albicans* was found in [47], we found.

In this study, due to the components having biological activity in the *C. crinita* structure, it was thought that it is likely to display scolicidal effects along with the antimicrobial activity. Results from the scolicidal effect study of the algae sample extracts in four organic solvents (Ch, DCM, H and M), was observed to decrease over time except for the control group. Additionally, the extract dose prepared at a concentration of 15,000 µg/mL was observed to have a strong scolicidal effect.

According to the reference information reached in different studies, a scolicidal agent, formol, hypertonic glucose solution, alcohol, hypertonic NaCl (3%;10;20), chlorine hexidine, cetrimide, AgNO₃, povidone-iodine, alcohol, 3% H₂O₂, albendazole solution, iodine were used [48–50]. Among the scolicidal agents used, formol, hypertonic saline, and alcohol were found to have the highest risk [51, 52]. In this study, the effect of Ch, DCM, H, and M extracts of *C. crinita* on protoscolices was investigated in parallel using NaCl as a scolicidal agent.

Ozcelik et al. [53] reported that *Allium sativum* was effective on protoscolices. Again, Ozcelik et al. [54] found that propolis also affected daughter vesicles within 10 min. In the study, the M extract of *C. crinita* showed no live parasites at the 7th h. Additionally, the DCM extract showed no parasites at the 9th h. No viable protoscolices were observed at the

Table 11. LD₅₀ and LD₉₀ values of NaCl per time.

| Time | | NaCl | | |
|------|------------------|----------------------|----------------------|---------|
| | | Value | 95% confidence limit | |
| | | | Lower | Upper |
| 3 h | LD ₅₀ | 1305.8 ^a | 1258.1 | 1354.6 |
| | LD ₉₀ | 40220.2 ^a | 37256.1 | 43563.4 |
| 4 h | LD ₅₀ | 907.8 ^a | 872.9 | 943.4 |
| | LD ₉₀ | 23128.4 ^a | 21672.9 | 24746.5 |
| 7 h | LD ₅₀ | 558.6 ^a | 534.3 | 583.2 |
| | LD ₉₀ | 15365.0 | 14468.1 | 16356.5 |
| 9 h | LD ₅₀ | 370.0 ^a | 356.4 | 383.7 |
| | LD ₉₀ | 4331.7 | 4177.7 | 4495.9 |
| 11 h | LD ₅₀ | 245.6 ^a | 234.5 | 256.8 |
| | LD ₉₀ | 3593.4 | 3457.3 | 3739.0 |
| 17 h | LD ₅₀ | 176.8 ^a | 166.4 | 187.4 |
| | LD ₉₀ | 4885.6 ^a | 4662.5 | 5127.3 |
| 20 h | LD ₅₀ | 63.6 ^a | 58.4 | 69.0 |
| | LD ₉₀ | 1971.1 ^a | 1888.1 | 2060.3 |
| 24 h | LD ₅₀ | 23.2 ^a | 20.3 | 26.3 |
| | LD ₉₀ | 1327.9 ^a | 1266.0 | 1394.3 |

a: µg/mL

11th h for the Ch and H extracts. When the application doses were compared to the Tukey test, application dose rates showed a significant change according to the doses ($p < 0.05$). In the study, LD₅₀ and LD₉₀ values were also examined by probit analysis. Accordingly, the LD₅₀ value observed in the protoscolices at the 9th h of DCM extract of *C. crinita* was determined to be 498.5 µg/mL; LD₉₀ value of H extract was determined to be 6800.0µg/mL, LD₅₀ value observed at the 11th h as 368.2 µg/mL; LD₉₀ value was 13675.4µg/mL, the LD₅₀ value observed in protoscolices at the 11th h of Ch extract was 270.4 µg/mL; LD₉₀ value was 10436.6; (please remove coma) µg/mL; LD₅₀ value was 32.0 µg/mL, and LD₉₀ value was 1863.2 µg/mL at the 7th h of M extract. With this information in mind, it can be concluded that *C. crinita* extracts can be used as potential scolical agents.

4. Conclusion

The volatile components of *C. crinita* were observed with both HD and SPME in the GC-MS/FID device, and their structures were clarified in this study. Terpene and terpenoids, carboxylic acid, aldehyde class compounds, which were not identified in previous studies, were found to be present here. In the antimicrobial activity study of essential oils, *C. crinita* was observed to have a high activity level on our study's microorganisms. Most of the extracts prepared with organic solvents were affected with a value of 6.5 to 16 mm zone diameter.

Furthermore, organic solvent extracts of *C. crinita* were determined to have an additional use as bactericidal and scolical agents. Thus, it was concluded that LD₉₀ dose can be increased for a faster effect or decreased for a slower effect. For *C. crinita* to be used on living cells, it is recommended that controlled experiments are performed with experimental animals and that dosage levels are made to adjust the amount of antimicrobial effect.

Acknowledgments

The author thanks Assoc. Dr. Ülkü Karaman for scolical analysis, Ass. Dr. Tuğba Cebeci for antimicrobial activity and Ass. Dr. Yeliz Kaşko Arıcı for statistical analysis. This study was supported by a grant from Giresun University, Turkey (GRU-FEN-BAP-A-170417-111).

References

1. Miyashita K, Mikamia N, Hosokawa M. Chemical and nutritional characteristics of brown seaweed lipids: A review. *Journal of Functional Foods* 2013; 5: 1507-1517. doi: 10.1016/j.jff.2013.09.019
2. Aktar S, Cebe GE. General specifications, using areas of algae and their importance on pharmacy. *Journal of Faculty of Pharmacy of Ankara University* 2010; 39 (3):237-264. doi: 10.1501/Eczfak_0000000568
3. Olasehinde AT, Olaniran AO, Okoh AI. Macroalgae as a valuable source of naturally occurring bioactive compounds for the treatment of Alzheimer's Disease. *Marine Drugs* 2019; 17: 609-627. doi: 10.3390/md17110609
4. Taşkin E, Öztürk M, Kurt O, Öztürk M. Check-list of Marine Flora of Turkey. 7th International Phycological Congress, *Phycologia* 40 (4), p.71. 18-25 August 2001 Thessaloniki, Greece.
5. Brownlee IA, Allen A, Pearson JP, Dettmar PW, Havler ME et al. Alginate as a source of dietary fiber. *Critical Reviews in Food Science and Nutrition* 2005; 45: 497-510. doi: 10.1080/10408390500285673
6. Al-Saif SSA, Abdel-Raouf N, El-Wazanani HA, Aref IA. Antibacterial substances from marine algae isolated from Jeddah coast of Red sea. Saudi Arabia. *Saudi Journal of Biological Science* 2014; 21 (1):57-64. doi: 10.1016/j.sjbs.2013.06.001
7. Senthilkumar K, Manivasagana P, Venkatesana J, Kim SK. Brown seaweed fucoidan: Biological activity and apoptosis, growth signaling mechanism in cancer. *International Journal of Biological Macromolecules* 2013; 60: 366-374. doi: 10.1016/j.ijbiomac.2013.06.030
8. Sanjeeva KKA, Kima EA, Son KT, Jeon YJ. Bioactive properties and potentials cosmeceutical applications of phlorotannins isolated from brown seaweeds: A review. *Journal of Photochemistry and Photobiology B: Biology* 2016; 162: 100-105. doi: 10.1016/j.jphotobiol.2016.06.027
9. Laurienzo P. Marine polysaccharides in pharmaceutical applications: an overview. *Marine Drugs* 2010; 8: 2435-2465. doi: 10.3390/md8092435.
10. Manial A, Sujith S, Selvin J, Shakir C, Kiran GS et al. Antibacterial activity of *Falkenbergia hillebrandii* (Born) from the Indian coast against human pathogeny. *Int. Journal of Experimental Botany* 2009; 78: 161-166.
11. Pereira HS, Lea-o-Ferreira LR, Moussatche N, Teixeira VL, Cavalcanti DN et al. Antiviral activity of diterpenes isolated from the Brazilian marine alga *Dictyota menstrualis* against human immunodeficiency virus type 1 (HIV-1). *Antiviral Research* 2004; 64: 69-76. doi: 10.1016/j.antiviral.2004.06.006
12. Zerrifi SA, Khalloufi EF, Oudra B, Vasconcelos V. Seaweed bioactive compounds against pathogens and microalgae: potential uses on pharmacology and harmful algae bloom control. *Marine Drugs* 2018; 16 (2): 55-65. doi: 10.3390/md16020055
13. Duan E. Bazı Deniz Makroalglerinden (*Ulva* sp., *Cystoseira* sp. ve *Corallina* sp.) Fermente Sıvı Organik Gübre Üretimi ve Taze Fasülye (*Phaseolus vulgaris*) Verimine Etkisinin Belirlenmesi", Giresun Üniversitesi, Fen Bilimleri Enstitüsü, Yüksek Lisans Tezi, 2013, pp. 64, Giresun, in Turkish.
14. Agregán R, Munekata PES, Franco D, Carballo J, Barba FJ et al. Antioxidant potential of extracts obtained from macro- (*Ascophyllum nodosum*, *Fucus vesiculosus* and *Bifurcaria bifurcata*) and micro-algae (*Chlorella vulgaris* and *Spirulina platensis*) assisted by ultrasound. *Medicines (Basel)* 2018; 5: 33-42. doi: 10.3390/medicines5020033
15. Kılıç A. Methods of essential oil production. *Journal of Bartın Faculty of Forestry* 2008; 10 (13): 37-45.
16. Rodríguez AGG, Portilla CJ, Bañuelos TO, Zepeda RC. Anticancer activity of seaweeds. *Drug Discovery Today* 2018; 23 (2): 434-445. doi: 10.1016/j.drudis.2017.10.019
17. A.Arslan B, Yilancioglu K, Kalkan Z, Timucin AC, Gür H et al. Screening of new antileukemic agents from essential oils of algae extracts and computational modeling of their interactions with intracellular signaling nodes. *European Journal of Pharmaceutical Sciences* 2016; 83: 120-131. doi: 10.1016/j.ejps.2015.12.001
18. Moubayed MS, Al Hourri HJ, Khulaifi MMA, Al Farraj DA. Antimicrobial, antioxidant properties and chemical composition of seaweeds collected from Saudi Arabia (Red Sea and Arabian Gulf). *Saudi Journal of Biological Sciences* 2017; 24: 162-169. doi: 10.1016/j.sjbs.2016.05.018
19. Roohinejad S, Koubaa M, Barba JF, Saljoughian S, Amid M et al. Application of seaweeds to develop new food products with enhanced shelf-life, quality and health-related beneficial properties. *Food Research International* 2017; 99: 1066-1083. doi: 10.1016/j.foodres.2016.08.016
20. Gümüş B, Ünlüsayın M, Gümüş E. Determination of antimicrobial activity of two macro algae extracts. *Ege Journal of Fisheries and Aquatic Sciences* 2016; 33 (4): 389-395. doi: 10.12714/egejfas.2016.33.4.13
21. Asakawa Y, Toyota M, Takemoto T, Kubo I, Nakanishi K. Insect antifeedant secoaromadendrane-type sesquiterpenes from *Plagiochila* species. *Phytochemistry* 1980; 19: 2147-2154.
22. Cho JY, Kwon EH, Choi JS, Hong SY, Shin HW et al. Antifouling activity of seaweed extracts on the green algae *Enteromorpha prolifera* and the mussel *Mytilus edulis*. *Journal of Applied Phycology* 2001; 13 (2): 117-125.

23. Mayer AM, Rodríguez AD, Berlinck RG, Fusetani N. Marine pharmacology in 2007–8: Marine compounds with antibacterial, anticoagulant, antifungal, anti-inflammatory, antimalarial, antiprotozoal, antituberculosis and antiviral activities; affecting the immune and nervous system, and other miscellaneous mechanisms of action. *Comparative Biochemistry and Physiology Part C: Toxicology & Pharmacology* 2011; 153 (2): 191-222. doi: 10.1016/j.cbpc.2010.08.008
24. Davis G, Alvarez SM, Ochoa IM, Torres MC, Padron EG et al. Bactericide, Antioxidant and Cytotoxic Activities from Marine Algae of Genus *Laurencia* Collected in Baja California Sur, Mexico. *International Journal of Pharmacology* 2018; 14 (3): 391-396. DOI: 10.3923/ijp.2018.391.396
25. Zbakh H, Chiheb I, Motilva V, Riadi H. Antibacterial, cytotoxic and antioxidant potentials of *Cladophora prolifera* (Roth) Kützing collected from the Mediterranean coast of Morocco. *American Journal of Phytomedicine and Clinical Therapeutics* 2014; 2: 1187–1199.
26. Kamenarskaa Z, Yalçın FN, Ersöz T, Çalış İ, Stefanov K et al. Chemical Composition of *Cystoseira crinita* Bory from the Eastern Mediterranean. *Z. Naturforsch. C Biosci.*, 2002; 57 (7-8): 584-590. doi: 10.1515/znc-2002-7-806
27. Ibtissam C, Hassane R, José ML, Francisco DSJ, Antonio GVJ et al. Screening of antibacterial activity in marine green and brown macroalgae from the coast of Morocco. *African Journal of Biotechnology* 2009;8 (7): 1258-1262.
28. Renda G, Özel A, Barut B, Korkmaz B, Yayli N. The volatile chemical compositions of the essential oil/spme and enzyme inhibitory and radical scavenging activities of solvent extracts and the essential oils from *Coronilla orientalis* miller and *C. varia* L. grows in Turkey. *Iranian Journal of Pharmaceutical Research* 2019; 18 (4): 1831-1842. doi: 10.22037/ijpr.2019.1100802
29. Demirel Z, Y.Koz, FF, K.Yavaşoğlu NU, Özdemir G, Sukatar A. Antimicrobial and antioxidant activities of solvent extracts and the essential oil composition of *Laurencia obtusa* and *Laurencia obtusa* var. *pyramidata*. *Romanian Biotechnological Letters* 2011; 16 (11): 5927-5936.
30. Filice C, Pirola F, Brunetti E, Dughetti S, Strosselli M et al. A new therapeutic approach for hydatid liver cysts. Aspiration and alcohol injection under sonographic guidance. *Gastroenterology* 1990; 98: 1366-1368. doi: 10.1016/0016-5085(90)90358-8
31. Yagci G, Ustunsoz B, Kaymakcioglu N, Bozlar U, Gorgulu S et al. Results of surgical, laparoscopic, and percutaneous treatment for hydatid disease of the liver: 10 years experience with 355 patients. *World Journal of Surgery* 2005; 29 (12): 1670-1679. DOI: 10.1007/s00268-005-0058-1
32. N.Moghaddam S, Abrishami A, Taefi A, Malekzadeh R. Percutaneous needle aspiration, injection, and re-aspiration with or without benzimidazole coverage for uncomplicated hepatic hydatid cysts. *Cochrane Database of Systematic Reviews* 2011; 19 (1): CD003623. doi: 10.1002/14651858.CD003623.pub3
33. Klegin C, Moura NF, Sousa MHO, Frassini R, E.Ely M. et al. Chemical composition and cytotoxic evaluation of the essential oil of *Phyllogonium viride* (Phyllogoniaceae, Bryophyta). *Chemistry & Biodiversity* 2021; 18: e2000794 2-8. doi: 10.1002/cbdv.202000794.
34. Amorati R, Foti CM, Valgimigli L. Antioxidant Activity of Essential Oils. *Journal of Agricultural and Food Chemistry* 2013; 61 (46): 10835-10847. doi: 10.1021/jf403496k
35. Karadağ AE, Demirci B, Kültür Ş, Demirci F, Başer KH. Antimicrobial, anticholinesterase evaluation and chemical characterization of essential oil *Phlomis kurdica* Rech. fil. Growing in Turkey. *Journal of Essential Oil Bearing Plants* 2019; 32 (3): 242-246. doi: 10.1080/10412905.2020.1743786
36. Yilar M, Bayar Y, A.Bayar AA. Allelopathic and Antifungal potentials of endemic *Salvia absconditiflora* Greuter & Burdet collected from different locations in Turkey. *Allelopathy Journal* 2020; 49 (2): 243-256. doi: 10.26651/allelo.j/2020-49-2-1268
37. Briskin DP. Medicinal plants and phytomedicines: Linking plant biochemistry and physiology to human health. *Plant Physiology* 2000; 124 (2): 507-514. doi: 10.1104/pp.124.2.507
38. Xiao S, Yu H, Xie Y, Guo Y, Fan J et al. The anti-inflammatory potential of *Cinnamomum camphora* (L.) J. Presl essential oil in vitro and in vivo. *Journal of Ethnopharmacology* 2021; 267: 113516. doi: 10.1016/j.jep.2020.113516
39. Spavieri J, Allmendinger A, Kaiser M, Itoe MA, Blunden G. et al. Assessment of Dual Life Stage Antiplasmodial Activity of British Seaweeds. *Marine Drugs* 2013; 11 (10): 4019–4034. doi: 10.3390/md11104019
40. Orhan I, Sener B, Atıcı T, Brun R, Perozzo R et al. Turkish freshwater and marine macrophyte extracts show in vitro antiprotozoal activity and inhibit FabI, a key enzyme of *Plasmodium falciparum* fatty acid biosynthesis. *Phytomedicine* 2006; 13 (6): 388-393.
41. Manivannan K, Karthikai DG, Anantharaman P, Balasubramanian T. Antimicrobial potential of selected brown seaweeds from Vedalai coastal waters, Gulf of Mannar. *Asian Pacific Journal of Tropical Biomedicine* 2011; 1 (2): 114-120. doi: 10.1016/S2221-1691(11)60007-5
42. Kim YH, Kim JH, Jin HJ, Lee SY. Antimicrobial activity of ethanol extracts of *Laminaria japonica* against oral microorganism. *Anaerobe* 2013; 21: 34-38. doi: 10.1016/j.anaerobe.2013.03.012
43. Peres JCF, D.Carvalho LR, Gonçalves E, Berian LOS, Felicio JD. Evaluation of antifungal activity of seaweed extracts. *Agricultural Sciences* 2012; 36 (3): 294-299. doi: 10.1590/S1413-70542012000300004

44. Ely R, Supriya T, Naik CG. Antimicrobial activity of marine organisms collected off the coast of South East India. *Journal of Experimental Marine Biology and Ecology* 2004; 309: 121-127. doi: 10.1016/j.jembe.2004.03.010
45. Mhadhebia L, Chaieb K, Bouraoui A. Evaluation of antimicrobial activity of organic fractions of six marine algae from tunisian mediterranean coasts. *International Journal of Pharmacy and Pharmaceutical Sciences* 2012; 4 (1): 534-537.
46. S.Selçuk S, Meriçli AH, Güven KC, Kaiser M, Casey R et al. Evaluation of Turkish seaweeds for antiprotozoal, antimycobacterial and cytotoxic activities. *Phytotherapy Research* 2011; 25 (5): 778-83. doi: 10.1002/ptr.3330
47. Berber İ, Avşar C and Koyuncu H. Antimicrobial and antioxidant activities of *Cystoseira crinita* Duby and *Ulva intestinalis* Linnaeus from the coastal region of Sinop, Turkey. *Journal of Coastal Life Medicine* 2015; 3 (6):441-445. doi: 10.12980/JCLM.3.2015JCLM-2015-0013
48. Xiao SH, Yang YO, Guo HF, Zhang CW, Jiao PY et al. Effects of mebendazole, albendazole and albendazole sulfoxide on glycogen contents of *Echinococcus granulosus* cysts harbored in mice. *Chung Kuo Yao Li Hsueh Pao* 1990; 11: 546-549.
49. Moreno MJ, U.París MA, Casado N, Caabeiro FR. Praziquantel and albendazole in the combined treatment of experimental hydatid disease. *Parasitol Research* 2001; 87 (3): 235-238. doi: 10.1007/s004360000334
50. Yetim I, Erzurumlu K. Current Approaches in Treatment of Hydatid Cysts of Liver. *Journal of Clinical and Analytical Medicine* 2013; 4 (1): 64-71.
51. Belghiti J, Benhamou JP, Houry S, Grenier P, Huguier M. et al. Caustic sclerosing cholangitis. *Archives of Surgery* 1986; 121 (10): 1162-1165. doi: 10.1001/archsurg.1986.01400100070014
52. Kathkouda N, Fabiani P, Benizri E, Mouiel J. Laser resection of a liver hydatid cyst under videolaparoscopy. *British Journal of Surgery* 1992; 79 (6): 560-1. doi: 10.1002/bjs.1800790628
53. Özcelik S, Sümer Z, Değerli S, Ozan F, Sökmen A. Can Garlic (*Allium sativum*) Extract Used As Scolocidal Agent?. *Turkish Journal of Parasitology* 2007; 31 (4): 318-321 (in Turkish).
54. Özcelik S, Sümer Z, Değerli S. The scolocidal effect of propolis on protoscoleces and daughter cysts. *Cumhuriyet Medical Journal* 2015; 37 (1): 4-8. doi: 10.7197/CMJ.V37I1.5000071890