

# Generalizations of Zassenhaus lemma and Jordan-Hölder theorem for 2–crossed modules

Selim ÇETİN\*

\*Department of Mathematics, Faculty of Arts & Humanities, Burdur Mehmet Akif Ersoy University, Burdur, Türkiye  
 ORCID iD: <https://orcid.org/0000-0002-9017-1465>

Received: .202

Accepted/Published Online: .202

Final Version: ..202

**Abstract:** We present a comprehensive generalization of the Zassenhaus Lemma, the Schrier Refinement Theorem, and the Jordan-Hölder Theorem, extending their applicability to both crossed modules and 2–crossed modules. It is discovered that the previously established normality conditions are insufficient for forming quotient objects of 2–subcrossed modules, as demonstrated through an illustrative example, and these conditions are rigorously revised to allow these generalizations. These new conditions now yield results that are categorically accurate. Additionally, the study has led to the derivation of several supplementary results, including isomorphism theorems for 2–crossed modules.

**Key words:** crossed module, normality, Zassenhaus lemma, isomorphism theorems.

## 1. Introduction and Preliminaries

Crossed modules, in conjunction with their topological background, constitute structures that are algebraically widespread and significant. One of the most notable reasons for their algebraic importance is the fact that any group  $G$  can be analyzed through the crossed modules  $(T, G, \partial)$  that can be established on it. This situation have some similarities with the intimate connection between modules and rings, as it was stated in [22] that to study on rings without some reference to modules is inconceivable.

First introduced by Whitehead [23, 24], crossed module are structures that model the homotopy 2-type, just as homotopy 1-types can be classified by groups. While crossed modules, can be studied within the framework of homotopy theory, which is the origin of the concept, they can also be studied with categorical or pure algebraic approaches. In fact, Lue and Norrie have addressed crossed modules from the axiomatic algebraic perspective and showed in their studies that many concepts in group theory surprisingly correspond to similar concepts in crossed modules, which have sufficiently well-behaved properties [13, 19]. Thus developing an interesting and comprehensive algebraic theory of crossed modules have been possible, which directly generalize the theory of groups in many aspects.

Crossed modules, which are one of the best-known structures of higher-dimensional group theory among with others [1, 2, 4–10, 14–17] and which have many naturally equivalent structures as Cat –1 groups, simplicial groups, strict 2–groups, have many equivalent definitions that make use of different mathematical tools. For the purpose of the study, here we prefer to give the axiomatic definition, which is most commonly encountered in the literature.

\*Correspondence: selimcetin@gmail.com

2020 AMS Mathematics Subject Classification: 18G45, 20E34, 20E32, 20E07

A crossed module is a triple  $(T, G, \partial)$ , where  $T$  is a  $G$ -group (i.e. the group  $G$  acts on the group  $T$  by automorphisms) and  $\partial : T \rightarrow G$  is a  $G$ -equivariant group homomorphism satisfying the Peiffer rule. More explicitly  $\partial$  fulfills the properties

$$\text{CM1)} \quad \partial(g \triangleright t) = g \partial t g^{-1}$$

$$\text{CM2)} \quad \partial(s) \triangleright t = s t s^{-1}$$

for all  $s, t \in T$  and  $g \in G$ , where  $g \triangleright t$  stands for the action of the element  $g$  on  $t$ .

Crossed modules generalize both the notion of a group and the notion of a normal subgroup of a group. Indeed any group  $G$  can naturally be identified with both the crossed modules  $(1, G, inc)$  and  $(G, G, id)$ , where in the latter crossed module  $G$  acts on itself by conjugation, and these identifications specify two fully faithful functors from the category of groups  $\mathfrak{Gr}$  to the category of crossed modules  $\mathfrak{XMod}$ . On the other hand, any normal subgroup  $N$  of a group  $G$  naturally gives rise to the crossed module  $(N, G, inc)$ , where  $G$  acts on  $N$  by conjugation, and as a crossed module  $(T, G, \partial)$  is sometimes referred as a  $G$ -crossed module, a normal subgroup of  $G$  defines a  $G$ -crossed module, making possible to interpret  $G$ -crossed modules as generalizations of normal subgroups of  $G$ .

In internal viewpoint, it is also possible to define subobjects and subobjects normal subobjects in  $\mathfrak{XMod}$ . Given two crossed modules  $(T, G, \partial)$  and  $(S, H, \partial')$ . We call  $(S, H, \partial')$  a subcrossed module of  $(T, G, \partial)$  if  $S$  and  $H$  are subgroups of  $T$  and  $G$ , respectively,  $\partial'$  is the restriction of  $\partial$  and also the action of  $H$  on  $S$  is inherited from the action of  $G$  on  $T$ . For practical purposes, we often denote also the restriction of  $\partial$  with again  $\partial$ . The situation, where  $(S, H, \partial)$  is a subcrossed module of  $(T, G, \partial)$  is denoted with  $(S, H, \partial) \leq (T, G, \partial)$ . In this case, the normality of  $(S, H, \partial)$  in  $(T, G, \partial)$  is defined by the conditions

i)  $H$  is normal subgroup of  $G$ ,

ii)  $g \triangleright s \in S$  for all  $g \in G$  and  $s \in S$ ,

iii)  $(h \triangleright t)t^{-1} \in S$  for all  $h \in H$  and  $t \in T$ ,

and denoted by  $(S, H, \partial) \trianglelefteq (T, G, \partial)$ .

If  $(S, H, \partial) \trianglelefteq (T, G, \partial)$ , then we have by definition  $H \trianglelefteq G$ . Additionally, it is easily seen that also  $S \trianglelefteq T$ . Thus it is possible to form a crossed module  $(T/S, G/H, \bar{\partial})$ , where  $\bar{\partial}(tS) = \partial(t)H$  and  $G/H$  acts on  $T/S$  by the formula  $(gH) \triangleright (tS) = (g \triangleright t)S$ . This is called the quotient crossed module of  $(T, G, \partial)$  by  $(S, H, \partial)$  and is denoted by  $\frac{(T, G, \partial)}{(S, H, \partial)}$ .

A crossed module homomorphism from a crossed module  $(T, G, \partial)$  to a crossed module  $(T', G', \partial')$  is a pair  $(\theta, \sigma)$  of group homomorphisms  $\theta : T \rightarrow T'$ ,  $\sigma : G \rightarrow G'$ , such that  $\sigma \partial = \partial' \theta$  and  $\theta(g \triangleright t) = \sigma(g) \triangleright \theta(t)$  for all  $g \in G$  and  $t \in T$ . Therefore a crossed module homomorphism can be summarized by the following commutative diagrams.

$$\begin{array}{ccc} T & \xrightarrow{\partial} & G \\ \theta \downarrow & & \downarrow \sigma \\ T' & \xrightarrow{\partial'} & G' \end{array} \quad \begin{array}{ccc} G \times T & \xrightarrow{\triangleright} & T \\ (\sigma, \theta) \downarrow & & \downarrow \theta \\ G' \times T' & \xrightarrow{\triangleright'} & T' \end{array}$$

Such an crossed module homomorphism defines a normal subcrossed module  $(\ker \theta, \ker \sigma, \partial)$  of  $(T, G, \partial)$  and a subcrossed module  $(\operatorname{im} \theta, \operatorname{im} \sigma, \partial')$  of  $(T', G', \partial')$ . While the former is called the kernel of the crossed module homomorphism  $(\theta, \sigma)$ , the latter is called the image of  $(\theta, \sigma)$ .

It worths noting that, there is a close relationship between the concepts of a normal subcrossed module and the kernel of a crossed module homomorphism. Namely, as we stated in the previous paragraph, the kernel of a crossed module homomorphism  $(\theta, \sigma) : (T, G, \partial) \longrightarrow (T', G', \partial')$  is a normal subcrossed module of  $(T, G, \partial)$  and conversely, any normal subcrossed module  $(S, H, \partial)$  of  $(T, G, \partial)$  gives rise a crossed module homomorphism  $(\theta_q, \sigma_q) : (T, G, \partial) \longrightarrow (T/S, G/H, \bar{\partial})$  defined as  $\theta_q(t) = tS$ ,  $\sigma_q(g) = gH$ , whose kernel is equal to  $(S, H, \partial)$ .

As a significant result of the discussion above, it can be stated that normal subcrossed modules of a crossed module  $(T, G, \partial)$  are exactly the kernels of crossed module homomorphisms originating from  $(T, G, \partial)$ . This provides us an important criterion for defining normal objects. Extending this approach from the category of crossed modules to the category of 2-crossed modules is the reason why, we need to revise the definition of the normal 2-subcrossed module in the literature. Indeed, a counterexample is also presented in this article, illustrating that the present definition may not always be sufficient to construct a well-defined quotient 2-crossed module. For more details, see Example 3.11.

In the context of high-dimensional group theory 2-crossed modules, which are first introduced by Conduché in [3], model 3-groups, just as crossed modules model 2-groups [18]. Some fundamental definitions regarding 2-crossed modules are given in Section 3. Despite their seemingly complex structure, it is quite interesting to encounter examples of 2-crossed modules in natural sciences, for instance in axion electrodynamics in the framework of particle physics [11].

Isomorphism theorems are well-known major results of group theory. As crossed modules generalize the group theory, the question arises whether if one can generalize isomorphism theorems to crossed modules. The answer is positive, and in fact the isomorphism theorems for crossed modules have really been well established and found their places in the literature. Relevant results can be found collectively in [19]. One way to further generalize these theorems is to investigate the validity of the results for 2-crossed modules. A part of the current work includes to identify the structures that correctly describe the quotient objects of the 2-crossed modules, and then to prove the generalizations of the isomorphism theorems.

Some of important mathematical results that are somewhat related to the isomorphism theorems are Zassenhaus lemma, Schreier refinement theorem and Jordan-Hölder theorem. Zassenhaus lemma is also known as Butterfly lemma in the literature, and this nomenclature is based on the shape of the Hasse diagram for the lattice of the subgroups involved in the lemma, given by Lang [12]. Schreier refinement theorem is an important result on refinements of subnormal series of subgroups of a group, whose proof depends to Zassenhaus lemma. The Jordan-Hölder theorem, on the other hand, is an important result that can be qualified as the fundamental theorem for the composition series of groups. Some of the outputs of this article are the generalizations of the above mentioned lemma and theorems, which have an important role in the group theory, both in to the theory of crossed modules and the theory of 2-crossed modules.

## 2. The Case of Crossed Modules

**Lemma 2.1** *Given two subcrossed modules  $(S, H, \partial)$ ,  $(R, K, \partial)$  of a crossed module  $(T, G, \partial)$ .  $(P, N, \partial)$  and  $(Q, M, \partial)$  be normal subcrossed modules of  $(S, H, \partial)$  and  $(R, K, \partial)$ , respectively. Then*

$$(P, N, \partial)((S, H, \partial) \cap (Q, M, \partial))$$

is a normal subcrossed module of

$$(P, N, \partial)((S, H, \partial) \cap (R, K, \partial))$$

and

$$(Q, M, \partial)((P, N, \partial) \cap (R, K, \partial))$$

is a normal subcrossed module of

$$(Q, M, \partial)((S, H, \partial) \cap (R, K, \partial)).$$

Moreover,

$$\frac{(P, N, \partial)((S, H, \partial) \cap (R, K, \partial))}{(P, N, \partial)((S, H, \partial) \cap (Q, M, \partial))} \cong \frac{(Q, M, \partial)((S, H, \partial) \cap (R, K, \partial))}{(Q, M, \partial)((P, N, \partial) \cap (R, K, \partial))}.$$

**Proof** From the definitions of intersection of two subcrossed modules, product of a subcrossed module and a normal subcrossed module and quotient crossed module we must show that

$$\left( \frac{P(S \cap R)}{P(S \cap Q)}, \frac{N(H \cap K)}{N(H \cap M)}, \bar{\partial}_1 \right) \cong \left( \frac{Q(S \cap R)}{Q(P \cap R)}, \frac{M(H \cap K)}{M(N \cap K)}, \bar{\partial}_2 \right).$$

For the purpose we consider the crossed module

$$\frac{(S, H, \partial) \cap (R, K, \partial)}{((P, N, \partial) \cap (R, K, \partial))((S, H, \partial) \cap (Q, M, \partial))} = \left( \frac{S \cap R}{(P \cap R)(S \cap Q)}, \frac{H \cap K}{(N \cap K)(H \cap M)}, \bar{\partial} \right).$$

1 It is known from the group theory that the functions

$$\begin{aligned} \theta &: \frac{P(S \cap R)}{P(S \cap Q)} \longrightarrow \frac{S \cap R}{(P \cap R)(S \cap Q)} \\ \sigma &: \frac{N(H \cap K)}{N(H \cap M)} \longrightarrow \frac{H \cap K}{(N \cap K)(H \cap M)} \end{aligned}$$

2 given by  $\theta(ptP(S \cap Q)) = t(P \cap R)(S \cap Q)$  and  $\sigma(ngN(H \cap M)) = g(N \cap K)(H \cap M)$  are group isomorphisms  
3 [21].

4 Then for each  $ptP(S \cap Q) \in \frac{P(S \cap R)}{P(S \cap Q)}$ ,

$$\begin{aligned} \sigma \bar{\partial}_1(ptP(S \cap Q)) &= \sigma(\partial(pt)N(H \cap M)) = \sigma(\partial p \partial t N(H \cap M)) \\ &= \partial t(N \cap K)(H \cap M) = \bar{\partial}(t(P \cap R)(S \cap Q)) = \bar{\partial} \theta(ptP(S \cap Q)) \end{aligned}$$

5 and for each  $ngN(H \cap M) \in \frac{N(H \cap K)}{N(H \cap M)}$ ,

$$\begin{aligned} \theta(ngN(H \cap M) \triangleright ptP(S \cap Q)) &= \theta((ng \triangleright pt)P(S \cap Q)) \\ &= \theta((ng \triangleright pt)(g \triangleright t)^{-1}(g \triangleright t)P(S \cap Q)) \\ &= \theta((ng \triangleright p)(n \triangleright (g \triangleright t))(g \triangleright t)^{-1}(g \triangleright t)P(S \cap Q)) \\ &= \theta((p'(g \triangleright t)P(S \cap Q)) = (g \triangleright t)(P \cap R)(S \cap Q)) \\ &= g(N \cap K)(H \cap M) \triangleright t(P \cap R)(S \cap Q) \end{aligned}$$

$$= \sigma\left(\text{ng}N(H \cap M)\right) \triangleright \theta\left(\text{pt}P(S \cap Q)\right)$$

1 which proves that  $(\theta, \sigma)$  is a crossed module isomorphism.

2  
3 The isomorphism  $\left(\frac{Q(S \cap R)}{Q(P \cap R)}, \frac{M(H \cap K)}{M(N \cap K)}, \overline{\partial_2}\right) \cong \left(\frac{S \cap R}{(P \cap R)(S \cap Q)}, \frac{H \cap K}{(N \cap K)(H \cap M)}, \overline{\partial}\right)$  is similarly seen from the  
4 symmetry.  $\square$

5 **Definition 2.2** [19] A normal series (series of lenght  $n$ ) of a crossed module  $(T, G, \partial)$  consists of subcrossed  
6 modules  $(T_i, G_i, \partial)$ ,  $i = 0, \dots, n$ , such that,  $(T_i, G_i, \partial) \trianglelefteq (T_{i-1}, G_{i-1}, \partial)$  for  $i = 1, \dots, n$  and

$$\begin{array}{ccccccc} 1 = T_n & \longrightarrow & T_{n-1} & \longrightarrow & \cdots & \longrightarrow & T_1 & \longrightarrow & T_0 = T \\ \downarrow & & \downarrow & & & & \downarrow & & \downarrow \partial \\ 1 = G_n & \longrightarrow & G_{n-1} & \longrightarrow & \cdots & \longrightarrow & G_1 & \longrightarrow & G_0 = G \end{array}$$

In this case the quotient crossed modules

$$\frac{(T_0, G_0, \partial_0)}{(T_1, G_1, \partial_1)}, \frac{(T_1, G_1, \partial_1)}{(T_2, G_2, \partial_2)}, \dots, \frac{(T_{n-1}, G_{n-1}, \partial_{n-1})}{(T_n, G_n, \partial_n)}$$

7 are called factor crossed modules of  $(T, G, \partial)$ .

8 **Definition 2.3** Given two normal series

$$\begin{array}{ccccccc} 1 = T_n & \longrightarrow & T_{n-1} & \longrightarrow & \cdots & \longrightarrow & T_1 & \longrightarrow & T_0 = T \\ \downarrow & & \downarrow & & & & \downarrow & & \downarrow \partial \\ 1 = G_n & \longrightarrow & G_{n-1} & \longrightarrow & \cdots & \longrightarrow & G_1 & \longrightarrow & G_0 = G \end{array}$$

9 and

$$\begin{array}{ccccccc} 1 = T_{i_m} & \longrightarrow & T_{i_{m-1}} & \longrightarrow & \cdots & \longrightarrow & T_{i_1} & \longrightarrow & T_{i_0} = T \\ \downarrow & & \downarrow & & & & \downarrow & & \downarrow \partial \\ 1 = G_{i_m} & \longrightarrow & G_{i_{m-1}} & \longrightarrow & \cdots & \longrightarrow & G_{i_1} & \longrightarrow & G_{i_0} = G \end{array}$$

1 of a crossed module  $(T, G, \partial)$ . If  $i_0 = 0$ ,  $i_m = n$  and  $i_{k-1} < i_k$  for each  $k = 1, \dots, m$ , then the former series  
 2 is said to be a refinement of the latter one.

3 **Definition 2.4** A crossed module  $(T, G, \partial)$  is called simple, if its only normal subcrossed modules are  $(1, 1, 1)$   
 4 and  $(T, G, \partial)$ .

5 **Definition 2.5** A normal series of crossed modules, of which all nontrivial factor crossed modules are simple,  
 6 is called composition series.

7 **Definition 2.6** Two normal series of a crossed module  $(T, G, \partial)$  are said to be equivalent, if for each factor  
 8 crossed module corresponding to one of the normal series is isomorphic to some factor crossed modules in the  
 9 other normal series.

10 **Theorem 2.7 (Scherier Refinement Theorem for crossed modules)** Given two normal series

$$\begin{array}{ccccccc}
 1 = S_n & \longrightarrow & S_{n-1} & \longrightarrow & \cdots & \longrightarrow & S_1 & \longrightarrow & S_0 = T \\
 \downarrow & & \downarrow & & & & \downarrow & & \downarrow \partial \\
 1 = H_n & \longrightarrow & H_{n-1} & \longrightarrow & \cdots & \longrightarrow & H_1 & \longrightarrow & H_0 = G
 \end{array}$$

11 and

$$\begin{array}{ccccccc}
 1 = R_m & \longrightarrow & R_{m-1} & \longrightarrow & \cdots & \longrightarrow & R_1 & \longrightarrow & R_0 = T \\
 \downarrow & & \downarrow & & & & \downarrow & & \downarrow \partial \\
 1 = K_m & \longrightarrow & K_{m-1} & \longrightarrow & \cdots & \longrightarrow & K_1 & \longrightarrow & K_0 = G
 \end{array}$$

12 of a crossed module  $(T, G, \partial)$ . Then these normal series have equivalent refinements.

**Proof** We define  $T_{ij} = S_{i+1}(S_i \cap R_j)$ ,  $G_{ij} = H_{i+1}(H_i \cap K_j)$  for all  $0 \leq i \leq n-1$  and  $0 \leq j \leq m$ . Then  $(T_{ij}, G_{ij}, \partial_{ij})$  are subcrossed module of  $(T, G, \partial)$ , since  $(S_{i+1}, H_{i+1}, \partial_{i+1})$  is normal in  $(S_i, H_i, \partial_i)$  where  $\partial_{ij}$  is the restriction of  $\partial$ . Note that,

$$T_{i0} = S_{i+1}(S_i \cap R_0) = S_{i+1}(S_i \cap T) = S_{i+1}S_i = S_i$$

$$T_{im} = S_{i+1}(S_i \cap R_m) = S_{i+1}(S_i \cap 1) = S_{i+1}1 = S_{i+1}$$

and  $G_{i0} = H_i$ ,  $G_{im} = H_{i+1}$ , similarly. Thus we obtain a refinement

$$\begin{array}{cccccccccccccccc} 1 = S_n = T_{(n-1)m} & \twoheadrightarrow & \cdots & \twoheadrightarrow & T_{(n-1)0} = S_{n-1} = T_{(n-2)m} & \twoheadrightarrow & \cdots & \twoheadrightarrow & T_{10} = S_1 = T_{0m} & \twoheadrightarrow & \cdots & \twoheadrightarrow & T_{00} = S_0 = T \\ \downarrow & & & & \downarrow & & & & \downarrow & & & & \downarrow \partial \\ 1 = H_n = G_{(n-1)m} & \twoheadrightarrow & \cdots & \twoheadrightarrow & G_{(n-1)0} = H_{n-1} = G_{(n-2)m} & \twoheadrightarrow & \cdots & \twoheadrightarrow & G_{10} = H_1 = G_{0m} & \twoheadrightarrow & \cdots & \twoheadrightarrow & G_{00} = H_0 = G \end{array}$$

of the first normal series by the second. Similarly we obtain a refinement of the second normal series by constructing the crossed module  $T'_{ji} = R_{j+1}(R_j \cap S_i)$ ,  $G'_{ji} = K_{j+1}(K_j \cap H_i)$ . By Lemma 2.1, we have isomorphisms

$$\frac{(S_{i+1}, H_{i+1}, \partial_{i+1})((S_i, H_i, \partial_i) \cap (R_j, K_j, \partial_j))}{(S_{i+1}, H_{i+1}, \partial_{i+1})((S_i, H_i, \partial_i) \cap (R_{j+1}, K_{j+1}, \partial_{j+1}))} \cong \frac{(R_{j+1}, K_{j+1}, \partial_{j+1})((R_j, K_j, \partial_j) \cap (S_i, H_i, \partial_i))}{(R_{j+1}, K_{j+1}, \partial_{j+1})((R_j, K_j, \partial_j) \cap (S_{i+1}, H_{i+1}, \partial_{i+1}))}$$

1 as desired. □

2 **Theorem 2.8 (Jordan-Hölder Theorem for crossed modules)** *All composition series of a crossed mod-*  
3 *ule are equivalent to each other and are of the same minimal lenght.*

4 **Proof** This is direct consequence of Scherier Refinement Theorem for crossed modules and the definition of a  
5 composition series. □

6 **Example 2.9** *Let  $G$  be an abelian group given with a composition series of minimal lenght  $n$ , that is*

$$1 = G_n \longrightarrow G_{n-1} \longrightarrow \cdots \longrightarrow G_1 \longrightarrow G_0 = G.$$

7 *Then it is routine to check that the crossed module  $(1, G, inc)$  also has a composition series of minimal lenght*  
8  *$n$ :*

$$\begin{array}{cccccccc} 1 = 1 & \longrightarrow & 1 & \longrightarrow & \cdots & \longrightarrow & 1 & \longrightarrow & 1 = 1 \\ \downarrow & & \downarrow & & & & \downarrow & & \downarrow inc \\ 1 = G_n & \longrightarrow & G_{n-1} & \longrightarrow & \cdots & \longrightarrow & G_1 & \longrightarrow & G_0 = G \end{array}$$

9 *However, the minimal lenght of a composite series of the abelian crossed module  $(G, G, id)$  is  $2n$ , since*

$$\begin{array}{cccccccc} 1 = G_n & \longrightarrow & G_{n-1} & \longrightarrow & G_{n-1} & \longrightarrow & \cdots & \longrightarrow & G_1 & \longrightarrow & G_0 & \longrightarrow & G_0 = G \\ \downarrow & & \downarrow & & \downarrow & & & & \downarrow & & \downarrow & & \downarrow id \\ 1 = G_n & \longrightarrow & G_n & \longrightarrow & G_{n-1} & \longrightarrow & \cdots & \longrightarrow & G_1 & \longrightarrow & G_1 & \longrightarrow & G_0 = G \end{array}$$

10 *is a composition series of  $(G, G, id)$ . Note that this is not always true for a nonabelian group  $G$ , since  $(N, G, inc)$*   
11 *is not have to be a normal subcrossed module of  $(G, G, id)$  where  $N \trianglelefteq G$ , if  $[G, G]$  is not contained in  $N$ .*

### 3. The Case of 2–Crossed Modules

**Definition 3.1** A 2–chain complex of groups consists of groups  $S$ ,  $L$ ,  $M$  and group homomorphisms  $\partial_1 : L \rightarrow M$ ,  $\partial_2 : S \rightarrow L$

$$S \xrightarrow{\partial_2} L \xrightarrow{\partial_1} M$$

such that  $\text{im } \partial_2 \subseteq \ker \partial_1$ . If in addition  $\text{im } \partial_2 \trianglelefteq L$  and  $\text{im } \partial_1 \trianglelefteq M$ , then it is called a normal 2–complex of groups.

**Definition 3.2** A normal 2–complex of groups

$$S \xrightarrow{\partial_2} L \xrightarrow{\partial_1} M$$

with a mapping

$$\{ , \} : L \times L \longrightarrow S$$

and action of  $M$  on  $S$ ,  $L$  and  $M$ , where the action of  $M$  on itself is given by conjugation, is called a 2–crossed module, if the following conditions are hold, where  $m \triangleright l$  and  $m \triangleright s$  denotes action of  $m \in M$  on  $l \in L$  and  $s \in S$ , respectively and  $l \triangleright s$  is defined to be equal to  $\{\partial_2 s, l\} s$ .

1.  $\partial_1$  and  $\partial_2$  are  $M$ –equivariant, that is

$$i) \quad \partial_1(m \triangleright l) = m \triangleright \partial_1(l)$$

$$ii) \quad \partial_2(m \triangleright s) = m \triangleright \partial_2(s)$$

$$2. \quad \partial_2\{l_1, l_2\} = (\partial_1 l_1 \triangleright l_2)(l_1 l_2^{-1} l_1^{-1})$$

$$3. \quad \{\partial_2 s_1, \partial_2 s_2\} = [s_2, s_1]$$

$$4. \quad i) \quad \{l_1 l_2, l\} = (\partial_1 l_1 \triangleright \{l_2, l\})\{l_1, l_2 l_2^{-1}\}$$

$$ii) \quad \{l, l_1 l_2\} = \{l, l_1\}((l l_1 l^{-1}) \triangleright \{l, l_2\})$$

$$5. \quad \{l, \partial_2 s\} = (\partial_1 l \triangleright s)(l \triangleright s^{-1})$$

$$6. \quad m \triangleright \{l_1, l_2\} = \{m \triangleright l_1, m \triangleright l_2\}$$

for all  $s, s_1, s_2 \in S$ ,  $l, l_1, l_2 \in L$  and  $m \in M$ .

¶A 2–crossed module

$$S \xrightarrow{\partial_2} L \xrightarrow{\partial_1} M$$

is sometimes denoted as  $(S, L, M, \partial_2, \partial_1)$  or  $(S, L, M)$  shortly, when no risk of confusion is present.

Here the curly bracket  $\{ , \}$  is named Peiffer Lifting.

As shown in [3], the notation  $l \triangleright s$  introduced in the definition gives an action of  $L$  on  $S$ , keeping in the mind that  $\{\partial_2 s, l\}^{-1} = \partial_2 s \triangleright \{\partial_2 s^{-1}, l\}$ . On the other hand  $L$  also acts on  $S$  by  $\partial_1$ . Note that general

$$l \triangleright s \neq \partial_1 l \triangleright s$$

where action on the left hand side is the action given by  $\{\partial_2 s, l\} s$  and the one on the right hand side is derived from the action of  $M$  on  $S$  via  $\partial_1$ . While we use the same notation for both actions, this will not cause confusion, since the context clarifies it.

It easy to verify that the condition (5) can be given equivalently

$$(5') \{l, \partial_2 s\} \{\partial_2 s, l\} = (\partial_1 l \triangleright s) s^{-1}$$

by using the definition of the action of  $L$  on  $S$ .

Given a 2-crossed module

$$S \xrightarrow{\partial_2} L \xrightarrow{\partial_1} M$$

we have a crossed module

$$S \xrightarrow{\partial_2} L$$

with the action  $l \triangleright s = \{\partial_2 s, l\} s$  and two pre-crossed modules

$$L \xrightarrow{\partial_1} M$$

and

$$S \xrightarrow{\partial_1 \partial_2} M$$

where the latter has trivial boundary map  $\partial_1 \partial_2 = 1_M$ .

**Definition 3.3** *Given two 2-crossed modules*

$$S \xrightarrow{\partial_2} L \xrightarrow{\partial_1} M$$

and

$$T \xrightarrow{\partial'_2} K \xrightarrow{\partial'_1} N.$$

A homomorphism

$$\Phi : (S, L, M, \partial_2, \partial_1) \longrightarrow (T, K, N, \partial'_2, \partial'_1)$$

is a triple  $\Phi = (\alpha, \beta, \gamma)$  of group homomorphisms

$$\alpha : S \longrightarrow T$$

$$\beta : L \longrightarrow K$$

$$\gamma : M \longrightarrow N$$

1 making the diagram

$$\begin{array}{ccccc} S & \xrightarrow{\partial_2} & L & \xrightarrow{\partial_1} & M \\ \alpha \downarrow & & \downarrow \beta & & \downarrow \gamma \\ T & \xrightarrow{\partial'_2} & K & \xrightarrow{\partial'_1} & N \end{array}$$

commuting ( $\beta\partial_2 = \partial'_2\alpha$ ,  $\gamma\partial_1 = \partial'_1\beta$ ) and having the properties

$$\alpha(m \triangleright s) = \gamma(m) \triangleright \alpha(s)$$

$$\beta(m \triangleright l) = \gamma(m) \triangleright \beta(l)$$

$$\alpha\{l, l'\} = \{\beta(l), \beta(l')\}.$$

2 If each of  $\alpha$ ,  $\beta$  and  $\gamma$  is an isomorphism of groups, then  $\Phi$  is said to be a 2-crossed module isomorphism.

¶A homomorphism

$$\Phi : (S, L, M, \partial_2, \partial_1) \longrightarrow (T, K, N, \partial'_2, \partial'_1)$$

also preserves both actions of  $L$  on  $S$  since,

$$\alpha(l \triangleright s) = \alpha(\{\partial_2 s, l\}s) = \{\beta\partial_2 s, \beta l\}\alpha s = \{\partial'_2 \alpha s, \beta l\}\alpha s = \beta(l) \triangleright \alpha(s)$$

and

$$\alpha(\partial_1 l \triangleright s) = \gamma\partial_1 l \triangleright \alpha s = \partial'_1 \beta(l) \triangleright \alpha(s).$$

Also if  $\Phi = (\alpha, \beta, \gamma)$  is a 2-crossed module homomorphism, then

$$(\alpha, \beta) : (S, L, \partial_2) \longrightarrow (T, K, \partial'_2)$$

3 is a crossed module homomorphism.

4 **Definition 3.4** Let

$$S \xrightarrow{\partial_2} L \xrightarrow{\partial_1} M$$

5 be a 2-crossed module. If also

$$T \xrightarrow{\partial'_2} K \xrightarrow{\partial'_1} N$$

6 is a 2-crossed module with  $T \subseteq S$ ,  $K \subseteq L$ ,  $N \subseteq M$  with  $\partial'_1 = \partial_1|_K$ ,  $\partial'_2 = \partial_2|_T$  and restricted actions  
7 and Peiffer Lifting, then  $(T, K, N, \partial'_2, \partial'_1)$  is said to be a 2-subcrossed module of  $(S, L, M, \partial_2, \partial_1)$ . This case is  
8 denoted as  $(T, K, N) \leq (S, L, M)$ .

1 **Definition 3.5** [20] Given a 2-subcrossed module  $(T, K, N)$  of a 2-crossed module

$$S \xrightarrow{\partial_2} L \xrightarrow{\partial_1} M.$$

2 If

- 3 1.  $K \trianglelefteq L$  and  $N \trianglelefteq M$
- 4 2.  $m \triangleright k \in K$ ,  $m \triangleright t \in T$
- 5 3.  $(n \triangleright l)l^{-1} \in K$ ,  $(n \triangleright s)s^{-1} \in T$
- 6 4.  $\{k, l\}, \{l, k\} \in T$

7 for all  $s \in S$ ,  $l \in L$ ,  $m \in M$ ,  $t \in T$ ,  $k \in K$  and  $n \in N$ , then  $(T, K, N)$  is said to be a normal 2-subcrossed  
8 module of  $(S, L, M)$  and this is denoted by  $(T, K, N) \trianglelefteq (S, L, M)$ .

Note that from the condition (4) we also have  $l \triangleright t \in T$  and  $(k \triangleright s)s^{-1} \in T$  for  $l \in L$ ,  $t \in T$ ,  $k \in K$  and  $s \in S$ . Indeed, if  $(T, K, N)$  is normal 2-subcrossed module of  $(S, L, M, \partial_2, \partial_1)$  then for all  $l \in L$ ,  $t \in T$ ,  $k \in K$  and  $s \in S$  we have

$$l \triangleright t = \{\partial t, l\}t \in T,$$

$$(k \triangleright s)s^{-1} = \{\partial s, k\}ss^{-1} = \{\partial s, k\} \in T$$

and also

$$\partial_1 l \triangleright t \in T, (\partial_1 k \triangleright s)s^{-1} \in T.$$

In addition,  $T \trianglelefteq S$  since

$$sts^{-1} = \partial_2(s) \triangleright t \in T.$$

9 As a consequence of these arguments,  $(T, K)$  is a normal subcrossed module of  $(S, L)$ , if  $(T, K, N)$  is normal  
10 2-subcrossed module of  $(S, L, M, \partial_2, \partial_1)$ .

11 Note that the definition of normality given here is somewhat different than the definition given in [20].  
12 In addition to the definition given there, we add conditions on Peiffer Lifting of a quotient 2-crossed module.  
13 See Lemma 3.9 and Example 3.11.  
14

**Definition 3.6** [20] Given a 2-crossed module homomorphism

$$\Phi = (\alpha, \beta, \gamma) : (S, L, M, \partial_2, \partial_1) \longrightarrow (T, K, N, \partial'_2, \partial'_1).$$

15 The kernel of  $\Phi$  is defined by  $\ker \Phi = (\ker \alpha, \ker \beta, \ker \gamma)$  and the image of  $\Phi$  is defined by  $\text{im } \Phi = (\text{im } \alpha, \text{im } \beta, \text{im } \gamma)$ .

It is known that  $\ker \Phi$  is normal 2-subcrossed module of  $(S, L, M)$  in the sense of the definition given in [20], and also  $\text{im } \Phi$  is a 2-subcrossed module of  $(T, K, N)$ . Normality of  $\ker \Phi$  in the sense of the definition given here is seen showing in addition that

$$\{k, l\}, \{l, k\} \in \ker \alpha$$

16 for all  $k \in \ker \beta$  and  $l \in L$ . Indeed,

$$\alpha\{k, l\} = \{\beta(k), \beta(l)\} = \{1, \beta(l)\}$$

$$= \{\partial_2(1), \beta(l)\}1 = l \triangleright 1 = 1,$$

1

$$\begin{aligned} \alpha\{l, k\} &= \{\beta(l), \beta(k)\} = \{\beta(l), 1\}\{\beta(l), \partial_2(1)\} \\ &= (\partial'_1\beta(l) \triangleright 1)(\beta(l) \triangleright 1^{-1}) = 1 \cdot 1 = 1 \end{aligned}$$

2 gives  $\{k, l\}, \{l, k\} \in \ker \alpha$ .

3 **Definition 3.7** [20] Let  $(S, L, M, \partial_2, \partial_1)$  be a 2-crossed module and  $(T, K, N)$  be a normal 2-subcrossed  
4 module of  $(S, L, M)$ . Then the quotient 2-crossed module  $\frac{(S, L, M)}{(T, K, N)}$  is given by the quintuple  $(\frac{S}{T}, \frac{L}{K}, \frac{M}{N}, \overline{\partial_2}, \overline{\partial_1})$   
5 where  $\overline{\partial_2}, \overline{\partial_1}$  and Peiffer Lifting are given with

$$\overline{\partial_2}(sT) = \partial_2(s)K$$

$$\overline{\partial_1}(lK) = \partial_1(l)N$$

$$\{l_1K, l_2K\} = \{l_1, l_2\}T,$$

6 and the actions of  $\frac{M}{N}$  on  $\frac{S}{T}$  and  $\frac{L}{K}$  are given with

$$mN \triangleright sT = (m \triangleright s)T$$

$$mN \triangleright lK = (m \triangleright l)K.$$

7 **Proposition 3.8** Let  $(S, L, M, \partial_2, \partial_1)$  be a 2-crossed module and  $(T, K, N)$  be a normal 2-subcrossed module  
8 of  $(S, L, M)$ . Then, the triple  $\mathbf{Q} = (\mathbf{q}_1, \mathbf{q}_2, \mathbf{q}_3)$  given by  $\mathbf{q}_1(s) = sT$ ,  $\mathbf{q}_2(l) = lK$ ,  $\mathbf{q}_3(m) = mN$  for all  $s \in S$ ,  
9  $l \in L$  and  $m \in M$  gives a 2-crossed module homomorphism from  $(S, L, M, \partial_2, \partial_1)$  to  $(\frac{S}{T}, \frac{L}{K}, \frac{M}{N}, \overline{\partial_2}, \overline{\partial_1})$  and  
10 kernel of  $\mathbf{Q}$  is equal to  $(T, K, N)$ .

11 **Proof** It is known that  $\mathbf{q}_1, \mathbf{q}_2$  and  $\mathbf{q}_3$  are group homomorphisms, namely canonical mappings. The other  
12 conditions are easily seen from the following equations, for all  $s \in S$ ,  $l \in L$ ,  $m \in M$ .

$$\overline{\partial_2}\mathbf{q}_1(s) = \overline{\partial_2}(sT) = \partial_2(s)K = \mathbf{q}_2\partial_2(s)$$

$$\overline{\partial_1}\mathbf{q}_2(l) = \overline{\partial_1}(lK) = \partial_1(l)N = \mathbf{q}_3\partial_1(l)$$

$$\{\mathbf{q}_2(l_1), \mathbf{q}_2(l_2)\} = \{l_1K, l_2K\} = \{l_1, l_2\}T = \mathbf{q}_1(\{l_1, l_2\})$$

$$\mathbf{q}_3(m) \triangleright \mathbf{q}_1(s) = mN \triangleright sT = (m \triangleright s)T = \mathbf{q}_1(m \triangleright s)$$

$$\mathbf{q}_3(m) \triangleright \mathbf{q}_2(l) = mN \triangleright lK = (m \triangleright l)K = \mathbf{q}_2(m \triangleright l).$$

13 As a result of group theoretical identities we also have

$$\ker \mathbf{Q} = (\ker \mathbf{q}_1, \ker \mathbf{q}_2, \ker \mathbf{q}_3) = (T, K, N).$$

14

□

15 The additional requirements on normality that  $\{k, l\}, \{l, k\} \in T$  for  $k \in K$ ,  $l \in L$  are needed to ensure the  
16 well-definedness of quotient Peiffer Lifting. In fact, these are necessary and sufficient conditions for the Peiffer  
17 Lifting to be well-defined as seen in the following Lemma.

**Lemma 3.9** Given a 2-crossed module  $(S, L, M, \partial_2, \partial_1)$  and a 2-subcrossed module  $(T, K, N)$  of  $(S, L, M)$ , which satisfies all conditions of normality definition in [20]. Then the Peiffer Lifting of the quotient is well-defined iff and only iff  $\{k, l\}, \{l, k\} \in T$  for all  $k \in K, l \in L$ .

**Proof** Let  $\{k, l\}, \{l, k\} \in T$  for all  $k \in K$  and  $l \in L$ . Assume that  $l_1 K = l_2 K$  in  $L/K$ . Then  $l_1 l_2^{-1} \in K$ . Observing that

$$\begin{aligned} \{l_2, l\} \{l_1, l\}^{-1} &= \{l_2, l\} \{l_1 l_2^{-1} l_2, l\}^{-1} \\ &= \{l_2, l\} \left[ (\partial_1(l_1 l_2^{-1}) \triangleright \{l_2, l\}) \{l_1 l_2^{-1}, l_2 l_2^{-1}\} \right]^{-1} \\ &= \{l_2, l\} \{l_1 l_2^{-1}, l_2 l_2^{-1}\}^{-1} (\partial_1(l_1 l_2^{-1}) \triangleright \{l_2, l\}^{-1}) \\ &= \{l_2, l\} \{l_1 l_2^{-1}, l_2 l_2^{-1}\}^{-1} \left( (\partial_1(l_1 l_2^{-1}) \triangleright \{l_2, l\}^{-1}) \{l_2, l\} \right) \{l_2, l\} \end{aligned}$$

we see  $\{l_2, l\} \{l_1, l\}^{-1} \in T$ , hence  $\{l_1 K, l K\} = \{l_2 K, l K\}$  and from

$$\begin{aligned} \{l, l_1\}^{-1} \{l, l_2\} &= ((l_1 l^{-1}) \triangleright \{l, l_1^{-1}\}) \{l, l_2\} \\ &= (l_1 l^{-1}) \triangleright \left( \{l, l_1^{-1}\} ((l_1^{-1} l^{-1}) \triangleright \{l, l_2\}) \right) \\ &= (l_1 l^{-1}) \triangleright \{l, l_1^{-1} l_2\} \end{aligned}$$

we have  $\{l, l_1\}^{-1} \{l, l_2\} \in T$  so that  $\{l K, l_1 K\} = \{l K, l_2 K\}$ . For the converse, we have  $\{l_1, l\} \{l_2, l\}^{-1}, \{l, l_1\} \{l, l_2\}^{-1} \in T$  if  $l_1 l_2^{-1} \in K$ . Then for all  $k \in K$  and  $l \in L$ , since  $k = k l^{-1} \in K$ ,

$$\begin{aligned} \{k, l\} &= \{k, l\} 1 = \{k, l\} \{1, l\}^{-1} \in T, \\ \{l, k\} &= \{l, k\} 1 = \{l, k\} \{l, 1\}^{-1} \in T. \end{aligned}$$

□

As a result of Proposition 3.8, Lemma 3.9 and the discussion following Definition 3.6, the following corollary immediately seen, which further emphasize that the given definition of normality is categorically correct.

**Corollary 3.10** Let  $(S, L, M, \partial_2, \partial_1)$  be a 2-crossed module and  $(T, K, N)$  be a 2-subcrossed module of  $(S, L, M)$ . The necessary and sufficient condition for  $(T, K, N)$  to be normal, is that there exists a 2-crossed module homomorphism  $\Phi : (S, L, M, \partial_2, \partial_1) \rightarrow (S', L', M', \partial'_2, \partial'_1)$ , where  $(S', L', M', \partial'_2, \partial'_1)$  is an any 2-crossed module, such that  $\ker \Phi = (T, K, N)$ .

The following example illustrates that there are 2-subcrossed modules, which are not normal but satisfy all conditions of normality except the condition on the Peiffer Lifting. Therefore, the condition on the Peiffer Lifting is essential for an accurate definition for normality on 2-subcrossed modules.

**Example 3.11** Consider the multiplicative groups  $T = M = N = \{1\}$ ,  $S = \{-1, 1\}$ ,  $K = \{-1, 1, i, -i\}$  and  $L = \{1, i, j, k, -1, -i, -j, -k\}$ , as subgroups of multiplicative group of nonzero quaternions. Then  $S \hookrightarrow L \rightarrow M$  and  $T \hookrightarrow K \rightarrow N$  are 2-crossed modules with the homomorphisms are given as inclusions and trivial homomorphisms, all actions of  $M = N$  are trivial, and both Peiffer Liftings are inverse commutator, that is

$$\{l_1, l_2\} = l_2 l_1 l_2^{-1} l_1^{-1}$$

$$\{k_1, k_2\} = k_2 k_1 k_2^{-1} k_1^{-1}.$$

- 1 Thus  $l \triangleright s = \{\partial s, l\}s = l \partial s l^{-1} \partial s^{-1} s = l s l^{-1} s^{-1} s = l s l^{-1}$  and similarly  $k \triangleright t = k t k^{-1}$  are actions by conjugation.  
2 It is routine to check that  $(S, L, M)$  and  $(T, K, N)$  are 2-crossed module and  $(T, K, N)$  is a 2-subcrossed  
3 module of  $(S, L, M)$ .  
4  $(T, K, N)$  satisfies the first three conditions in Definition 3.5. In fact,  
5 1.  $N \trianglelefteq M$  trivially, and observing that  $jij^{-1} = -i$ ,  $kik^{-1} = -i$  in particular, guides the normality of  $K$   
6 in  $L$ .  
7 2.  $1 \triangleright z \in K$ ,  $1 \triangleright 1 \in T$  and  $l \triangleright 1 \in T$  for  $1 \in M$ ,  $z \in K$ ,  $1 \in T$ ,  $l \in L$ .  
8 3.  $(1 \triangleright l)l^{-1} = 1 \in K$ ,  $(1 \triangleright s)s^{-1} = 1 \in T$ ,  $(z \triangleright 1)1^{-1} = z1z^{-1}1 = 1 \in T$  and  $(z \triangleright (-1))(-1)^{-1} =$   
9  $z(-1)z^{-1}(-1) = 1 \in T$  for  $1 \in N$ ,  $l \in L$ ,  $s \in S$ ,  $z \in K$  and  $1, -1 \in S$ .

However, note that the condition (4) is not satisfied since

$$\{i, j\} = jij^{-1}i^{-1} = ji(-j)(-i) = jiji = (-k)(-k) = k^2 = -1 \notin T.$$

In this circumstance, if we try to form the quotient  $(S/T, L/K, M/N)$ , then

$$jK = \{j1, ji, j(-1), j(-i)\} = \{j, -k, -j, k\}$$

$$kK = \{k1, ki, k(-1), k(-i)\} = \{k, j, -k, -j\}$$

hence  $jK = kK$ . But

$$\{jK, jK\} = \{j, j\}T = jjj^{-1}j^{-1}T = 1T = T = \{1\}$$

$$\begin{aligned} \{jK, kK\} &= \{j, k\}T = kjk^{-1}j^{-1}T = kj(-k)(-j)T \\ &= kjkjT = (-i)(-i)T = (-1)T = \{-1\} \end{aligned}$$

- 11 so that  $\{jK, jK\} \neq \{jK, kK\}$ , which is a matter of being not well-defined.

**Theorem 3.12** [20] Given a 2-crossed module homomorphism

$$\Phi = (\alpha, \beta, \gamma) : (S, L, M, \partial_2, \partial_1) \longrightarrow (T, K, N, \partial'_2, \partial'_1).$$

Then

$$\frac{(S, L, M)}{\ker \Phi} \cong \text{im } \Phi.$$

- 12 The above theorem is the First Isomorphism Theorem for 2-crossed modules, which have been readily stated  
13 and proven in [20] and its proof is valid also under the definitions given here. Now, we give some preliminary  
14 definitions and propositions as a preparation of statements of Second and Third Isomorphism Theorems for  
15 2-crossed modules and their some consequences.

- 16 **Definition 3.13** [20] Let  $(T_1, K_1, N_1)$  and  $(T_2, K_2, N_2)$  be two 2-subcrossed modules of  $(S, L, M, \partial_2, \partial_1)$ .  
17 Then the intersection of  $(T_1, K_1, N_1)$  and  $(T_2, K_2, N_2)$  is given by  $(T_1, K_1, N_1) \cap (T_2, K_2, N_2) = (T_1 \cap T_2, K_1 \cap$   
18  $K_2, N_1 \cap N_2)$ .

Note that the additional condition on Peiffer Lifting for normality follows immediately, which makes the following result valid also under the definitions given here.

**Proposition 3.14** [20] *Intersection of 2-subcrossed modules is also a 2-subcrossed module. In addition, if each 2-subcrossed module is normal, then the intersection is also normal.*

**Definition 3.15** *Let  $(S, L, M, \partial_2, \partial_1)$  be a 2-crossed module and  $(T, K, N), (V, J, P)$  be 2-subcrossed modules of  $(S, L, M)$ . Then the multiplication of  $(T, K, N)$  and  $(V, J, P)$  is given by  $(T, K, N)(V, J, P) := (TV, KJ, NP)$ .*

**Proposition 3.16**  *$(T, K, N)$  be a normal 2-subcrossed module and  $(V, J, P)$  be a 2-subcrossed module of  $(S, L, M, \partial_2, \partial_1)$ .*

(i)  $(T, K, N)(V, J, P)$  is a 2-subcrossed module of  $(S, L, M)$

(ii) If in addition  $(V, J, P)$  is normal, then also  $(T, K, N)(V, J, P)$  is a normal 2-subcrossed module of  $(S, L, M)$ .

**Proof**

(i)  $\partial_2(TV) \subseteq KJ$  and  $\partial_1(KJ) \subseteq NP$  is clear. For  $t \in T, k \in K, n \in N, v \in V, j \in J$  and  $p \in P$  we have

$$\begin{aligned} (np) \triangleright (tv) &= ((np) \triangleright t)((np) \triangleright v)(p \triangleright v)^{-1}(p \triangleright v) \\ &= ((np) \triangleright t)(n \triangleright (p \triangleright v))(p \triangleright v)^{-1}(p \triangleright v) \in TV \end{aligned}$$

and

$$\begin{aligned} (np) \triangleright (kj) &= ((np) \triangleright k)((np) \triangleright j)(p \triangleright j)^{-1}(p \triangleright j) \\ &= ((np) \triangleright k)(n \triangleright (p \triangleright j))(p \triangleright j)^{-1}(p \triangleright j) \in KJ. \end{aligned}$$

Also for  $k_1, k_2 \in K, j_1, j_2 \in J$

$$\begin{aligned} \{k_1 j_1, k_2 j_2\} &= (\partial_1 k_1 \triangleright \{j_1, k_2 j_2\})\{k_1, j_1 k_2 j_2 j_1^{-1}\} \\ &= \left( n \triangleright \left( \{j_1, k_2\}((j_1 k_2 j_1^{-1}) \triangleright \{j_1, j_2\}) \right) \right) t \\ &= (n \triangleright t')(n \triangleright (k' \triangleright v))t \\ &= t''(n \triangleright l)l^{-1}(k' \triangleright v)v^{-1}vt \\ &= t''t'''t''''vt \\ &= t''t'''t''''t'''''v \in TV \end{aligned}$$

where  $n = \partial_1 k, t = \{k_1, j_1 k_2 j_2 j_1^{-1}\}, t' = \{j_1, k_2\}, k' = j_1 k_2 j_1^{-1}, t'' = n \triangleright t', l = k' \triangleright v, t''' = (n \triangleright l)l^{-1}, t'''' = (k' \triangleright v)v^{-1}, t''''' = vt v^{-1}$ .

(ii) (1)  $KJ \trianglelefteq L$  and  $NP \trianglelefteq M$  follows from  $K, J \trianglelefteq L$  and  $N, P \trianglelefteq M$ .

(2)  $m \triangleright (kj) = (m \triangleright k)(m \triangleright j) \in KJ$  and  $m \triangleright (tv) = (m \triangleright t)(m \triangleright v) \in TV$ .

$$(3) \quad (np) \triangleright l) l^{-1} = (n \triangleright (p \triangleright l))(p \triangleright l)^{-1} (p \triangleright l) l^{-1} \in KJ \text{ and} \\ (np) \triangleright s) s^{-1} = (n \triangleright (p \triangleright s))(p \triangleright s)^{-1} (p \triangleright s) s^{-1} \in TV.$$

$$(4) \quad \{kj, l\} = (\partial_1 k \triangleright \{j, l\}) \{k, jl j^{-1}\} \in VT = TV \\ \{l, kj\} = \{l, k\} ((lkl^{-1}) \triangleright \{l, j\}) \in TV.$$

□

**Definition 3.17** [20] *Direct product of two 2-crossed modules  $(S, L, M, \partial_2, \partial_1)$  and  $(S', L', M', \partial'_2, \partial'_1)$  is defined as the 2-crossed module*

$$(S, L, M, \partial_2, \partial_1) \times (S', L', M', \partial'_2, \partial'_1) = (S \times S', L \times L', M \times M', \partial_2 \times \partial'_2, \partial_1 \times \partial'_1)$$

where

$$(\partial_1 \times \partial'_1)(l, l') = (\partial_1 l, \partial'_1 l'), \\ (\partial_2 \times \partial'_2)(s, s') = (\partial_2 s, \partial'_2 s'), \\ (m, m') \triangleright (s, s') = (m \triangleright s, m' \triangleright s'), \\ (m, m') \triangleright (l, l') = (m \triangleright l, m' \triangleright l'), \\ \{(l_1, l'_1), (l_2, l'_2)\} = (\{l_1, l_2\}, \{l'_1, l'_2\}).$$

**Proposition 3.18** *Let  $(S, L, M, \partial_2, \partial_1)$  be a 2-crossed module and  $(T, K, N), (V, J, P)$  be normal 2-subcrossed modules of  $(S, L, M)$  such that*

$$i) \quad (T, K, N)(V, J, P) = (S, L, M),$$

$$ii) \quad (T, K, N) \cap (V, J, P) = (1, 1, 1).$$

Then,

$$(S, L, M) \cong (T, K, N) \times (V, J, P).$$

**Proof** Noting that  $T, V \trianglelefteq S$ ,  $K, J \trianglelefteq L$ ,  $N, P \trianglelefteq M$ ,  $TV = S$ ,  $KJ = L$ ,  $NP = M$ ,  $T \cap V = 1$ ,  $K \cap J = 1$  and  $N \cap P = 1$ , we can conclude from the group theory that

$$\alpha : T \times V \longrightarrow S, \quad \alpha(t, v) = tv \\ \beta : K \times J \longrightarrow L, \quad \beta(k, j) = kj \\ \gamma : N \times P \longrightarrow M, \quad \gamma(n, p) = np$$

are group isomorphisms. We see that

$$\Phi = (\alpha, \beta, \gamma) : (T, K, N) \times (V, J, P) \longrightarrow (S, L, M)$$

1 satisfies other remaining conditions for 2-crossed module homomorphisms.

$$\begin{array}{ccccc}
 T \times V & \xrightarrow{\partial_2 \times \partial_2} & K \times J & \xrightarrow{\partial_1 \times \partial_1} & N \times P \\
 \downarrow \alpha & & \downarrow \beta & & \downarrow \gamma \\
 S & \xrightarrow{\partial_2} & L & \xrightarrow{\partial_1} & M
 \end{array}$$

2

$$\begin{aligned}
 \beta(\partial_2 \times \partial_2)(t, v) &= \beta(\partial_2 t, \partial_2 v) = \partial_2 t \partial_2 v = \partial_2(tv) = \partial_2 \alpha(t, v), \\
 \gamma(\partial_1 \times \partial_1)(k, j) &= \gamma(\partial_1 k, \partial_1 j) = \partial_1 k \partial_1 j = \partial_1(kj) = \partial_1 \beta(k, j).
 \end{aligned}$$

3 Note that from the normality, for all  $p \in P$ ,  $t \in T$  and  $k \in K$ ,  $(p \triangleright t)t^{-1} \in V$  and also  $p \triangleright t \in T$  and  $t^{-1} \in T$   
4 gives  $(p \triangleright t)t^{-1} \in T \cap V = \{1\}$  which implies  $p \triangleright t = t$  and similarly  $p \triangleright k = k$ . So  $P$  acts on  $T$  and  $K$  trivially.  
5 On the other side  $N$  acts on  $V$  and  $J$  trivially. Hence,

$$\begin{aligned}
 \alpha((n, p) \triangleright (t, v)) &= \alpha(n \triangleright t, p \triangleright v) = (n \triangleright t)(p \triangleright v) = (n \triangleright (p \triangleright t))(n \triangleright (p \triangleright v)) \\
 &= (np \triangleright t)(np \triangleright v) = np \triangleright tv = \gamma(n, p) \triangleright \alpha(t, v).
 \end{aligned}$$

6

$$\begin{aligned}
 \beta((n, p) \triangleright (k, j)) &= \alpha(n \triangleright k, p \triangleright j) = (n \triangleright k)(p \triangleright j) = (n \triangleright (p \triangleright k))(n \triangleright (p \triangleright j)) \\
 &= (np \triangleright k)(np \triangleright j) = np \triangleright kj = \gamma(n, p) \triangleright \beta(k, j).
 \end{aligned}$$

Let  $k_1, k_2 \in K$  and  $j_1, j_2 \in J$ . Observe that

$$\{k_1 j_1, k_2 j_2\} = \{k_1 j_1, k_2\} \left( (k_1 j_1 k_2 (k_1 j_1)^{-1}) \triangleright \{k_1 j_1, j_2\} \right)$$

7 Now,  $K$  acts on  $V$  trivially since for  $k \in K$  and  $v \in V$ ,  $k \triangleright v = \{\partial_2 v, k\}v$  and  $k \in K$ ,  $\partial_2 v \in J$  together  
8 implies that  $\{\partial_2 v, k\} \in T \cap V = \{1\}$  so that  $k \triangleright v = v$ . Here we have  $(k_1 j_1)k_2(k_1 j_1)^{-1} \in K$  and by  $j_2 \in J$ ,  
9  $\{k_1 j_1, j_2\} \in V$ . Thus

$$\begin{aligned}
 \{k_1 j_1, k_2 j_2\} &= \{k_1 j_1, k_2\} \{k_1 j_1, j_2\} \\
 &= (\partial_1 k_1 \triangleright \{j_1, k_2\}) \{k_1, j_1 k_2 j_1^{-1}\} (\partial_1 k_1 \triangleright \{j_1, j_2\}) \{k_1, j_1 j_2 j_1^{-1}\}.
 \end{aligned}$$

Note that since  $j_1 \in J$  and  $k_2 \in K$ ,  $\{j_1, k_2\} \in V \cap T = \{1\}$  and similarly  $\{k_1, j_1 j_2 j_1^{-1}\} = 1$ . Also,  $\partial_1 k_1 \in N$  and  $\{j_1, j_2\} \in V$  gives  $\partial_1 k_1 \triangleright \{j_1, j_2\} = \{j_1, j_2\}$  from the triviality of the action. Lastly,  $j_1 k_2 j_1^{-1} k_2^{-1} \in J \cap K = \{1\}$  so that  $j_1 k_2 j_1^{-1} = k_2$ . So we have

$$\{k_1 j_1, k_2 j_2\} = \{k_1, k_2\} \{j_1, j_2\}$$

which means that

$$\{\beta(k_1, j_1), \beta(k_2, j_2)\} = \alpha(\{k_1, k_2\}, \{j_1, j_2\}).$$

10

□

**Definition 3.19** If a 2-crossed module  $(S, L, M, \partial_2, \partial_1)$  and its given two normal 2-subcrossed modules  $(T, K, N)$  and  $(V, J, P)$  satisfies the conditions

i)  $(T, K, N)(V, J, P) = (S, L, M),$

ii)  $(T, K, N) \cap (V, J, P) = (1, 1, 1),$

then  $(S, L, M)$  is said to be internal direct product of  $(T, K, N)$  and  $(V, J, P)$ .

**Theorem 3.20** Let  $(S, L, M, \partial_2, \partial_1)$  be a 2-crossed module,  $(T, K, N)$  be a normal 2-subcrossed module of  $(S, L, M)$  and  $(V, J, P)$  be a 2-subcrossed module of  $(S, L, M)$ . Then,

$$\frac{(V, J, P)}{(T, K, N) \cap (V, J, P)} \cong \frac{(T, K, N)(V, J, P)}{(T, K, N)}.$$

**Proof** We shall show that

$$\left( \frac{V}{T \cap V}, \frac{J}{K \cap J}, \frac{P}{N \cap P} \right) \cong \left( \frac{TV}{T}, \frac{KJ}{K}, \frac{NP}{N} \right).$$

Consider the functions,

$$\alpha : \frac{V}{T \cap V} \longrightarrow \frac{TV}{T}, \quad \alpha(v(T \cap V)) = vT$$

$$\beta : \frac{J}{K \cap J} \longrightarrow \frac{KJ}{K}, \quad \beta(j(K \cap J)) = jK$$

$$\gamma : \frac{P}{N \cap P} \longrightarrow \frac{NP}{N}, \quad \gamma(p(N \cap P)) = pN.$$

It is well-known that  $\alpha, \beta$  and  $\gamma$  are group isomorphisms. We want to show that  $\Phi = (\alpha, \beta, \gamma)$  is a 2-crossed module homomorphism.

Consider first the diagram

$$\begin{array}{ccccc} \frac{V}{T \cap V} & \xrightarrow{\quad \overline{\partial_2} \quad} & \frac{J}{K \cap J} & \xrightarrow{\quad \overline{\partial_1} \quad} & \frac{P}{N \cap P} \\ \downarrow \alpha & & \downarrow \beta & & \downarrow \gamma \\ \frac{TV}{T} & \xrightarrow{\quad \widetilde{\partial_2} \quad} & \frac{KJ}{K} & \xrightarrow{\quad \widetilde{\partial_1} \quad} & \frac{NP}{N} \end{array}$$

For all  $v \in V, k \in K$  and  $p \in P$

$$\beta \overline{\partial_2}(v(T \cap V)) = \beta(\partial_2 v(K \cap J)) = \partial_2 vK = \widetilde{\partial_2}(vT) = \widetilde{\partial_2} \alpha(v(T \cap V)).$$

and

$$\gamma \overline{\partial_1}(j(K \cap J)) = \gamma(\partial_1 j(N \cap P)) = \partial_1 jN = \widetilde{\partial_1}(jK) = \widetilde{\partial_1} \beta(j(K \cap J)).$$

1 In addition,

$$\begin{aligned}\alpha(p(N \cap P) \triangleright v(T \cap V)) &= \alpha((p \triangleright v)(T \cap V)) = (p \triangleright v)T \\ &= pN \triangleright vT = \gamma(p(N \cap P)) \triangleright \alpha(v(T \cap V))\end{aligned}$$

2 and

$$\begin{aligned}\beta(p(N \cap P) \triangleright j(K \cap J)) &= \beta((p \triangleright j)(K \cap J)) = (p \triangleright j)K \\ &= pN \triangleright jK = \gamma(p(N \cap P)) \triangleright \beta(j(K \cap J)).\end{aligned}$$

3 Lastly, for all  $j_1, j_2 \in J$ , we have

$$\begin{aligned}\alpha\{j_1(K \cap J), j_2(K \cap J)\} &= \alpha(\{j_1, j_2\}(T \cap V)) = \{j_1, j_2\}T \\ &= \{j_1K, j_2K\} = \left\{ \beta(j_1(K \cap J)), \beta(j_2(K \cap J)) \right\}.\end{aligned}$$

4

□

5 **Proposition 3.21** *Given a normal 2-subcrossed module  $(T, K, N)$  and a 2-subcrossed module  $(V, J, P)$  of*  
 6  *$(S, L, M, \partial_2, \partial_1)$  such that  $T \subseteq V$ ,  $K \subseteq J$  and  $N \subseteq P$ . Then,  $\frac{(V, J, P)}{(T, K, N)}$  is a normal 2-subcrossed module of*  
 7  *$\frac{(S, L, M)}{(T, K, N)}$  if and only if  $(V, J, P)$  is a normal 2-subcrossed module of  $(S, L, M)$ .*

8 **Proof** It is obvious that  $\frac{(V, J, P)}{(T, K, N)}$  is a 2-subcrossed module of  $\frac{(S, L, M)}{(T, K, N)}$ .

9 1. For groups it is known that  $J \trianglelefteq L \iff \frac{J}{K} \trianglelefteq \frac{L}{K}$  and  $P \trianglelefteq M \iff \frac{P}{N} \trianglelefteq \frac{M}{N}$ .

10 2.  $m \triangleright j \in J \iff (m \triangleright j)K \in \frac{J}{K} \iff mN \triangleright jK \in \frac{J}{K}$   
 11  $m \triangleright v \in V \iff (m \triangleright v)T \in \frac{V}{T} \iff mN \triangleright vT \in \frac{V}{T}.$

12 3.  $(p \triangleright l)l^{-1} \in J \iff (p \triangleright l)l^{-1}K \in \frac{J}{K} \iff (pN \triangleright lK)(lK)^{-1} \in \frac{J}{K}$   
 13  $(p \triangleright s)s^{-1} \in V \iff (p \triangleright s)s^{-1}T \in \frac{V}{T} \iff (pN \triangleright sT)(sT)^{-1} \in \frac{V}{T}.$

14 4.  $\{j, l\} \in V \iff \{j, l\}T \in \frac{V}{T} \iff \{jK, lK\} \in \frac{V}{T}$   
 15  $\{l, j\} \in V \iff \{l, j\}T \in \frac{V}{T} \iff \{lK, jK\} \in \frac{V}{T}.$

16

□

**Theorem 3.22** *Let  $(S, L, M, \partial_2, \partial_1)$  be a 2-crossed module,  $(T, K, N)$  and  $(V, J, P)$  be two normal 2-subcrossed module of  $(S, L, M)$ , such that  $T \subseteq V$ ,  $K \subseteq J$  and  $N \subseteq P$ . Then,*

$$\frac{\frac{(S, L, M)}{(T, K, N)}}{\frac{(V, J, P)}{(T, K, N)}} \cong \frac{(S, L, M)}{(V, J, P)}.$$

17 **Proof** Define  $\alpha$ ,  $\beta$  and  $\gamma$  as

$$\alpha: \frac{S/T}{V/T} \longrightarrow \frac{S}{V}, \quad \alpha(sT(V/T)) = sV$$

$$\beta : \frac{L/K}{J/K} \longrightarrow \frac{L}{J}, \quad \beta(lK(J/K)) = lJ$$

$$\gamma : \frac{M/N}{P/N} \longrightarrow \frac{M}{P}, \quad \gamma(mN(P/N)) = mP.$$

1 which are known to be group isomorphisms.

2 Consider the diagram:

$$\begin{array}{ccccc} \frac{S/T}{V/T} & \xrightarrow{\overline{\partial}_2} & \frac{L/K}{J/K} & \xrightarrow{\overline{\partial}_1} & \frac{M/N}{P/N} \\ \downarrow \alpha & & \downarrow \beta & & \downarrow \gamma \\ \frac{S}{V} & \xrightarrow{\overline{\partial}_2} & \frac{L}{J} & \xrightarrow{\overline{\partial}_1} & \frac{M}{P} \end{array}$$

3 For all  $s \in S$ ,  $l, l_1, l_2 \in L$  and  $m \in M$

$$\begin{aligned} \beta \overline{\partial}_2(sT(V/T)) &= \beta(\tilde{\partial}_2(sT)J/K) = \beta(\partial_2 sK(J/K)) \\ &= \partial_2 sJ = \overline{\partial}_2(sV) = \overline{\partial}_2 \alpha(sT(V/T)), \end{aligned}$$

4

$$\begin{aligned} \gamma \overline{\partial}_1(lK(J/K)) &= \gamma(\tilde{\partial}_1(lK)P/N) = \gamma(\partial_1 lN(P/N)) \\ &= \partial_1 lP = \overline{\partial}_1(lJ) = \overline{\partial}_1 \beta(lK(J/K)), \end{aligned}$$

5 and

$$\begin{aligned} \alpha(mN(P/N) \triangleright sT(V/T)) &= \alpha((mN \triangleright sT)V/T) = \alpha((m \triangleright s)T(V/T)) \\ &= (m \triangleright s)V = mP \triangleright sV \\ &= \gamma(mN(P/N)) \triangleright \alpha(sT(V/T)), \end{aligned}$$

6

$$\begin{aligned} \beta(mN(P/N) \triangleright lK(J/K)) &= \beta((mN \triangleright lK)J/K) = \beta((m \triangleright l)K(J/K)) \\ &= (m \triangleright l)J = mP \triangleright lJ \\ &= \gamma(mN(P/N)) \triangleright \beta(lK(J/K)). \end{aligned}$$

7 In addition,

$$\begin{aligned} \alpha\{l_1K(J/K), l_2K(J/K)\} &= \alpha(\{l_1K, l_2K\}(V/T)) = \alpha(\{l_1, l_2\}T(V/T)) \\ &= \{l_1, l_2\}V = \{l_1J, l_2J\} \\ &= \left\{ \beta(l_1K(J/K)), \beta(l_2K(J/K)) \right\}. \end{aligned}$$

8

□

**Lemma 3.23** Let  $(S, L, M, \partial_2, \partial_1)$  be a 2-crossed module,  $(T, K, N)$  and  $(V, J, P)$  be 2-subcrossed modules of  $(S, L, M)$ ,  $(Y, H, Q)$  be a normal 2-subcrossed module of  $(T, K, N)$  and  $(Z, G, R)$  be a normal 2-subcrossed module of  $(V, J, P)$ . Then

$$(Y, H, Q)((T, K, N) \cap (Z, G, R))$$

is a normal 2-subcrossed module of

$$(Y, H, Q)((T, K, N) \cap (V, J, P))$$

and

$$(Z, G, R)((Y, H, Q) \cap (V, J, P))$$


is a normal 2-subcrossed module of

$$(Z, G, R)((T, K, N) \cap (V, J, P))$$

and also,

$$\frac{(Y, H, Q)((T, K, N) \cap (V, J, P))}{(Y, H, Q)((T, K, N) \cap (Z, G, R))} \cong \frac{(Z, G, R)((T, K, N) \cap (V, J, P))}{(Z, G, R)((Y, H, Q) \cap (V, J, P))}.$$

- 1 Note that the elements of the lattice of 2-subcrossed modules of  $(S, L, M)$ , which also includes some 2-subcrossed
- 2 modules in this lemma, can be represented in the diagram below. This is why the group-theoretical analogue of
- 3 this lemma is sometimes called the Butterfly Lemma. For a more economical use of the spaces in the diagram,
- 4 we represent each 2-subcrossed module with calligraphic version of the first letter involved.



5

6 **Proof** We consider the 2-crossed module

$$\frac{(T, K, N) \cap (V, J, P)}{(Y, H, Q) \cap (V, J, P)((T, K, N) \cap (Z, G, R))} \cong \left( \frac{T \cap V}{(Y \cap V)(T \cap Z)}, \frac{K \cap J}{(H \cap J)(K \cap G)}, \frac{N \cap P}{(Q \cap P)(N \cap R)} \right).$$

1 From the group theory it is known that

$$\begin{aligned}\alpha &: \frac{Y(T \cap V)}{Y(T \cap Z)} \longrightarrow \frac{T \cap V}{(Y \cap V)(T \cap Z)}, \quad \alpha(ysY(T \cap Z)) = s(Y \cap V)(T \cap Z) \\ \beta &: \frac{H(K \cap J)}{H(K \cap G)} \longrightarrow \frac{K \cap J}{(H \cap J)(K \cap G)}, \quad \beta(hlH(K \cap G)) = l(H \cap J)(K \cap G) \\ \gamma &: \frac{Q(N \cap P)}{Q(N \cap R)} \longrightarrow \frac{N \cap P}{(Q \cap P)(N \cap R)}, \quad \gamma(qmQ(N \cap R)) = m(Q \cap P)(N \cap R).\end{aligned}$$

2 are group isomorphisms [21]. Then for  $ysY(T \cap Z) \in \frac{Y(T \cap V)}{Y(T \cap Z)}$

$$\begin{aligned}\beta \bar{\partial}_2(ysY(T \cap Z)) &= \beta(\partial_2(ys)H(K \cap G)) = \partial_2 s(H \cap J)(K \cap G) \\ &= \tilde{\partial}_2(s(Y \cap V)(T \cap Z)) = \tilde{\partial}_2 \alpha(ysY(T \cap Z))\end{aligned}$$

3 and for  $hlH(K \cap G) \in \frac{H(K \cap J)}{H(K \cap G)}$

$$\begin{aligned}\gamma \bar{\partial}_1(hlH(K \cap G)) &= \gamma(\partial_1(hl)Q(N \cap R)) = \partial_1 l(Q \cap P)(N \cap R) \\ &= \tilde{\partial}_1(l(H \cap J)(K \cap G)) = \tilde{\partial}_1 \beta(hlH(K \cap G)).\end{aligned}$$

4 For  $ysY(T \cap Z) \in \frac{Y(T \cap V)}{Y(T \cap Z)}$ ,  $qmQ(N \cap R) \in \frac{Q(N \cap P)}{Q(N \cap R)}$

$$\begin{aligned}\alpha(qmQ(N \cap R) \triangleright ysY(T \cap Z)) &= \alpha((qm \triangleright ys)Y(T \cap Z)) \\ &= \alpha((qm \triangleright ys)(m \triangleright s)^{-1}(m \triangleright s)Y(T \cap Z)) \\ &= \alpha((qm \triangleright y)(q \triangleright (m \triangleright s))(m \triangleright s)^{-1}(m \triangleright s)Y(T \cap Z)) \\ &= \alpha(y'(m \triangleright s)Y(T \cap Z)) = (m \triangleright s)(Y \cap V)(T \cap Z) \\ &= m(Q \cap P)(N \cap R) \triangleright s(Y \cap V)(T \cap Z) \\ &= \gamma(qmQ(N \cap R)) \triangleright \alpha(ysY(T \cap Z)).\end{aligned}$$

5 where  $y' = (qm \triangleright y)(q \triangleright (m \triangleright s))(m \triangleright s)^{-1}$ . For  $hlH(K \cap G) \in \frac{H(K \cap J)}{H(K \cap G)}$ ,  $qmQ(N \cap R) \in \frac{Q(N \cap P)}{Q(N \cap R)}$

$$\begin{aligned}\beta(qmQ(N \cap R) \triangleright hlH(K \cap G)) &= \beta((qm \triangleright hl)H(K \cap G)) \\ &= \beta((qm \triangleright hl)(m \triangleright l)^{-1}(m \triangleright l)H(K \cap G)) \\ &= \beta((qm \triangleright h)(q \triangleright (m \triangleright l))(m \triangleright l)^{-1}(m \triangleright l)H(K \cap G)) \\ &= \beta(h'(m \triangleright l)H(K \cap G)) = (m \triangleright l)(H \cap J)(K \cap G) \\ &= m(Q \cap P)(N \cap R) \triangleright l(H \cap J)(K \cap G) \\ &= \gamma(qmQ(N \cap R)) \triangleright \beta(hlH(K \cap G)),\end{aligned}$$

6 where  $h' = (qm \triangleright h)(q \triangleright (m \triangleright l))(m \triangleright l)^{-1}$ .

For  $h_1 l_1 H(K \cap G), h_2 l_2 H(K \cap G) \in \frac{H(K \cap J)}{H(K \cap G)}$  noting that

$$(h_i l_i)(l_i h_i)^{-1} = h_i l_i h_i^{-1} l_i^{-1} \in H \subseteq H(K \cap G)$$

we have

$$h_i l_i H(K \cap G) = l_i h_i H(K \cap G) = l_i H(K \cap G)$$

and

$$\begin{aligned} \alpha\{h_1 l_1 H(K \cap G), h_2 l_2 H(K \cap G)\} &= \alpha\{l_1 H(K \cap G), l_2 H(K \cap G)\} \\ &= \alpha(\{l_1, l_2\}Y(T \cap Z)) = \alpha(1\{l_1, l_2\}Y(T \cap Z)) \\ &= \{l_1, l_2\}(Y \cap V)(T \cap Z) \\ &= \{l_1(H \cap J)(K \cap G), l_2(H \cap J)(K \cap G)\} \\ &= \left\{ \beta(h_1 l_1 H(K \cap G)), \beta(h_2 l_2 H(K \cap G)) \right\}. \end{aligned}$$

□

**Definition 3.24** A normal series (of lenght  $n$ ) of a 2-crossed module  $(S, L, M, \partial_2, \partial_1)$  consists of 2-subcrossed modules  $(S_i, L_i, M_i)$ ,  $i = 0, \dots, n$ , of  $(S, L, M)$  such that for each  $i = 1, \dots, n$ ,  $(S_i, L_i, M_i)$  is a normal 2-subcrossed module of  $(S_{i-1}, L_{i-1}, M_{i-1})$ ,  $(S_n, L_n, M_n) = (1, 1, 1)$  and  $(S_0, L_0, M_0) = (S, L, M)$ .

$$\begin{array}{ccccccc} 1 = S_n & \longrightarrow & S_{n-1} & \longrightarrow & \cdots & \longrightarrow & S_1 & \longrightarrow & S_0 = S \\ \downarrow & & \downarrow & & & & \downarrow & & \downarrow \partial_2 \\ 1 = L_n & \longrightarrow & L_{n-1} & \longrightarrow & \cdots & \longrightarrow & L_1 & \longrightarrow & L_0 = L \\ \downarrow & & \downarrow & & & & \downarrow & & \downarrow \partial_1 \\ 1 = M_n & \longrightarrow & M_{n-1} & \longrightarrow & \cdots & \longrightarrow & M_1 & \longrightarrow & M_0 = M \end{array}$$

If  $(S_i, L_i, M_i)$  is a normal series of  $(S, L, M)$ , then the quotients

$$\frac{(S_0, L_0, M_0)}{(S_1, L_1, M_1)}, \frac{(S_1, L_1, M_1)}{(S_2, L_2, M_2)}, \dots, \frac{(S_{n-1}, L_{n-1}, M_{n-1})}{(S_n, L_n, M_n)}$$

are called factor 2-crossed modules of  $(S, L, M)$ . If a normal series  $(S_i, L_i, M_i)$  of  $(S, L, M)$  can be obtained by eliminating some 2-crossed modules in a normal series  $(S'_i, L'_i, M'_i)$  of  $(S, L, M)$  then  $(S'_i, L'_i, M'_i)$  is called a refinement of  $(S_i, L_i, M_i)$ .

**Definition 3.25** A 2-crossed module  $(S, L, M, \partial_2, \partial_1)$  is called simple, if its only normal 2-subcrossed modules are  $(1, 1, 1)$  and  $(S, L, M)$ .

**Definition 3.26** A normal series of a 2-crossed module is said to be a composition series, if all nontrivial factor 2-crossed modules are simple.

**Definition 3.27** Two normal series of a 2-crossed module are said to be equivalent, if for each factor 2-crossed module corresponding to one of the normal series is isomorphic to some factor 2-crossed modules in the other normal series.

1 **Theorem 3.28 (Scherier Refinement Theorem for 2-crossed modules)** *Let*

$$\begin{array}{ccccccc}
 1 = T_n & \longrightarrow & T_{n-1} & \longrightarrow & \cdots & \longrightarrow & T_1 & \longrightarrow & T_0 = S \\
 \downarrow & & \downarrow & & & & \downarrow & & \downarrow \partial_2 \\
 1 = K_n & \longrightarrow & K_{n-1} & \longrightarrow & \cdots & \longrightarrow & K_1 & \longrightarrow & K_0 = L \\
 \downarrow & & \downarrow & & & & \downarrow & & \downarrow \partial_1 \\
 1 = N_n & \longrightarrow & N_{n-1} & \longrightarrow & \cdots & \longrightarrow & N_1 & \longrightarrow & N_0 = M
 \end{array}$$

2 *and*

$$\begin{array}{ccccccc}
 1 = V_m & \longrightarrow & V_{m-1} & \longrightarrow & \cdots & \longrightarrow & V_1 & \longrightarrow & V_0 = S \\
 \downarrow & & \downarrow & & & & \downarrow & & \downarrow \partial_2 \\
 1 = J_m & \longrightarrow & J_{m-1} & \longrightarrow & \cdots & \longrightarrow & J_1 & \longrightarrow & J_0 = L \\
 \downarrow & & \downarrow & & & & \downarrow & & \downarrow \partial_1 \\
 1 = P_m & \longrightarrow & P_{m-1} & \longrightarrow & \cdots & \longrightarrow & P_1 & \longrightarrow & P_0 = M
 \end{array}$$

3 *be two normal series of a 2-crossed module  $(S, L, M, \partial_2, \partial_1)$ . Then  $(T_i, K_i, N_i)$  and  $(V_i, J_i, M_i)$  have equivalent*  
 4 *refinements.*

**Proof** Define  $S_{ij} = T_{i+1}(T_i \cap V_j)$ ,  $L_{ij} = K_{i+1}(K_i \cap J_j)$  and  $M_{ij} = N_{i+1}(N_i \cap P_j)$  for  $0 \leq i \leq n-1$  and  $0 \leq j \leq m$ . Note that each  $(S_{ij}, L_{ij}, M_{ij})$  is a 2-subcrossed module of  $(S, L, M, \partial_2, \partial_1)$ , since  $(T_{i+1}, K_{i+1}, N_{i+1})$  is normal in  $(T_i, K_i, N_i)$ . Moreover

$$S_{i0} = T_{i+1}(T_i \cap V_0) = T_{i+1}(T_i \cap S) = T_{i+1}T_i = T_i$$

and

$$S_{im} = T_{i+1}(T_i \cap V_m) = T_{i+1}(T_i \cap 1) = T_{i+1}1 = T_{i+1}$$

5 and similarly  $L_{i0} = K_i$ ,  $M_{i0} = N_i$ ,  $L_{im} = K_{i+1}$ ,  $M_{im} = N_{i+1}$ . Thus we obtain a refinement of the normal  
 6 series  $(T_i, K_i, N_i)$ , namely

$$\begin{array}{ccccccc}
 1 = T_n = S_{(n-1)m} \twoheadrightarrow \cdots \twoheadrightarrow S_{(n-1)0} = T_{n-1} = S_{(n-2)m} \twoheadrightarrow \cdots \twoheadrightarrow S_{10} = T_1 = S_{0m} \twoheadrightarrow \cdots \twoheadrightarrow S_{00} = T_0 = S \\
 \downarrow & & \downarrow & & \downarrow & & \downarrow \partial_2 \\
 1 = K_n = L_{(n-1)m} \twoheadrightarrow \cdots \twoheadrightarrow L_{(n-1)0} = K_{n-1} = L_{(n-2)m} \twoheadrightarrow \cdots \twoheadrightarrow L_{10} = K_1 = L_{0m} \twoheadrightarrow \cdots \twoheadrightarrow L_{00} = K_0 = L \\
 \downarrow & & \downarrow & & \downarrow & & \downarrow \partial_1 \\
 1 = N_n = M_{(n-1)m} \twoheadrightarrow \cdots \twoheadrightarrow M_{(n-1)0} = N_{n-1} = M_{(n-2)m} \twoheadrightarrow \cdots \twoheadrightarrow M_{10} = N_1 = M_{0m} \twoheadrightarrow \cdots \twoheadrightarrow M_{00} = N_0 = M.
 \end{array}$$

In the same way we can construct a refinement of the normal series  $(V_i, J_i, M_i)$  by defining  $S'_{ij} =$

$V_{j+1}(V_j \cap T_i)$ ,  $L'_{ij} = J_{j+1}(J_j \cap K_i)$  and  $M'_{ij} = P_{j+1}(P_j \cap N_i)$ . Then by Lemma 3.23, we have isomorphisms

$$\frac{(T_{i+1}, K_{i+1}, N_{i+1})((T_i, K_i, N_i) \cap (V_j, J_j, P_j))}{(T_{i+1}, K_{i+1}, N_{i+1})((T_i, K_i, N_i) \cap (V_{j+1}, J_{j+1}, P_{j+1}))} \cong \frac{(V_{j+1}, J_{j+1}, P_{j+1})((V_j, J_j, P_j) \cap (T_i, K_i, N_i))}{(V_{j+1}, J_{j+1}, P_{j+1})((V_j, J_j, P_j) \cap (T_{i+1}, K_{i+1}, N_{i+1}))}$$

which completes the proof.  $\square$

Considering Definition 3.26, the following theorem is a direct consequence of the Schrier Refinement Theorem for 2-crossed modules.

**Theorem 3.29 (Jordan-Hölder Theorem for 2-crossed modules)** *All composition series of a 2-crossed module are equivalent to each other and have the same minimal length.*

## References

- [1] Boyacı Y, Casas JM, Datuashvili T, Uslu EÖ. Actions in modified categories of interest with application to crossed modules. *Theory and Applications of Categories* 2015; 30: 882-908.
- [2] Casas JM, Datuashvili T, Ladra M, Uslu EÖ. Actions in the category of precrossed modules in Lie algebras. *Communications in Algebra* 2012; 40: 2962-2982. <https://doi.org/10.1080/00927872.2011.588632>
- [3] Conduché D. Modules croisés généralisés de longueur 2. *Journal of Pure and Applied Algebra* 1984; 34 (2-3): 155-178. [https://doi.org/10.1016/0022-4049\(84\)90034-3](https://doi.org/10.1016/0022-4049(84)90034-3)
- [4] Datuashvili T, Şahan T. Actions and semi-direct products in categories of groups with action. *Hacettepe Journal of Mathematics and Statistics* 2023; 52 (1): 103-113. <https://doi.org/10.15672/hujms.1028848>
- [5] Datuashvili T, Şahan T. Pentactions and action representability in the category of reduced groups with action. *Georgian Mathematical Journal* 2023; 30 (2): 51-60. <https://doi.org/10.1515/gmj-2022-2205>
- [6] Davvaz B, Alp M. Simplicial polygroups and the generalized Moore complexes Quasigroups and Related Systems 2022; 30 (1): 47-62.
- [7] Davvaz B, Emir K. On morphisms of crossed polymodules. *Journal of Linear and Topological Algebra* 2020; 9 (1): 95-103.
- [8] Dehghani MA, Davvaz B, Alp M. On crossed polysquare version of homotopy kernels. *Journal of Mathematical Extension* 2021; 16: 1-37.
- [9] Dehghani MA, Davvaz B, Alp M. On the 2-crossed polymodules, pullback and pushout 2-crossed polymodules. *Advanced Studies: Euro-Tbilisi Mathematical Journal* 2023; 16 (1): 27-45. <https://doi.org/10.32513/asetmj/1932200823105>
- [10] Gürdal U. A Jordan-Hölder theorem for crossed squares. *Kuwait Journal of Science* 2023; 50 (2): 83-90. <https://doi.org/10.1016/j.kjs.2023.02.012>
- [11] Hidaka Y, Nitta M, Yokokura R. Global 3-group symmetry and 't Hooft anomalies in axion electrodynamics. *Journal of High Energy Physics* 2021; 1: 1-53. [https://doi.org/10.1007/JHEP01\(2021\)173](https://doi.org/10.1007/JHEP01(2021)173)
- [12] Lang S. *Algebra*. New York, NY, USA: Springer-Verlag, 2002. <https://doi.org/10.1007/978-1-4613-0041-0>
- [13] Lue AST. Semi-complete crossed modules and holomorphs of groups. *Bulletin of the London Mathematical Society* 1979; 11: 8-16. <https://doi.org/10.1112/blms/11.1.8>
- [14] Mucuk O, Şahan T, Alemdar N. Normality and quotients in crossed modules and group-groupoids. *Applied Categorical Structures* 2015; 23 (3): 415-428. <https://doi.org/10.1007/s10485-013-9335-6>
- [15] Mutlu A, Mutlu B. Freeness conditions for quasi 3-crossed modules and complexes of using simplicial algebras with CW-bases. *Mathematical Sciences* 2013; 7: 1-12. <https://doi.org/10.1186/2251-7456-7-35>

- 1 [16] Mutlu A, Mutlu B. On Semi 3-crossed module by using simplicial algebra. Global Journal of Science Frontier  
2 Research 2012; 12: 81-90. <https://doi.org/10.1186/2251-7456-7-35>
- 3 [17] Mutlu A, Mutlu B. Quasi 3-crossed modules. Theoretical Mathematics & Applications 2012; 2: 101-118.
- 4 [18] Mutlu A, Porter T. Freeness conditions for 2-crossed modules and complexes. Theory and Applications of Cate-  
5 gories 1998; 4 (8): 174-194.
- 6 [19] Norrie KJ. Crossed modules and analogues of group Theorems. PhD, University of London, London, United  
7 Kingdom, 1987.
- 8 [20] Onarlı G. 2-çaprazlanmış modüller üzerine. PhD, Eskişehir Osmangazi University, Eskişehir, Türkiye, 2013.
- 9 [21] Rotman JJ. Advanced Modern Algebra. Urbana, IL, USA: American Mathematical Society, 2002.
- 10 [22] Tercan A, Yücel CC. Module Theory, Extending Modules and Generalizations. Basel, Switzerland:  
11 Birkhäuser-Springer, 2016. <https://doi.org/10.1007/978-3-0348-0952-8>
- 12 [23] Whitehead JHC. Combinatorial homotopy II. Bulletin of the American Mathematical Society 1944; 55: 453-496.  
13 <https://doi.org/10.1090/S0002-9904-1949-09213-3>
- 14 [24] Whitehead JHC. On operators in relative homotopy groups. Annals of Mathematics. 1948; 2 (49): 610-640. <https://doi.org/10.2307/1969048>  
15