


Solvability of Gripenberg's equations of fractional order with perturbation term in weighted L_p -spaces on \mathbb{R}^+

Mohamed M. A. METWALI* 

Department of Mathematics, Faculty of Sciences, Damanhour University, Damanhour, Egypt

Received: 20.06.2021

Accepted/Published Online: 08.09.2021

Final Version: ..2021

Abstract: This article deals with the solvability of Gripenberg's equations of fractional order with a perturbation term in weighted Lebesgue spaces on $\mathbb{R}^+ = [0, \infty)$ via the fixed point hypothesis and the measure of noncompactness. The uniqueness of the solutions for the studied problem is discussed. An example is included to validate our results. The results presented in the article extend and generalize some former results in the available literature.

Key words: Measure of noncompactness, Gripenberg's equations, fractional calculus, weighted Lebesgue spaces, Schauder fixed point theorem

1. Introduction

The integral equations of Gripenberg's type of the form

$$x(\tau) = k \left(g_1(\tau) + \int_0^\tau a_1(\tau - s)x(s)ds \right) \left(g_2(\tau) + \int_0^\tau a_2(\tau - s)x(s)ds \right), \tau \in \mathbb{R}^+$$

have numerous applications in mathematical modeling, in particular in the models of the spread of diseases, that do not induce permanent immunity. Some qualitative properties of the solutions of models such as existence and uniqueness of continuous solutions were studied in [1, 12, 20, 29].

Additionally, numerous types of infectious diseases models consist of discontinuous data functions, so it is recommended to examine the integrable solutions of models.

The purpose of the article is to examine the solutions of the equation

$$x(\tau) = h(\tau, x(\varphi_3(\tau))) + \left(g_1(\tau) + g_3(\tau) \cdot (Gx)(\varphi_1(\tau)) \right) \left(g_2(\tau) + \frac{1}{\Gamma(\alpha)} \int_0^\tau \frac{u(s, x(\varphi_2(s)))}{(\tau - s)^{1-\alpha}} ds \right), \tau \in \mathbb{R}^+ \quad (1.1)$$

in weighted Lebesgue spaces $L_1^N(\mathbb{R}^+)$, where $N > 0$, $0 < \alpha < 1$.

In [22] the authors have discussed the existence of solutions in weighted L_p -spaces for Hammerstein integral equations

$$x(\tau) = h(\tau) + \int_0^\infty a(\tau, s)f(s, x(s)) ds$$

*Correspondence: metwali@sci.dmu.edu.eg

2010 AMS Mathematics Subject Classification: 45G10, 47H30, 47N20

and integral equations of Urysohn-type

$$x(\tau) = h(\tau) + \int_0^\infty u(\tau, s, x(s)) ds$$

using Schauder fixed point hypothesis. The authors in [24] have discussed the existence results of Volterra integral equation by utilizing the Banach contraction mapping principle in the considered spaces.

Recall that, the idea of the measure of noncompactness were utilized to demonstrate the presence of continuous or L_p -solutions of the integral equations on compact or noncompact domains in [3, 5, 13, 15–17, 21, 25, 33] and for qualitative properties of the outcomes (cf. [9, 10, 26, 32]).

Further, in [8] the authors examined the solutions of the equation

$$x(\tau) = h(\tau, x(\tau)) + \left(p_1(\tau) + \int_0^\tau k_1(\tau, s)g_1(s, x(s))ds \right) \left(p_2(\tau) + \int_0^\tau k_2(\tau, s)g_2(s, x(s))ds \right)$$

in the space $L_1([0, 1])$ and these results were extended to an unbounded interval in [11].

In [27] the author used the measure of weak noncompactness with Krasnoselskii fixed point theory to find $L_1(\mathbb{R}^+)$ -solutions of the equation

$$x(\tau) = h(\tau, x(\varphi_3(\tau))) + g\left(\tau, f(\tau, x(\varphi_2(\tau))) \int_0^\tau u(\tau, s, x(\varphi_1(s)))\right), \tau \in \mathbb{R}^+.$$

Nevertheless, to investigate our results, we use triple of weighted spaces (L_1^N, L_p^N, L_q^N) . One more challenge is to examine the integrable solutions for fractional integral equation on unbounded domains \mathbb{R}^+ .

Hence, we dedicate our investigations to expand the previously mentioned results as particular cases of Equation (1.1) in the weighted Lebesgue spaces on $L_1^N(\mathbb{R}^+)$ using a different set of assumptions. Moreover, the uniqueness of the solutions of Equation (1.1) are also discussed. At the end, we give an example to validate the applicability of our results. Our methodology utilizes Schauder’s fixed point hypothesis and appropriate measure of noncompactness.

2. Notation and auxiliary facts

Let $\mathbb{R} = (-\infty, \infty)$, $\mathbb{R}^+ = [0, \infty)$ and Φ be the empty set. Denote by $L_p^N = L_p^N(\mathbb{R}^+)$, $1 \leq p < \infty$ the weighted Lebesgue spaces, which is the spaces of measurable functions x with

$$\|x\|_{L_p^N} = \left(\int_0^\infty e^{-N\tau} |x(\tau)|^p d\tau \right)^{\frac{1}{p}} < \infty, \quad N > 0.$$

When $N = 0$ we obtain classical L_p -spaces for $1 \leq p < \infty$ with standard norm.

Let $C(D)$ refer to the space of real and continuous functions on a closed and bounded set $\Phi \neq D \subset \mathbb{R}$. Let $\Phi \neq X \subset C(D)$ be bounded set and $T > 0$.

For $\varepsilon > 0$ and $x \in X$, we denote by $\omega^T(x, \varepsilon)$ the modulus of continuity of the function x , on the interval $[0, T]$, where

$$\omega^T(x, \varepsilon) = \sup\{|x(\tau_2) - x(\tau_1)| : \tau_1, \tau_2 \in [0, T], |\tau_1 - \tau_2| \leq \varepsilon\}.$$

Suppose that $(E, \|\cdot\|)$ is a Banach space and has zero element θ and $B_r = \{x \in L_1^N : \|x\|_{L_1^N} \leq r\}$, $r > 0$. Let $\Phi \neq \mathcal{M}_E$ and \mathcal{N}_E^W be the family of all bounded subsets and the subfamily containing all relatively weakly compact sets of E , respectively. The symbols $\text{Conv}Y$ and \bar{Y}^W stand for the convex closed hull and the weak closure of a set Y , respectively.

Definition 2.1 [6] *The operator $\mu : \mathcal{M}_E \rightarrow [0, \infty)$ refers to a regular measure of weak noncompactness in E if it fulfills:*

- (i) $\mu(Y) = 0 \iff Y \in \mathcal{N}_E^W$.
- (ii) $Y \subset X \Rightarrow \mu(Y) \leq \mu(X)$.
- (iii) $\mu(\bar{Y}) = \mu(\text{Conv}Y) = \mu(Y)$.
- (iv) $\mu(\lambda Y) = |\lambda| \mu(Y)$, for $\lambda \in \mathbb{R}$.
- (v) $\mu(Y + X) \leq \mu(Y) + \mu(X)$.
- (vi) $\mu(Y \cup X) = \max\{\mu(Y), \mu(X)\}$.
- (vii) *If $\Phi \neq Y_n \subset E$ is a sequence of closed and bounded sets, $Y_n = \bar{Y}_n^W$ s.t. $Y_{n+1} \subset Y_n$, $n = 1, 2, 3, \dots$, and $\lim_{n \rightarrow \infty} \mu(Y_n) = 0$, then $Y_\infty = \bigcap_{n=1}^\infty Y_n \neq \Phi$.*

De Blasi presented the next measure of weak noncompactness β (cf. [7]):

$$\beta(X) = \inf\{r > 0 : \text{there exists a weakly compact subset } Y \text{ of } E \text{ s.t. } X \subset Y + B_r\}.$$

Now, for a bounded set $\Phi \neq X \subset L_1$, $\varepsilon > 0$ and $T > 0$, consider the following:

$$c^T(X) = \lim_{\varepsilon \rightarrow 0} \left\{ \sup_{x \in X} \left\{ \sup \left(\int_D |x(\tau)| d\tau : D \subset [0, T], \text{meas}(D) \leq \varepsilon \right) \right\} \right\},$$

$$c(X) = \lim_{T \rightarrow \infty} c^T(X), \tag{2.1}$$

where $\text{meas}(D)$ refers to the Lebesgue measure of the set D and let

$$d(X) = \lim_{T \rightarrow \infty} \sup \left\{ \int_T^\infty |x(\tau)| d\tau : x \in X \right\}. \tag{2.2}$$

Put

$$\gamma(X) = c(X) + d(X). \tag{2.3}$$

From [2, 7] we know the function $\gamma(X)$ represents a measure of weak noncompactness in L_1 and $\beta(X) \leq \gamma(X) \leq 2\beta(X)$.

Theorem 2.2 [18] *Let $Q \neq \Phi$ be a convex, bounded, and closed subset of the Banach space E . Let $H : Q \rightarrow Q$ be a completely continuous mapping. Hence, H has at least one fixed point in Q .*

Definition 2.3 [4] Suppose that a function $u(\tau, x) : \mathbb{R}^+ \times \mathbb{R} \rightarrow \mathbb{R}$ fulfills the Carathéodory conditions, i.e. it is continuous in x for almost all $\tau \in \mathbb{R}^+$ and measurable in τ for any $x \in \mathbb{R}$. Then to every measurable function x , we can assign the superposition (Nemytskii) operator F_u as follows.

$$F_u(x)(\tau) = u(\tau, x(\tau)), \quad \tau \in \mathbb{R}^+.$$

Moreover, for every functions $u \in L_1$ and $\varphi : \mathbb{R}^+ \rightarrow \mathbb{R}^+$, we denote the operator $F_{\varphi, u}$, as

$$F_{\varphi, u}(x)(\tau) = u(\tau, x(\varphi(\tau))), \quad \tau \in \mathbb{R}^+.$$

Lemma 2.4 Assume that the function u fulfills the Carathéodory conditions and \exists a function $a \in L_q^N$ and $b \geq 0$ s.t.

$$|u(\tau, x)| \leq a(\tau) + b \cdot |x|^{\frac{p}{q}} \tag{2.4}$$

for all $\tau \in \mathbb{R}^+$ and $x \in \mathbb{R}$. Then the operator $F_u : L_p^N \rightarrow L_q^N$ ($p, q \geq 1$) is continuous.

Proof By using (2.4), we have

$$\begin{aligned} \|F_u\|_{L_q^N} &= \left(\int_0^\infty e^{-N\tau} |u(\tau, x(\tau))|^q d\tau \right)^{\frac{1}{q}} \\ &\leq \left(\int_0^\infty e^{-N\tau} \left(a(\tau) + b \cdot |x(\tau)|^{\frac{p}{q}} \right)^q d\tau \right)^{\frac{1}{q}} \\ &\leq \left(\int_0^\infty e^{-N\tau} |a(\tau)|^q d\tau \right)^{\frac{1}{q}} + b \left(\int_0^\infty e^{-N\tau} |x(\tau)|^p d\tau \right)^{\frac{p}{pq}} \\ &\leq \|a\|_{L_q^N} + b \|x\|_{L_p^N}^{\frac{p}{q}}. \end{aligned}$$

Thus $F_u : L_p^N \rightarrow L_q^N$ and is continuous by using the well known results in [4]. □

Furthermore, let D^c stands for the complement of D and let J be an interval in the next theorems.

Theorem 2.5 Let $m : J \rightarrow \mathbb{R}$ be a measurable function. \exists a closed set $D_\varepsilon \subset J$, $\varepsilon > 0$ s.t. $m|_{D_\varepsilon}$ is continuous with $meas(D_\varepsilon^c) \leq \varepsilon$.

Theorem 2.6 [31] Let $u : J \times \mathbb{R} \rightarrow \mathbb{R}$ be a function fulfilling Carathéodory conditions. \exists a closed set $D_\varepsilon \subset J$, $\varepsilon > 0$ s.t. $f|_{D_\varepsilon \times \mathbb{R}}$ is continuous with $meas(D_\varepsilon^c) \leq \varepsilon$.

These results were proved in [28] for general operator $G : Z \times X \rightarrow Y$, where Z is a topological space with finite Borel regular measure, X is a metrizable separable locally compact space, and Y is topological spaces.

Next, we give the definition of fractional operator.

Definition 2.7 [23, 26, 30] Let $y \in L_1$, and $\alpha \in \mathbb{R}^+$. The fractional integral of Riemman-Liouville type (R-L) of the function y of order α is given by

$$I^\alpha y(\tau) = \frac{1}{\Gamma(\alpha)} \int_0^\tau \frac{y(s)}{(\tau - s)^{1-\alpha}} ds, \quad \alpha > 0, \quad \tau > 0,$$

where $\Gamma(\alpha) = \int_0^\infty e^{-s} s^{\alpha-1} ds$.

3. Main results

Let us write (1.1) as follows:

$$x = (Hx) = F_{\varphi_3, h}(x) + (Bx) \cdot (Ax),$$

$$(Bx)(\tau) = g_1(\tau) + g_3(\tau) \cdot (Gx)(\varphi_1(\tau)), \quad (Ax)(\tau) = g_2(\tau) + I^\alpha F_{\varphi_2, u}(x)(\tau),$$

where $F_{\varphi_3, h}, F_{\varphi_2, u}$ are the superposition operators, as in Definition 2.3, I^α as in Definition 2.7 and (Gx) is an operator maps L_1^N into L_q^N continuously.

Let $\frac{1}{p} + \frac{1}{q} = 1$ and write the assumptions:

- (i) $g_i : \mathbb{R}^+ \rightarrow \mathbb{R}$, $i = 1, 2, 3$, where $g_1 \in L_q^N$, $g_2 \in L_p^N$ and g_3 is bounded function s.t. $\sup_{\tau \in \mathbb{R}^+} |g_3(\tau)| \leq M$.
- (ii) Suppose that $u, h, G : \mathbb{R}^+ \times \mathbb{R} \rightarrow \mathbb{R}$ fulfill Carathéodory conditions. Further, assume that $G : L_1^N \rightarrow L_q^N$ is continuous.
- (iii) $\exists b_i \geq 0$ ($i = 1, 2, 3$) and positive functions $a_1 \in L_q^N$, $a_2 \in L_p^N$, and $a_3 \in L_1^N$ s.t.

$$|Gx| \leq a_1(\tau) + b_1|x|^{\frac{1}{q}}, \quad |u(\tau, x)| \leq a_2(\tau) + b_2|x|^{\frac{1}{p}}$$

and

$$|h(\tau, x)| \leq a_3(\tau) + b_3|x|$$

for all $\tau \in \mathbb{R}^+$ and $x \in \mathbb{R}$.

- (iv) $\varphi_i : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ are increasing and absolutely continuous functions. Further, $\exists M_i > 0$ s.t. $\varphi_i' \geq M_i$ a.e. on $(0, \infty)$ and $\varphi_i^{-1}(\tau) \geq \tau$, $i = 1, 2, 3$.
- (v) Assume that $\exists N > 0$, s.t.

$$W = e^{N^2} \left(\frac{b_3}{M_3} + \frac{Mb_1b_2p^\alpha}{N^\alpha M_1^{\frac{1}{q}} M_2^{\frac{1}{p}}} \right) < 1.$$

- (vi) Assume that r is a positive number satisfying

$$\begin{aligned} \|a_3\|_{L_1^N} + \left(\|g_1\|_{L_q^N} + M\|a_1\|_{L_q^N} \right) \left(\|g_2\|_{L_p^N} + \left(\frac{p}{N} \right)^\alpha \|a_2\|_{L_p^N} \right) + \left(\|g_1\|_{L_q^N} + M\|a_1\|_{L_q^N} \right) \frac{b_2p^\alpha}{N^\alpha M_2^{\frac{1}{p}}} \cdot r^{\frac{1}{p}} \\ + \left(\|g_2\|_{L_p^N} + \left(\frac{p}{N} \right)^\alpha \|a_2\|_{L_p^N} \right) \frac{Mb_1}{M_1^{\frac{1}{q}}} \cdot r^{\frac{1}{q}} + \left(\frac{b_3}{M_3} + \frac{Mb_1b_2p^\alpha}{N^\alpha M_1^{\frac{1}{q}} M_2^{\frac{1}{p}}} \right) \cdot r \leq r. \end{aligned}$$

Proposition 3.1 *We can easily check the followings:*

(a) Assume that the functions $g \in L_q^N$ and $f \in L_p^N$ s.t. $\frac{1}{p} + \frac{1}{q} = 1$, then we have the following inequality

$$\begin{aligned} \|g \cdot f\|_{L_1^N} &= \int_0^\infty e^{-N\tau} |g(\tau)f(\tau)| d\tau \\ &= \int_0^\infty (e^{-N\tau} * e^{-\frac{N\tau}{p}}) e^{-\frac{N\tau}{q}} |g(\tau)f(\tau)| d\tau \\ &\leq \left(\int_0^\infty \left(e^{-N\tau(1-\frac{1}{p})} |g(\tau)| \right)^q d\tau \right)^{\frac{1}{q}} \left(\int_0^\infty \left(e^{-\frac{N\tau}{p}} |f(\tau)| \right)^p d\tau \right)^{\frac{1}{p}} \\ &\leq \|g\|_{L_q^N} \cdot \|f\|_{L_p^N}. \end{aligned}$$

(b) The operator I^α , $\alpha > 0$ is bounded in L_p^N , $1 \leq p \leq \infty$ (cf. [30, Theorem 5.7] see also [23]) s.t.

$$(1) \|I^\alpha f\|_{L_p^N} \leq \left(\frac{p}{N}\right)^\alpha \|f\|_{L_p^N} \text{ for } 1 \leq p < \infty.$$

$$(2) \|I^\alpha f\|_{L_p^N} \leq \left(\frac{1}{N}\right)^\alpha \|f\|_{L_p^N} \text{ for } p = \infty.$$

(3) The operator $I^\alpha : L_p^N \rightarrow L_p^N$ and is continuous.

(c) Let, $D \subset \mathbb{R}^+$ with $\text{meas}(D) \leq \varepsilon$, $\varepsilon > 0$ be arbitrary. For $0 \leq \tau \leq N$, $N > 0$ and by using assumption (iv), we have

$$(1) e^{-N^2} \int_D |x(\tau)| d\tau \leq \int_D e^{-N\tau} |x(\tau)| d\tau \leq \int_D |x(\tau)| d\tau.$$

$$(2) \|(x(\varphi_i))\|_{L_p^N}^{\frac{1}{p}} \leq \frac{1}{M_i^{\frac{1}{p}}} \left(\int_0^\infty e^{-Ns} |x(\varphi_i(s))| |\varphi_i'(s)| ds \right)^{\frac{1}{p}} \leq \frac{1}{M_i^{\frac{1}{p}}} \|x\|_{L_1^N}^{\frac{1}{p}}.$$

$$(3) \|(x(\varphi_i))\|_{L_p^N(D)}^{\frac{1}{p}} \leq \frac{1}{M_i^{\frac{1}{p}}} \left(\int_{\varphi_i(D)} e^{-Nu} |x(u)| du \right)^{\frac{1}{p}} \leq \frac{1}{M_i^{\frac{1}{p}}} \left(\int_{\varphi_i(D)} |x(u)| du \right)^{\frac{1}{p}}.$$

(d) Assumption (vi) takes the form $C + Dr^{\frac{1}{p}} + Kr^{\frac{1}{q}} + Lr \leq r$. If $r = 1$, we would have $C + D + K + L \leq 1$.

Theorem 3.2 Let the assumptions (i)–(vi) be fulfilled, then Equation (1.1) has at least one integrable solution $x \in L_1^N$ on \mathbb{R}^+ .

Proof Step 1. By assumptions (ii), (iii) and Lemma 2.4 we get $F_{\varphi_2, u}$ is continuous mappings from L_1^N into L_p^N and $F_{\varphi_3, h}$ is continuous mappings from L_1^N into itself. Since the operator $I^\alpha : L_p^N \rightarrow L_p^N$ is continuous, assumption (i) gives that $A : L_1^N \rightarrow L_p^N$ is continuous and $B : L_1^N \rightarrow L_q^N$ is continuous. At last, the Hölder inequality gives us that $H : L_1^N \rightarrow L_1^N$ is continuous.

Step 2. In view of our assumptions and Proposition 3.1, we get

$$\begin{aligned}
 \|Hx\|_{L_1^N} &= \|F_{\varphi_3, h} + Bx \cdot Ax\|_{L_1^N} \\
 &\leq \|F_{\varphi_3, h}\|_{L_1^N} + \|Bx \cdot Ax\|_{L_1^N} \\
 &\leq \|h(\tau, x(\varphi_3))\|_{L_1^N} + \|Bx\|_{L_q^N} \cdot \|Ax\|_{L_p^N} \\
 &\leq \left\| a_3 + b_3|x(\varphi_3)| \right\|_{L_1^N} + \left\| g_1 + g_3 \cdot Gx(\varphi_1) \right\|_{L_q^N} \cdot \left\| g_2 + I^\alpha F_{\varphi_2, u}(x) \right\|_{L_p^N} \\
 &\leq \|a_3\|_{L_1^N} + b_3\|x(\varphi_3)\|_{L_1^N} \\
 &\quad + \left(\|g_1\|_{L_q^N} + M\|Gx(\varphi_1)\|_{L_q^N} \right) \cdot \left(\|g_2\|_{L_p^N} + \|I^\alpha F_{\varphi_2, u}(x)\|_{L_p^N} \right) \\
 &\leq \|a_3\|_{L_1^N} + \frac{b_3}{M_3}\|x\|_{L_1^N} + \left(\|g_1\|_{L_q^N} + M \left(\|a_1\|_{L_q^N} + b_1\|(x(\varphi_1))\|^{\frac{1}{q}}\|_{L_q^N} \right) \right) \\
 &\quad \times \left(\|g_2\|_{L_p^N} + \left(\frac{p}{N} \right)^\alpha \left\| u(\tau, x(\varphi_2)) \right\|_{L_p^N} \right) \\
 &\leq \|a_3\|_{L_1^N} + \frac{b_3}{M_3}\|x\|_{L_1^N} + \left(\|g_1\|_{L_q^N} + M\|a_1\|_{L_q^N} + \frac{b_1M}{M_1^{\frac{1}{q}}}\|x\|_{L_1^N}^{\frac{1}{q}} \right) \\
 &\quad \times \left(\|g_2\|_{L_p^N} + \left(\frac{p}{N} \right)^\alpha \left\| a_2 + b_2(x(\varphi_2))^{\frac{1}{p}} \right\|_{L_p^N} \right) \\
 &\leq \|a_3\|_{L_1^N} + \frac{b_3}{M_3}\|x\|_{L_1^N} + \left(\|g_1\|_{L_q^N} + M\|a_1\|_{L_q^N} + \frac{b_1M}{M_1^{\frac{1}{q}}}\|x\|_{L_1^N}^{\frac{1}{q}} \right) \\
 &\quad \times \left(\|g_2\|_{L_p^N} + \left(\frac{p}{N} \right)^\alpha \left(\|a_2\|_{L_p^N} + \frac{b_2}{M_2^{\frac{1}{p}}}\|x\|_{L_1^N}^{\frac{1}{p}} \right) \right).
 \end{aligned}$$

Thus $H : L_1^N \rightarrow L_1^N$. For $x \in B_r$, where r is as in assumption (vi) and $B_r = \{m \in L_1^N : \|m\|_{L_1^N} \leq r\}$. Then

$$\begin{aligned}
 \|Hx\|_{L_1^N} &\leq \|a_3\|_{L_1^N} + \left(\|g_1\|_{L_q^N} + M\|a_1\|_{L_q^N} \right) \left(\|g_2\|_{L_p^N} + \left(\frac{p}{N} \right)^\alpha \|a_2\|_{L_p^N} \right) \\
 &\quad + \left(\|g_1\|_{L_q^N} + M\|a_1\|_{L_q^N} \right) \frac{b_2p^\alpha}{N^\alpha M_2^{\frac{1}{p}}} \cdot r^{\frac{1}{p}} + \left(\|g_2\|_{L_p^N} + \left(\frac{p}{N} \right)^\alpha \|a_2\|_{L_p^N} \right) \frac{Mb_1}{M_1^{\frac{1}{q}}} \cdot r^{\frac{1}{q}} \\
 &\quad + \left(\frac{b_3}{M_3} + \frac{Mb_1b_2p^\alpha}{N^\alpha M_1^{\frac{1}{q}} M_2^{\frac{1}{p}}} \right) \cdot r \leq r.
 \end{aligned}$$

Thus $H : B_r \rightarrow B_r$ is continuous.

Step 3. In what follows, let $\Phi \neq X \subset B_r$. For arbitrary fixed $\varepsilon > 0$, $T > 0$ s.t. for any $D \subset [0, T]$ with

$meas(D) \leq \varepsilon$. By using Proposition 3.1 for $x \in X$, we derive

$$\begin{aligned}
 e^{-N^2} \int_D |(Hx)(\tau)| d\tau &\leq \int_D e^{-N\tau} |(Hx)(\tau)| d\tau \\
 &\leq \|F_{\varphi_3, hx}\|_{L_1^N(D)} + \|Bx\|_{L_q^N(D)} \cdot \|Ax\|_{L_p^N(D)} \\
 &\leq \|a_3\|_{L_1^N(D)} + b_3 \|x(\varphi_3)\|_{L_1^N(D)} \\
 &+ \left[\|g_1\|_{L_q^N(D)} + M \|a_1\|_{L_q^N(D)} + Mb_1 \|(x(\varphi_1))^{\frac{1}{q}}\|_{L_q^N(D)} \right] \left[\|g_2\|_{L_p^N(D)} \right. \\
 &+ \left. \left(\frac{p}{N} \right)^\alpha \left(\|a_2\|_{L_p^N(D)} + b_2 \|(x(\varphi_2))^{\frac{1}{p}}\|_{L_p^N(D)} \right) \right] \\
 &\leq \|a_3\|_{L_1^N(D)} + \frac{b_3}{M_3} \int_{\varphi_3(D)} |x(u)| du \\
 &+ \left[\|g_1\|_{L_q^N(D)} + M \|a_1\|_{L_q^N(D)} + \frac{Mb_1}{M_1^{\frac{1}{q}}} \left(\int_{\varphi_1(D)} |x(u)| du \right)^{\frac{1}{q}} \right] \left[\|g_2\|_{L_p^N(D)} \right. \\
 &+ \left. \left(\frac{p}{N} \right)^\alpha \|a_2\|_{L_p^N(D)} + \frac{b_2 p^\alpha}{N^\alpha M_2^{\frac{1}{p}}} \left(\int_{\varphi_2(D)} |x(u)| du \right)^{\frac{1}{p}} \right].
 \end{aligned}$$

Since $g_1, a_1 \in L_q^N$, $g_2, a_2 \in L_p^N$ and $a_3 \in L_1^N$, we have

$$\lim_{\varepsilon \rightarrow 0} \left\{ \sup_{x \in X} \left\{ \sup \left(\|a_3\|_{L_1^N(D)} : D \subset \mathbb{R}^+, meas(D) \leq \varepsilon \right) \right\} \right\} = 0,$$

$$\lim_{\varepsilon \rightarrow 0} \left\{ \sup_{x \in X} \left\{ \sup \left(\|g_1\|_{L_q^N(D)} + M \|a_1\|_{L_q^N(D)} : D \subset \mathbb{R}^+, meas(D) \leq \varepsilon \right) \right\} \right\} = 0$$

and

$$\lim_{\varepsilon \rightarrow 0} \left\{ \sup_{x \in X} \left\{ \sup \left(\|g_2\|_{L_p^N(D)} + \left(\frac{p}{N} \right)^\alpha \|a_2\|_{L_p^N(D)} : D \subset \mathbb{R}^+, meas(D) \leq \varepsilon \right) \right\} \right\} = 0.$$

From Equation (2.1) and assumption (v), we have

$$c(HX) \leq W \cdot c(X). \tag{3.1}$$

For $T > 0$ and $x \in X$, we get

$$\begin{aligned}
 e^{-N^2} \int_T^\infty |(Hx)(\tau)| d\tau &\leq \|a_3\|_{L_1^N(T)} + \frac{b_3}{M_3} \int_{\varphi_3(T)}^\infty |x(u)| du \\
 &+ \left[\|g_1\|_{L_q^N(T)} + M \|a_1\|_{L_q^N(T)} + \frac{Mb_1}{M_1^{\frac{1}{q}}} \left(\int_{\varphi_1(T)}^\infty |x(u)| du \right)^{\frac{1}{q}} \right] \left[\|g_2\|_{L_p^N(T)} \right. \\
 &+ \left. \left(\frac{p}{N} \right)^\alpha \|a_2\|_{L_p^N(T)} + \frac{b_2 p^\alpha}{N^\alpha M_2^{\frac{1}{p}}} \left(\int_{\varphi_2(T)}^\infty |x(u)| du \right)^{\frac{1}{p}} \right].
 \end{aligned}$$

Then as $T \rightarrow \infty$, we have $\varphi_i(T) \rightarrow \infty$, $i = 1, 2, 3$ and from Equation (2.2), we obtain

$$d(HX) \leq W \cdot d(X). \tag{3.2}$$

By adding Equations (3.1) and (3.2) and from Equation (2.3), we have

$$\gamma(HX) \leq W \cdot \gamma(X). \tag{3.3}$$

Step 4. Recall that B_r is defined in Step 2 and define $B_r^1 = Conv(H(B_r))$, $B_r^2 = Conv(H(B_r^1))$ and so on. We have the sequence $\{B_r^n\}$, which is decreasing i.e. $B_r^{n+1} \subset B_r^n$ for $n = 1, 2, \dots$. Clearly, all sets from that sequence are convex and closed, so weakly closed. From Step 3, that $\gamma(HX) \leq W\gamma(X)$ for all bounded set $X \subset B_r$, we have

$$\gamma(B_r^n) \leq W^n \gamma(B_r),$$

which implies $\lim_{n \rightarrow \infty} \gamma(B_r^n) = 0$. Then, axiom (vii) of Definition 2.1 gives us the set $Q = \cap_{n=1}^{\infty} B_r^n \neq \Phi$ is convex, closed, and weakly compact (in virtue of $\gamma(Q) = 0$). Moreover, $H(Q) \subset Q$.

Step 5. Since $\gamma(Q) = 0$, and for arbitrary sequence $\{x_n\} \subset Q$, $\exists T, \forall n$, s.t. the next inequality holds

$$\int_T^{\infty} e^{-N\tau} |x_n(\tau)| d\tau \leq \frac{\varepsilon}{4}. \tag{3.4}$$

Considering the functions $u(\tau, x)$, $h(\tau, x)$ and $Gx(\tau)$ on $[0, T] \times \mathbb{R}$, then by Theorems 2.5 and 2.6 we can find a closed set $D_\varepsilon \subset [0, T]$, s.t. $meas(D_\varepsilon^c) \leq \varepsilon$, $g_i|_{D_\varepsilon}$, $i = 1, 2, 3$ are continuous and $u|_{D_\varepsilon \times \mathbb{R}}$, $h|_{D_\varepsilon \times \mathbb{R}}$, $G|_{D_\varepsilon \times \mathbb{R}}$ are continuous.

Denoting by $U_n(\tau) = (Bx_n) \cdot (Ax_n)(\tau)$, $n \in \mathbb{N}$. First, we will prove that the operator $(U_n)(\tau)$ is equicontinuous and equibounded in the space $C(D_\varepsilon)$. Thus

$$\begin{aligned} |U_n(\tau)| &\leq \left(|g_1(\tau)| + |g_3(\tau)| |(Gx_n)(\varphi_1(\tau))| \right) \left(|g_2(\tau)| + \frac{1}{\Gamma(\alpha)} \int_0^\tau \frac{|u(s, x_n(\varphi_2(s)))|}{(\tau - s)^{1-\alpha}} ds \right) \\ &\leq \left(d_5 + M \left(|a_1(\tau)| + b_1 |x_n(\varphi_1(\tau))|^{\frac{1}{q}} \right) \right) \cdot \left(d_6 + \frac{1}{\Gamma(\alpha)} \int_0^\tau \frac{|a_2(s)| + b_2 |x_n(\varphi_2(s))|^{\frac{1}{p}}}{(\tau - s)^{1-\alpha}} ds \right) \\ &\leq \left(d_5 + M \left(d_1 + b_1 d_3^{\frac{1}{q}} \right) \right) \cdot \left(d_6 + \frac{1}{\Gamma(\alpha)} \int_0^\tau \frac{d_2 + b_2 d_4^{\frac{1}{p}}}{(\tau - s)^{1-\alpha}} ds \right) \\ &\leq \left(d_5 + M \left(d_1 + b_1 d_3^{\frac{1}{q}} \right) \right) \cdot \left(d_6 + \frac{d_2 + b_2 d_4^{\frac{1}{p}}}{\Gamma(\alpha + 1)} T^\alpha \right), \end{aligned}$$

where $|a_1(\tau)| \leq d_1$, $|a_2(\tau)| \leq d_2$, $|x_n(\varphi_1(\tau))| \leq d_3$, $|x_n(\varphi_2(\tau))| \leq d_4$, $|g_1(\tau)| \leq d_5$, and $|g_2(\tau)| \leq d_6$ for $\tau \in D_\varepsilon$. The previous estimate yields that $\{U_n\}$ is equibounded in $C(D_\varepsilon)$.

Next, for arbitrary $\tau_1, \tau_2 \in D_\varepsilon$ with $\tau_1 < \tau_2$, we get

$$\begin{aligned}
 & |U_n(\tau_2) - U_n(\tau_1)| \\
 = & \left| \left(g_1(\tau_2) + g_3(\tau_2) \cdot (Gx_n)(\varphi_1(\tau_2)) \right) \left(g_2(\tau_2) + \frac{1}{\Gamma(\alpha)} \int_0^{\tau_2} \frac{u(s, x_n(\varphi_2(s)))}{(\tau_2 - s)^{1-\alpha}} ds \right) \right. \\
 & - \left. \left(g_1(\tau_1) + g_3(\tau_1) \cdot (Gx_n)(\varphi_1(\tau_1)) \right) \left(g_2(\tau_1) + \frac{1}{\Gamma(\alpha)} \int_0^{\tau_1} \frac{u(s, x_n(\varphi_2(s)))}{(\tau_1 - s)^{1-\alpha}} ds \right) \right| \\
 \leq & \left| g_1(\tau_2)g_2(\tau_2) - g_1(\tau_1)g_2(\tau_1) \right| + \left| \frac{g_3(\tau_2)(Gx_n)(\varphi_1(\tau_2))}{\Gamma(\alpha)} \int_0^{\tau_2} \frac{u(s, x_n(\varphi_2(s)))}{(\tau_2 - s)^{1-\alpha}} ds \right. \\
 & \left. - \frac{g_3(\tau_1)(Gx_n)(\varphi_1(\tau_1))}{\Gamma(\alpha)} \int_0^{\tau_1} \frac{u(s, x_n(\varphi_2(s)))}{(\tau_1 - s)^{1-\alpha}} ds \right| \\
 & + \left| \frac{g_1(\tau_2)}{\Gamma(\alpha)} \int_0^{\tau_2} \frac{u(s, x_n(\varphi_2(s)))}{(\tau_2 - s)^{1-\alpha}} ds - \frac{g_1(\tau_1)}{\Gamma(\alpha)} \int_0^{\tau_1} \frac{u(s, x_n(\varphi_2(s)))}{(\tau_1 - s)^{1-\alpha}} ds \right| \\
 & + \left| g_2(\tau_2)g_3(\tau_2) \cdot (Gx_n)(\varphi_1(\tau_2)) - g_2(\tau_1)g_3(\tau_1) \cdot (Gx_n)(\varphi_1(\tau_1)) \right| \\
 = & I_1 + I_2 + I_3 + I_4. \tag{3.5}
 \end{aligned}$$

First, we estimate I_2 :

$$\begin{aligned}
 I_2 & = \left| \frac{g_3(\tau_2)(Gx_n)(\varphi_1(\tau_2))}{\Gamma(\alpha)} \int_0^{\tau_2} \frac{u(s, x_n(\varphi_2(s)))}{(\tau_2 - s)^{1-\alpha}} ds - \frac{g_3(\tau_1)(Gx_n)(\varphi_1(\tau_1))}{\Gamma(\alpha)} \int_0^{\tau_1} \frac{u(s, x_n(\varphi_2(s)))}{(\tau_1 - s)^{1-\alpha}} ds \right| \\
 & \leq \left| \frac{g_3(\tau_2)(Gx_n)(\varphi_1(\tau_2))}{\Gamma(\alpha)} \int_0^{\tau_2} \frac{u(s, x_n(\varphi_2(s)))}{(\tau_2 - s)^{1-\alpha}} ds - \frac{g_3(\tau_2)(Gx_n)(\varphi_1(\tau_1))}{\Gamma(\alpha)} \int_0^{\tau_1} \frac{u(s, x_n(\varphi_2(s)))}{(\tau_1 - s)^{1-\alpha}} ds \right| \\
 & + \left| \frac{g_3(\tau_2)(Gx_n)(\varphi_1(\tau_1))}{\Gamma(\alpha)} \int_0^{\tau_1} \frac{u(s, x_n(\varphi_2(s)))}{(\tau_1 - s)^{1-\alpha}} ds - \frac{g_3(\tau_1)(Gx_n)(\varphi_1(\tau_1))}{\Gamma(\alpha)} \int_0^{\tau_1} \frac{u(s, x_n(\varphi_2(s)))}{(\tau_1 - s)^{1-\alpha}} ds \right| \\
 & \leq |g_3(\tau_2)| \left| \frac{(Gx_n)(\varphi_1(\tau_2))}{\Gamma(\alpha)} \int_0^{\tau_2} \frac{u(s, x_n(\varphi_2(s)))}{(\tau_2 - s)^{1-\alpha}} ds - \frac{(Gx_n)(\varphi_1(\tau_2))}{\Gamma(\alpha)} \int_0^{\tau_1} \frac{u(s, x_n(\varphi_2(s)))}{(\tau_1 - s)^{1-\alpha}} ds \right. \\
 & \left. + \frac{(Gx_n)(\varphi_1(\tau_2))}{\Gamma(\alpha)} \int_0^{\tau_1} \frac{u(s, x_n(\varphi_2(s)))}{(\tau_1 - s)^{1-\alpha}} ds - \frac{(Gx_n)(\varphi_1(\tau_1))}{\Gamma(\alpha)} \int_0^{\tau_1} \frac{u(s, x_n(\varphi_2(s)))}{(\tau_1 - s)^{1-\alpha}} ds \right| \\
 & + |g_3(\tau_2) - g_3(\tau_1)| \left| \frac{(Gx_n)(\varphi_1(\tau_1))}{\Gamma(\alpha)} \int_0^{\tau_1} \frac{u(s, x_n(\varphi_2(s)))}{(\tau_1 - s)^{1-\alpha}} ds \right| \\
 & \leq M \left| \frac{(Gx_n)(\varphi_1(\tau_2))}{\Gamma(\alpha)} \right| \left| \int_0^{\tau_2} \frac{u(s, x_n(\varphi_2(s)))}{(\tau_2 - s)^{1-\alpha}} ds - \int_0^{\tau_1} \frac{u(s, x_n(\varphi_2(s)))}{(\tau_1 - s)^{1-\alpha}} ds \right| \\
 & + \frac{M}{\Gamma(\alpha)} \left| (Gx_n)(\varphi_1(\tau_2)) - (Gx_n)(\varphi_1(\tau_1)) \right| \left| \int_0^{\tau_1} \frac{u(s, x_n(\varphi_2(s)))}{(\tau_1 - s)^{1-\alpha}} ds \right| \\
 & + \omega^T(g_3, |\tau_2 - \tau_1|)(d_1 + b_1 d_3^{\frac{1}{3}}) \frac{(d_2 + b_2 d_4^{\frac{1}{2}}) T^\alpha}{\Gamma(\alpha + 1)}
 \end{aligned}$$

$$\begin{aligned}
 &\leq \frac{M(d_1 + b_1 d_3^{\frac{1}{q}})}{\Gamma(\alpha)} \left| \int_0^{\tau_2} \frac{u(s, x_n(\varphi_2(s)))}{(\tau_2 - s)^{1-\alpha}} ds - \int_0^{\tau_2} \frac{u(s, x_n(\varphi_2(s)))}{(\tau_1 - s)^{1-\alpha}} ds \right| \\
 &\quad + \frac{M(d_1 + b_1 d_3^{\frac{1}{q}})}{\Gamma(\alpha)} \left| \int_0^{\tau_2} \frac{u(s, x_n(\varphi_2(s)))}{(\tau_1 - s)^{1-\alpha}} ds - \int_0^{\tau_1} \frac{u(s, x_n(\varphi_2(s)))}{(\tau_1 - s)^{1-\alpha}} ds \right| \\
 &\quad + M\omega^T(G, |\tau_2 - \tau_1|) \frac{(d_2 + b_2 d_4^{\frac{1}{p}})T^\alpha}{\Gamma(\alpha + 1)} + \omega^T(g_3, |\tau_2 - \tau_1|)(d_1 + b_1 d_3^{\frac{1}{q}}) \frac{(d_2 + b_2 d_4^{\frac{1}{p}})T^\alpha}{\Gamma(\alpha + 1)} \\
 &\leq \frac{M(d_1 + b_1 d_3^{\frac{1}{q}})(d_2 + b_2 d_4^{\frac{1}{p}})}{\Gamma(\alpha)} \left(\int_0^{\tau_2} |(\tau_2 - s)^{\alpha-1} - (\tau_1 - s)^{\alpha-1}| ds + \int_{\tau_1}^{\tau_2} (\tau_1 - s)^{\alpha-1} ds \right) \\
 &\quad + \frac{(d_2 + b_2 d_4^{\frac{1}{p}})T^\alpha}{\Gamma(\alpha + 1)} \left[M\omega^T(G, |\tau_2 - \tau_1|) + (d_1 + b_1 d_3^{\frac{1}{q}})\omega^T(g_3, |\tau_2 - \tau_1|) \right] \\
 &\leq \frac{M(d_1 + b_1 d_3^{\frac{1}{q}})(d_2 + b_2 d_4^{\frac{1}{p}})}{\alpha\Gamma(\alpha)} \left(-(\tau_2 - s)^\alpha \Big|_0^{\tau_2} + (\tau_1 - s)^\alpha \Big|_0^{\tau_2} - (\tau_1 - s)^\alpha \Big|_{\tau_1}^{\tau_2} \right) \\
 &\quad + \frac{(d_2 + b_2 d_4^{\frac{1}{p}})T^\alpha}{\Gamma(\alpha + 1)} \left[M\omega^T(G, |\tau_2 - \tau_1|) + (d_1 + b_1 d_3^{\frac{1}{q}})\omega^T(g_3, |\tau_2 - \tau_1|) \right] \\
 &\leq \frac{(d_2 + b_2 d_4^{\frac{1}{p}})}{\Gamma(\alpha + 1)} \left[M(d_1 + b_1 d_3^{\frac{1}{q}})(\tau_2^\alpha - \tau_1^\alpha) + MT^\alpha \omega^T(G, |\tau_2 - \tau_1|) \right. \\
 &\quad \left. + T^\alpha (d_1 + b_1 d_3^{\frac{1}{q}})\omega^T(g_3, |\tau_2 - \tau_1|) \right].
 \end{aligned}$$

By mean value theorem: $\exists z, \tau_1 < z < \tau_2$ s.t. $\tau_2^\alpha - \tau_1^\alpha = \alpha z^{\alpha-1}(\tau_2 - \tau_1) \leq (\tau_2 - \tau_1)$, then we have

$$I_2 \leq \frac{(d_2 + b_2 d_4^{\frac{1}{p}})}{\Gamma(\alpha + 1)} \left[M(d_1 + b_1 d_3^{\frac{1}{q}})(\tau_2 - \tau_1) + MT^\alpha \omega^T(G, |\tau_2 - \tau_1|) + T^\alpha (d_1 + b_1 d_3^{\frac{1}{q}})\omega^T(g_3, |\tau_2 - \tau_1|) \right]. \quad (3.6)$$

To evaluate I_1 , we have

$$I_1 = \left| g_1(\tau_2)g_2(\tau_2) - g_1(\tau_1)g_2(\tau_1) \right| \leq d_5 \omega^T(g_2, |\tau_2 - \tau_1|) + d_6 \omega^T(g_1, |\tau_2 - \tau_1|). \quad (3.7)$$

To evaluate I_3 , we have

$$\begin{aligned}
 I_3 &= \left| \frac{g_1(\tau_2)}{\Gamma(\alpha)} \int_0^{\tau_2} \frac{u(s, x_n(\varphi_2(s)))}{(\tau_2 - s)^{1-\alpha}} ds - \frac{g_1(\tau_1)}{\Gamma(\alpha)} \int_0^{\tau_1} \frac{u(s, x_n(\varphi_2(s)))}{(\tau_1 - s)^{1-\alpha}} ds \right| \\
 &\leq \frac{d_5 \left(d_2 + b_2 d_4^{\frac{1}{p}} \right)}{\Gamma(\alpha + 1)} |\tau_2 - \tau_1| + \frac{\left(d_2 + b_2 d_4^{\frac{1}{p}} \right) T^\alpha}{\Gamma(\alpha + 1)} \omega^T(g_1, |\tau_2 - \tau_1|)
 \end{aligned} \quad (3.8)$$

and to estimate I_4 , we have

$$\begin{aligned} I_4 &= \left| g_2(\tau_2)g_3(\tau_2) \cdot (Gx_n)(\varphi_1(\tau_2)) - g_2(\tau_1)g_3(\tau_1) \cdot (Gx_n)(\varphi_1(\tau_1)) \right| \\ &\leq d_6 M \omega^T(G, |\tau_2 - \tau_1|) + d_6(d_1 + b_2 d_3^{\frac{1}{q}}) \omega^T(g_3, |\tau_2 - \tau_1|) \\ &\quad + M(d_1 + b_2 d_3^{\frac{1}{q}}) \omega^T(g_2, |\tau_2 - \tau_1|), \end{aligned} \tag{3.9}$$

where $\omega^T(g_i, \cdot)$, $\omega^T(G, \cdot)$ denotes the modulus continuity of the functions $g_i (i = 1, 2, 3)$ and G on the sets D_ε and $D_\varepsilon \times \mathbb{R}$, respectively. By substituting from (3.6), (3.7), (3.8), and (3.9) in (3.5), we have an inequality which has been gotten since $Q \subset B_r$.

From the fact $\gamma(\{x_n\}) \leq \gamma(Q) = 0$, we deduce that, (3.5) tends to zero as $(\tau_2 - \tau_1) \rightarrow 0$ independently of x_n . We have $\{U_n\}$ is equicontinuous in $C(D_\varepsilon)$. Set

$$Y = \sup\{|U_n| : \tau \in D_\varepsilon, n \in \mathbb{N}\}.$$

Obviously, Y is finite and according to the choice of the set D_ε we conclude that the functions $u|_{D_\varepsilon \times [-Y, Y]}$, $h|_{D_\varepsilon \times [-Y, Y]}$, $G|_{D_\varepsilon \times [-Y, Y]}$ are uniformly continuous. So $\{Hx_n\} = \{F_{\varphi_3, h}(x_n) + U_n\}$ is equicontinuous and equibounded in $C(D_\varepsilon)$. Then, by Ascoli–Arzela theorem [18], we deduce that $\{Hx_n\}$ is a relatively compact set in $C(D_\varepsilon)$.

We note that the above thinking is independent of the choice of ε . Then, we shall reconstruct a sequence of closed sets $D_l \subset [0, T]$ s.t. $meas(D_l^c) \rightarrow 0$ as $l \rightarrow \infty$ and s.t. $\{Hx_n\}$ is relatively compact in each space $C(D_l)$.

By utilizing the weak compact set $H(Q)$, we select a number $\delta > 0$ s.t. for every closed set $D_\delta \subset [0, T]$ with $meas(D_\delta^c) \leq \delta$, we have

$$\int_{D_\delta^c} e^{-N\tau} |(Hx)(\tau)| d\tau \leq \frac{\varepsilon}{4}, \tag{3.10}$$

for any $x \in Q$. Since, the sequence $\{Hx_n\}$ is a Cauchy sequence in the spaces $C(D_l)$, $l = 1, 2, \dots$, we can select a number l_0 s.t. $meas(D_{l_0}^c) \leq \delta$ and $meas(D_{l_0}) > 0$, and for any numbers $n, m \geq l_0$, we have

$$e^{-N\tau} |(Hx_n)(\tau) - (Hx_m)(\tau)| \leq \frac{\varepsilon}{4meas(D_{l_0})} \tag{3.11}$$

for any $\tau \in D_{l_0}$. Then, by using (3.4), (3.10), and (3.11) we obtain

$$\begin{aligned} &\int_0^\infty e^{-N\tau} |(Hx_n)(\tau) - (Hx_m)(\tau)| d\tau = \int_T^\infty e^{-N\tau} |(Hx_n)(\tau) - (Hx_m)(\tau)| d\tau \\ &+ \int_{D_{l_0}} e^{-N\tau} |(Hx_n)(\tau) - (Hx_m)(\tau)| d\tau + \int_{D_{l_0}^c} e^{-N\tau} |(Hx_n)(\tau) - (Hx_m)(\tau)| d\tau \leq \varepsilon, \end{aligned}$$

which gives that $\{Hx_n\}$ is a Cauchy sequence in L_1^N . Hence, we obtain that the set (HQ) is relatively strongly compact in L_1^N .

Step 6. Let $Q_0 = Conv(H(Q))$ and by Mazur theorem, we have that Q_0 is compact in L_1^N . Further, the operator $H : Q_0 \rightarrow Q_0$ is continuous. Then, by applying Theorem 2.2, this fulfills the proof. \square

Remark 3.3 The results mentioned in Theorem 3.2 can be applied in the spaces $L_p^N(\mathbb{R}^+)$, ($p > 1$) or in Orlicz spaces L_φ under appropriate set of assumptions guarantee continuity and acting conditions of the studied problems (cf. [14]).

Remark 3.4 One can easily check that the operators $Gx = 1$, $Gx = x$ and $Gx = f(t, x)$ are examples of the operator Gx , which satisfy assumptions (ii) and (iii) of Theorem 3.2. Moreover, Gx shall take the form of linear, nonlinear, Hammerstein, Urysohn, or fraction integral operators.

Now, we will tackle the uniqueness of the solutions.

Theorem 3.5 Let assumptions of Theorem 3.2 are fulfilled, but replace assumption (iii) by the conditions:

(vii) $\exists b_i \geq 0$, ($i = 1, 3, 3$) and the functions $a_1 \in L_q^N$, $a_2 \in L_p^N$ and $a_3 \in L_1^N$ s.t.

$$|G(0)(\tau)| \leq a_1(\tau), \quad |u(\tau, 0)| \leq a_2(\tau), \quad |h(\tau, 0)| \leq a_3(\tau)$$

$$|(Gx)(\tau) - (Gy)(\tau)| \leq b_1|x - y|^{\frac{1}{q}}, \quad |u(\tau, x) - u(\tau, y)| \leq b_2|x - y|^{\frac{1}{p}},$$

$$\text{and } |h(\tau, x) - h(\tau, y)| \leq b_3|x - y|$$

for $x, y \in Q$, where Q is as in Theorem 3.2.

(viii) If for any constant $\mathbb{L} \geq 0$, we have

$$\begin{aligned} \mathbb{L} \leq & \frac{b_3}{M_3}\mathbb{L} + \frac{b_2p^\alpha \|g_1\|_{L_q^N}}{N^\alpha M_2^{\frac{1}{p}}}\mathbb{L}^{\frac{1}{p}} + \frac{Mb_1\|g_2\|_{L_p^N}}{M_1^{\frac{1}{q}}}\mathbb{L}^{\frac{1}{q}} + \frac{p^\alpha Mb_2}{N^\alpha M_2^{\frac{1}{p}}}\left(\|a_1\|_{L_q^N} + \frac{b_1}{M_1^{\frac{1}{q}}}r^{\frac{1}{q}}\right)\mathbb{L}^{\frac{1}{p}} \\ & + \frac{p^\alpha Mb_1}{N^\alpha M_1^{\frac{1}{q}}}\left(\|a_2\|_{L_p^N} + \frac{b_2}{M_2^{\frac{1}{p}}}r^{\frac{1}{p}}\right)\mathbb{L}^{\frac{1}{q}}, \text{ then } \mathbb{L} = 0, \end{aligned}$$

where r is defined in Equation (3.1), then (1.1) has a unique integrable solution $x \in L_1^N$ in Q .

Proof From assumption (vii), we derive

$$\begin{aligned} \left| |u(\tau, x)| - |u(\tau, 0)| \right| & \leq |u(\tau, x) - u(\tau, 0)| \leq b_2|x|^{\frac{1}{p}} \\ \Rightarrow |u(\tau, x)| & \leq |u(\tau, 0)| + b_2|x|^{\frac{1}{p}} \leq a_2(\tau) + b_2|x|^{\frac{1}{p}}. \end{aligned}$$

Similarly, we have $|h(\tau, x)| \leq a_3(\tau) + b_3|x|$ and $|Gx(\tau)| \leq a_1(\tau) + b_1|x|^{\frac{1}{q}}$. Then by Theorem 3.2, we have (1.1) has at least one solution $x \in L_1^N$ on \mathbb{R}^+ .

Next, assume that let x and y be any solutions of Equation (1.1). Thus we have

$$\begin{aligned}
 |x(\tau) - y(\tau)| &= \left| h(\tau, x(\varphi_3(\tau))) + \left(g_1(\tau) + g_3(\tau)(Gx(\varphi_1(\tau))) \right) \left(g_2(\tau) + \frac{1}{\Gamma(\alpha)} \int_0^\tau \frac{u(s, x(\varphi_2(s)))}{(\tau - s)^{1-\alpha}} ds \right) \right. \\
 &\quad \left. - h(\tau, y(\varphi_3(\tau))) - \left(g_1(\tau) + g_3(\tau)(Gy(\varphi_1(\tau))) \right) \cdot \left(g_2(\tau) + \frac{1}{\Gamma(\alpha)} \int_0^\tau \frac{u(s, y(\varphi_2(s)))}{(\tau - s)^{1-\alpha}} ds \right) \right| \\
 &\leq |h(\tau, x(\varphi_3(\tau))) - h(\tau, y(\varphi_3(\tau)))| + \frac{|g_1(\tau)|}{\Gamma(\alpha)} \int_0^\tau \frac{|u(s, x(\varphi_2(s))) - u(s, y(\varphi_2(s)))|}{(\tau - s)^{1-\alpha}} ds \\
 &\quad + |g_2(\tau)| |g_3(\tau)| |Gx(\varphi_1(\tau)) - Gy(\varphi_1(\tau))| \\
 &\quad + \frac{|g_3(\tau)(Gx(\varphi_1(\tau)))|}{\Gamma(\alpha)} \int_0^\tau \frac{|u(s, x(\varphi_2(s))) - u(s, y(\varphi_2(s)))|}{(\tau - s)^{1-\alpha}} ds \\
 &\quad + \frac{|g_3(\tau)|}{\Gamma(\alpha)} \left| Gx(\varphi_1(\tau)) - Gy(\varphi_1(\tau)) \right| \int_0^\tau \frac{|u(s, y(\varphi_2(s)))|}{(\tau - s)^{1-\alpha}} ds \\
 &\leq b_3 |x(\varphi_3(\tau)) - y(\varphi_3(\tau))| \\
 &\quad + \frac{|g_1(\tau)|}{\Gamma(\alpha)} \int_0^\tau \frac{b_2 |x(\varphi_2(s)) - y(\varphi_2(s))|^{\frac{1}{p}}}{(\tau - s)^{1-\alpha}} ds + b_1 M |g_2(\tau)| |x(\varphi_1(\tau)) - y(\varphi_1(\tau))|^{\frac{1}{q}} \\
 &\quad + \frac{M}{\Gamma(\alpha)} \left(a_1(\tau) + b_1 |x(\varphi_1(\tau))|^{\frac{1}{q}} \right) \cdot \int_0^\tau \frac{b_2 |x(\varphi_2(s)) - y(\varphi_2(s))|^{\frac{1}{p}}}{(\tau - s)^{1-\alpha}} ds \\
 &\quad + \frac{b_1 M}{\Gamma(\alpha)} \left| x(\varphi_1(\tau)) - y(\varphi_1(\tau)) \right|^{\frac{1}{q}} \int_0^\tau \frac{a_2(s) + b_2 |y(\varphi_2(s))|^{\frac{1}{p}}}{(\tau - s)^{1-\alpha}} ds.
 \end{aligned}$$

Therefore,

$$\begin{aligned}
 \|x - y\|_{L_1^N} &\leq b_3 \|x(\varphi_3) - y(\varphi_3)\|_{L_1^N} + \frac{b_2 \|g_1\|_{L_q^N}}{\Gamma(\alpha)} \left\| \int_0^\tau \frac{|x(\varphi_2(s)) - y(\varphi_2(s))|^{\frac{1}{p}}}{(\tau - s)^{1-\alpha}} ds \right\|_{L_p^N} \\
 &\quad + M b_1 \|g_2\|_{L_p^N} \left\| |x(\varphi_1) - y(\varphi_1)|^{\frac{1}{q}} \right\|_{L_q^N} \\
 &\quad + \frac{M b_2}{\Gamma(\alpha)} \left\| a_1 + b_1 |x(\varphi_1)|^{\frac{1}{q}} \right\|_{L_q^N} \cdot \left\| \int_0^\tau \frac{|x(\varphi_2(s)) - y(\varphi_2(s))|^{\frac{1}{p}}}{(\tau - s)^{1-\alpha}} ds \right\|_{L_p^N} \\
 &\quad + \frac{M b_1}{\Gamma(\alpha)} \left\| |x(\varphi_1) - y(\varphi_1)|^{\frac{1}{q}} \right\|_{L_q^N} \cdot \left\| \int_0^\tau \frac{a_2(s) + b_2 |y(\varphi_2(s))|^{\frac{1}{p}}}{(\tau - s)^{1-\alpha}} ds \right\|_{L_p^N} \\
 &\leq \frac{b_3}{M_3} \|x - y\|_{L_1^N} + \frac{p^\alpha b_2 \|g_1\|_{L_q^N}}{N^\alpha} \left\| \left(x(\varphi_2) - y(\varphi_2) \right)^{\frac{1}{p}} \right\|_{L_p^N} + \frac{M b_1 \|g_2\|_{L_p^N}}{M_1^{\frac{1}{q}}} \|x - y\|_{L_1^N}^{\frac{1}{q}} \\
 &\quad + \frac{p^\alpha M b_2}{N^\alpha} \left(\|a_1\|_{L_q^N} + \frac{b_1}{M_1^{\frac{1}{q}}} \|x\|_{L_1^N}^{\frac{1}{q}} \right) \left\| \left(x(\varphi_2) - y(\varphi_2) \right)^{\frac{1}{p}} \right\|_{L_p^N}
 \end{aligned}$$

$$\begin{aligned}
 & + \frac{p^\alpha M b_1}{N^\alpha M_1^{\frac{1}{q}}} \|x - y\|_{L_1^N}^{\frac{1}{q}} \cdot \left\| a_2 + b_2 |y(\varphi_2)|^{\frac{1}{p}} \right\|_{L_p^N} \\
 \leq & \frac{b_3}{M_3} \|x - y\|_{L_1^N} + \frac{b_2 p^\alpha \|g_1\|_{L_q^N}}{N^\alpha M_2^{\frac{1}{p}}} \|x - y\|_{L_1^N}^{\frac{1}{p}} + \frac{M b_1 \|g_2\|_{L_p^N}}{M_1^{\frac{1}{q}}} \|x - y\|_{L_1^N}^{\frac{1}{q}} \\
 & + \frac{p^\alpha M b_2}{N^\alpha M_2^{\frac{1}{p}}} \left(\|a_1\|_{L_q^N} + \frac{b_1}{M_1^{\frac{1}{q}}} \|x\|_{L_1^N}^{\frac{1}{q}} \right) \|x - y\|_{L_1^N}^{\frac{1}{p}} + \frac{p^\alpha M b_1}{N^\alpha M_1^{\frac{1}{q}}} \left(\|a_2\|_{L_p^N} + \frac{b_2}{M_2^{\frac{1}{p}}} \|y\|_{L_1^N}^{\frac{1}{p}} \right) \|x - y\|_{L_1^N}^{\frac{1}{q}} \\
 \leq & \frac{b_3}{M_3} \|x - y\|_{L_1^N} + \frac{b_2 p^\alpha \|g_1\|_{L_q^N}}{N^\alpha M_2^{\frac{1}{p}}} \|x - y\|_{L_1^N}^{\frac{1}{p}} + \frac{M b_1 \|g_2\|_{L_p^N}}{M_1^{\frac{1}{q}}} \|x - y\|_{L_1^N}^{\frac{1}{q}} \\
 & + \frac{p^\alpha M b_2}{N^\alpha M_2^{\frac{1}{p}}} \left(\|a_1\|_{L_q^N} + \frac{b_1}{M_1^{\frac{1}{q}}} r^{\frac{1}{q}} \right) \|x - y\|_{L_1^N}^{\frac{1}{p}} + \frac{p^\alpha M b_1}{N^\alpha M_1^{\frac{1}{q}}} \left(\|a_2\|_{L_p^N} + \frac{b_2}{M_2^{\frac{1}{p}}} r^{\frac{1}{p}} \right) \|x - y\|_{L_1^N}^{\frac{1}{q}}.
 \end{aligned}$$

The above inequality with assumption (viii) implies that $x = y$ (a.e.), and that fulfills the proof. \square

3.1. Example

We construct an example to indicate the validity of our outcomes.

Example 3.6 For $\tau \in [0, \infty)$, consider the equation

$$x(\tau) = \tau^3 + \frac{x(\sin(3\tau))}{e^{N^2}} + \left(e^{-\tau} + e^{-\tau^3} \left(\tau^{\frac{5}{2}} + \frac{\sqrt{x\left(\frac{\tau}{4+\tau}\right)}}{e^{N^2}} \right) \right) \left(e^\tau + \int_0^\tau \frac{(\tau-s)^{-\frac{1}{2}}}{\Gamma(\frac{1}{2})} \left(e^{5s} + \frac{\sqrt{x\left(\frac{2s}{2+s}\right)}}{2 + \left(x\left(\frac{2s}{2+s}\right)\right)^2} \right) ds \right). \quad (3.12)$$

For $p = q = 2$, we can see that

1. $g_1(\tau) = e^{-\tau} \in L_2^N$ and $\|g_1\|_{L_2^N} = \frac{1}{\sqrt{N+2}}$.
2. $g_2(\tau) = e^\tau \in L_2^N$ and $\|g_2\|_{L_2^N} = \frac{1}{\sqrt{N-2}}$.
3. $g_3(\tau) = e^{-\tau^3}$ and $M = \sup_{0 \leq \tau < \infty} |g_3(\tau)| \leq 1$.
4. $(Gx)(\tau) = \tau^{\frac{5}{2}} + \frac{1}{e^{N^2}} \sqrt{x}$, then $a_1(\tau) = \tau^{\frac{5}{2}}$, $b_1 = \frac{1}{e^{N^2}}$ with $\|a_1\|_{L_2^N} = \frac{\sqrt{120}}{N^3}$.
5. $u(\tau, x) = e^{5\tau} + \frac{\sqrt{x}}{2+x^2}$, then $a_2(\tau) = e^{5\tau}$, $b_2 = \frac{1}{2}$ with $\|a_2\|_{L_2^N} = \frac{1}{\sqrt{N-10}}$.
6. $h(\tau, x) = \tau^3 + \frac{1}{e^{N^2}} x$, then $a_3(\tau) = \tau^3$, $b_3 = \frac{1}{e^{N^2}}$ with $\|a_3\|_{L_1^N} = \frac{6}{N^4}$.
7. $\varphi_1(\tau) = \frac{\tau}{4+\tau}$, then $\varphi_1'(\tau) = \frac{4}{(4+\tau)^2} \geq \frac{1}{16} = M_1$ and $\varphi_1^{-1}(\tau) = \frac{4\tau}{1-\tau} \geq \tau$.
8. $\varphi_2(\tau) = \frac{2\tau}{2+\tau}$, then $\varphi_2'(\tau) = \frac{4}{(2+\tau)^2} \geq \frac{1}{4} = M_2$ and $\varphi_2^{-1}(\tau) = \frac{2\tau}{2-\tau} \geq \tau$.
9. $\varphi_3(\tau) = \sin(3\tau)$, then $\varphi_3'(\tau) = 3 \cos(3\tau) \geq 2 = M_3$ and $\varphi_3^{-1}(\tau) = \frac{\sin^{-1} \tau}{3} \geq \tau$.

10. For $N > 128$, we have

$$W = e^{N^2} \left(\frac{b_3}{M_3} + \frac{Mb_1b_2p^\alpha}{N^\alpha M_1^{\frac{1}{q}} M_2^{\frac{1}{p}}} \right) \leq \left(\frac{1}{2} + 4\sqrt{\frac{2}{N}} \right) < 1.$$

11. Let $r = 1$, and note that

$$\begin{aligned} & \|a_3\|_{L_1^N} + \left(\|g_1\|_{L_2^N} + M\|a_1\|_{L_2^N} \right) \left(\|g_2\|_{L_2^N} + \sqrt{\frac{2}{N}}\|a_2\|_{L_2^N} + b_2\sqrt{\frac{2}{NM_2}} \right) \\ & + \left(\|g_2\|_{L_2^N} + \sqrt{\frac{2}{N}}\|a_2\|_{L_2^N} \right) \frac{Mb_1}{\sqrt{M_1}} + \left(\frac{b_3}{M_3} + Mb_1b_2\sqrt{\frac{2}{NM_1M_2}} \right) \\ & \leq \frac{\sqrt{6}}{N^4} + \left(\frac{1}{\sqrt{N+2}} + \frac{\sqrt{120}}{N^3} \right) \left(\frac{1}{\sqrt{N-2}} + \sqrt{\frac{2}{N}} + \sqrt{\frac{2}{N(N-10)}} \right) \\ & + \frac{4}{e^{N^2}} \left(\frac{1}{\sqrt{N-2}} + \sqrt{\frac{2}{N(N-10)}} \right) + \frac{e^{-N^2}}{2} \left(1 + 8\sqrt{\frac{2}{N}} \right) < 1. \end{aligned}$$

Hence, due to Theorem 3.2, Equation (3.12) has at least one integrable solution $x \in L_1^N$ on \mathbb{R}^+ .

Moreover, we have

1. $|G(0)(\tau)| = \tau^{\frac{5}{2}}$ and $|G(x) - G(y)| \leq \frac{1}{e^{N^2}}|x - y|^{\frac{1}{2}}$.
2. $|u(\tau, 0)| = e^{5\tau}$ and $|u(\tau, x) - u(\tau, y)| \leq \frac{1}{2}|x - y|^{\frac{1}{2}}$.
3. $|h(\tau, 0)| = \tau^3$ and $|h(\tau, x) - h(\tau, y)| \leq \frac{1}{e^{N^2}}|x - y|$.
4. Assumption (viii) is satisfied for $\mathbb{L} = \|x - y\|_{L_1^N} = 0$, i.e. $x = y$ (a.e.).

Hence, due to Theorem 3.5, Equation (3.12) has a unique integrable solution $x \in L_1^N$.

4. Conclusion

This article dealt with the integral equations of the Gripenberg’s type of fractional order with a perturbation term (1.1) in weighted Lebesgue spaces $L_1^N, N > 0$ on $\mathbb{R}^+ = [0, \infty)$. With the aid of Schauder’s fixed point hypothesis and appropriate measure of noncompactness, we proved that considered problem has at least one integrable solution in the considered spaces under a general set of assumptions. Indeed, two new theorems, Theorems 3.2 and 3.5, are proved s.t. Equation (1.1) has at least one integrable solution and a unique integrable solution $x \in L_1^N$, respectively. At the end, an example is constructed to indicate the validity of these theorems. The results here extend and generalize some former results in the available literature.

References

- [1] Abdeldaim A. On some new Gronwall-Bellman-Ou-Iang type integral inequalities to study certain epidemic models. Journal of Integral Equations and Applications 2012; 24 (2): 149-166.

- [2] Aghajani A, O'Regan D, Shole Haghghi A. Measure of noncompactness on $L_p(\mathbb{R}^N)$ and applications. *Cubo. A Mathematical Journal* 2015; 17 (1): 85-97.
- [3] Awad H, Darwish M, Metwali M. On a cubic integral equation of Urysohn type with linear perturbation of second kind. *Journal of Mathematics and Applications* 2018; 41: 29-38.
- [4] Appell J, Zabrejko PP. *Nonlinear Superposition Operators*. Cambridge: Cambridge Tracts in Mathematics; 95, Cambridge University Press, 1990.
- [5] Banaś J, Chlebowicz A. On solutions of an infinite system of nonlinear integral equations on the real half-axis. *Banach Journal of Mathematical Analysis* 2019; 13 (4): 944-968.
- [6] Banaś J, Goebel K. *Measures of Noncompactness in Banach Spaces*. New York - Basel: Lecture Notes in Mathematics 60, M. Dekker, 1980.
- [7] Banaś J, Knap Z. Measures of weak noncompactness and nonlinear integral equations of convolution type. *Journal of Mathematical Analysis and Applications* 1990; 146 (2): 353-362.
- [8] Bellour A, Bousselsal M, Taoudi MA. Integrable solutions of a nonlinear integral equation related to some epidemic models. *Glasnik Matematički* 2014; 49 (2): 395-406.
- [9] Bohner M, Tunç O. Qualitative analysis of integro-differential equations with variable retardation. *Discrete and Continuous Dynamical Systems-B*, 2021; doi: 10.3934/dcdsb.2021059
- [10] Bohner M, Tunç O, Tunç C. Qualitative analysis of Caputo fractional integro-differential equations with constant delays. *Computational and Applied Mathematics* 2021; 40, 214. doi: 10.1007/s40314-021-01595-3
- [11] Boulfoul B, Bellour A, Djebali S. Solvability of nonlinear integral equations of product type. *Electronic Journal of Differential Equations* 2018; 2018 (19): 1-20.
- [12] Brestovanská E, Medved M. Fixed point theorems of the Banach and Krasnosel'skii type for mappings on m -tuple Cartesian product of Banach algebras and systems of generalized Gripenberg's equations. *Acta Universitatis Palackianae Olomucensis. Facultas Rerum Naturalium. Mathematica* 2012; 51 (2): 27-39.
- [13] Cichoń M, Metwali M. Existence of monotonic L_φ -solutions for quadratic Volterra functional-integral equations. *Electronic Journal of Qualitative Theory of Differential Equations* 2015; 13: 1-16.
- [14] Cichoń M, Metwali M. On a fixed point theorem for the product of operators. *Journal of Fixed Point Theory and Applications* 2016; 18 (4): 753-770.
- [15] Darwish M, Metwali M, O'Regan D. Unique solvability of fractional quadratic nonlinear integral equations. *Differential Equations and Applications* 2021; 13 (1): 1-13.
- [16] Deep A, Deepmala, Tunç C. On the existence of solutions of some non-linear functional integral equations in Banach algebra with applications. *Arab Journal of Basic and Applied Sciences* 2020; 27 (1): 279-286.
- [17] Deep A, Deepmala, Roshan JR. Solvability for generalized nonlinear functional integral equations in Banach spaces with applications. *Journal of Integral Equations and Applications* 2021; 33 (1): 19-30. doi: 10.1216/jie.2021.33.19
- [18] Deimling K. *Nonlinear Functional Analysis*. Berlin: Springer-Verlag, 1985.
- [19] Gorenflo R, Vessela S. *Abel Integral Equations*. Berlin-Heidelberg: Lecture Notes in Mathematics; 1461, Springer, 1991.
- [20] Gripenberg G. Periodic solutions of an epidemic model. *Journal of Mathematical Biology* 1980; 10 (3): 271-280.
- [21] Gu H, Zhou Y, Ahmad B, Alsaedi A. Integral solutions of fractional evolution equations with nondense domain. *Electronic Journal of Differential Equations* 2017; 2017 (145): 1-15.
- [22] Karoui A, Jawahdou A, Ben Aouicha H. Weighted L_p -solutions on unbounded intervals of nonlinear integral equations of the Hammerstein and Urysohn types. *Advances in Pure and Applied Mathematics* 2010; 2 (1): 1-22.
- [23] Kilbas AA, Srivastava HM, Trujillo JJ. *Theory and Applications of Fractional Differential Equations*. Elsevier, Amsterdam: 2006.

- [24] Kwapisz J. Wighted norms and Volterra integral equations in L^p spaces. *Journal of Applied Mathematics and Stochastic Analysis* 1991; 4 (2): 161-164.
- [25] Metwali M. On a class of quadratic Urysohn–Hammerstein integral equations of mixed type and initial value problem of fractional order. *Mediterranean Journal of Mathematics* 2016; 13 (5): 2691-2707.
- [26] Metwali M. On some qualitative properties of integrable solutions for Cauchy-type problem of fractional order. *Journal of Mathematics and Applications* 2017; 40: 121-134.
- [27] Metwali M. Solvability of functional quadratic integral equations with perturbation. *Opuscula Mathematica* 2013; 33: 725-739.
- [28] Metwali M. The solvability of functional quadratic Volterra-Urysohn integral equations on the half line. *Fasciculi Mathematici* 2018; 61: 109-123.
- [29] Pachpatte BG. On a new inequality suggested by the study of certain epidemic models. *Journal of Mathematical Analysis and Applications* 1995; 195 (3): 638-644.
- [30] Samko SG, Kilbas AA, Marichev OI. *Fractional Integrals and Derivative. Theory and Applications*. Gordon and Breach Science Publishers, 1993.
- [31] Scorza Dragoni G. Un teorema sulle funzioni continue rispetto ad una e misarubili rispetto ad un'altra variabile. *Rendiconti del Seminario Matematico della Università di Padova* 1948; 17: 102-106.
- [32] Tunç O, Atan Ö, Tunç C, Yao JC. Qualitative analyses of integro-fractional differential equations with Caputo derivatives and retardations via the Lyapunov-Razumikhin method. *Axioms* 2021; 10(2) 58. doi: 10.3390/axioms10020058
- [33] Weng S, Wang F. Existence of weak solutions for a boundary value problem of a second order ordinary differential equation. *Boundary Value Problems* 2018; 9 (2018). doi:10.1186/s13661-018-0929-7.