

Computing forgotten topological index of zero-divisor graphs of commutative rings

Arif GÜRSOY¹ , Necla KIRCALI GÜRSOY² , Alper ÜLKER^{3,*} 

¹Department of Mathematics, Faculty of Science, Ege University, İzmir, Turkey

²Tire Kutsan Vocational School, Ege University, İzmir, Turkey

³Department of Mathematics, Faculty of Science and Letters, Ağrı İbrahim Çeçen University, Ağrı, Turkey

Received: 04.11.2021

Accepted/Published Online: 10.12.2021

Final Version: ..2022

Abstract: The main objective of this paper is to calculate the forgotten topological index of the zero-divisor graph of \mathbb{Z}_n . Let p , q and r be distinct prime numbers. We calculate the forgotten topological index of the ring $\Gamma(\mathbb{Z}_n)$ where $n = p^\alpha, pq, p^2q, p^2q^2, pqr$. Also, we study the forgotten topological index of the product of rings of integers modulo n . We construct a polynomial algorithm to compute the forgotten topological index of $\Gamma(\mathbb{Z}_n)$.

Key words: Topological indices, forgotten topological index, zero-divisor graph, vertex degree, algorithm

1. Introduction and preliminaries

Zero-divisor graphs of commutative rings entered the area of algebraic combinatorics by the work of I. Beck [9]. His definition of zero-divisor graph has a vertex set on R and any two elements $x, y \in R$ are adjacent whenever $xy = 0$. This definition of a zero-divisor graph of a commutative ring was later modified by Anderson and Livingston in [5]. After the introduction of zero-divisor graphs, different types of graphs related to commutative rings emerged such as annihilating-ideal graphs, comaximal graphs, total graphs [1, 4, 6, 35, 37, 39–41, 43]. In recent years, many works have been done on topological indices of graphs. Indeed, these works include zero-divisor graphs of commutative rings. There are different types of topological indices such as distance-based and degree-based. One of the most well-studied distance-based index which is called Wiener index was studied for zero-divisor graphs in [7, 38]. Let $G = (V(G), E(G))$ be a simple and undirected graph. The Wiener index of a graph G is defined as:

$$W(G) = \sum_{u,v \in V(G)} d(u,v)$$

where $d(u,v)$ stands for the distance between the vertices u and v of G .

The first and the second Zagreb indices which are degree-based indices were introduced in [23] and more details about these indices can be found in [24]. These Zagreb indices are shown by M_1 and M_2 and defined as:

$$M_1 = M_1(G) = \sum_{u \in V(G)} d_u^2 \text{ and } M_2 = M_2(G) = \sum_{uv \in E(G)} d_u d_v$$

*Correspondence: aulker@agri.edu.tr

2010 AMS Mathematics Subject Classification: 05C25, 05C09, 13A70, 68R10

respectively. Many researchers have studied some recent results on these indices [10, 25, 28, 44]. Moreover, the Zagreb indices of zero-divisor graphs of $\Gamma(\mathbb{Z}_n)$ were calculated in [38].

Recently, the Sombor index was introduced by Gutman in [26] and defined as:

$$SO(G) = \sum_{uv \in E(G)} \sqrt{d_u^2 + d_v^2}$$

where d_u and d_v is degree of vertices u and v , respectively. This novel topological index is degree-based and has attracted authors and many research have been done about this index [11, 22, 27, 34, 45].

Moreover, the topological indices such as Wiener index, PI index, Zagreb index, and eccentric index measure the characters of drug molecules. Recently, there are many studies to determine these topological indices of special molecular graphs in chemical, nanomaterials and pharmaceutical engineering, etc. [13–20, 42].

Followed by the first and second Zagreb indices, Furtula and Gutman (2015) introduced the forgotten topological index (also called F-index) which was defined as,

$$FT(G) = \sum_{u \in V(G)} d_u^3 = \sum_{uv \in E(G)} (d_u^2 + d_v^2)$$

where d_v is denoted as the degree of vertex v (the number of vertices adjacent to vertex v) [12]. Furtula and Gutman analyzed that predictive ability of the forgotten topological index is almost similar to that of the first Zagreb index and for the eccentric factor and entropy, and both of them obtain correlation coefficients larger than 0.95. This fact implies the reason why the forgotten topological index is useful for testing the chemical and pharmacological properties of drug molecular structures. More recently, Gao et al. presented the forgotten topological index of some significant drug molecular structures [21]. Also, many researchers continue to examine the forgotten topological index with different studies [8, 29–33, 36].

The works mentioned above led us to study the forgotten topological index of $\Gamma(\mathbb{Z}_n)$, $\Gamma(\mathbb{Z}_n \times \mathbb{Z}_m)$, and $\Gamma(\mathbb{Z}_n \times \mathbb{Z}_m \times \mathbb{Z}_r)$. In section 2, we calculate the forgotten topological index of the graph $\Gamma(\mathbb{Z}_n)$ and give an algorithm for this calculation. Finally, in section 3, we examine the forgotten topological index of zero-divisor graphs of products of rings modulo integers n and m .

2. Forgotten topological index of zero-divisor graph of \mathbb{Z}_n

Recently, the zero-divisor graph of the ring \mathbb{Z}_n is a popular research in spectral graph and chemical graph theory. Many researchers have studied in this area. Singh and Bhat have examined adjacency matrix and Wiener index of zero-divisor graph $\Gamma(\mathbb{Z}_n)$ [38]. Later, Asir and Rabikka have studied Wiener index of zero-divisor graph of $\Gamma(\mathbb{Z}_n)$ [7]. Now, we analyze the forgotten topological index of zero-divisor graph $\Gamma(\mathbb{Z}_n)$ in this section.

Theorem 2.1 *Let p be a prime number, then followings hold:*

- (i) *If $p = 2$, then $FT(\Gamma(\mathbb{Z}_{p^2})) = 0$,*
- (ii) *If $p > 2$, then $FT(\Gamma(\mathbb{Z}_{p^2})) = 2(p-2)^2 \binom{p-1}{2}$.*

Proof

- (i) It is clear that \mathbb{Z}_4 has only one nonzero zero-divisor which is 2, then proof follows.

(ii) Since $\mathbb{Z}_{p^2} \cong K_{p-1}$, the proof follows. □

Now we are about to calculate the forgotten topological index of zero-divisor graph for powers of p greater than or equal to 3.

Theorem 2.2 *Let $p > 2$ be a prime number and $\alpha \in \mathbb{N}$ with $\alpha \geq 3$, then the forgotten topological index of $\Gamma(\mathbb{Z}_{p^\alpha})$ is*

$$FT(\Gamma(\mathbb{Z}_{p^\alpha})) = p^\alpha(p-1) \left[\sum_{i=1}^{\lfloor \frac{\alpha-1}{2} \rfloor} \frac{(p^i-1)^3}{p^{i+1}} + \sum_{i=\lceil \frac{\alpha}{2} \rceil}^{\alpha-1} \frac{(p^i-2)^3}{p^{i+1}} \right].$$

Proof We can show the zero-divisors of (\mathbb{Z}_{p^α}) as follows:

$$\begin{aligned} A_1 &= \{px \mid x = 1, \dots, p^{\alpha-1} - 1, p \nmid x\}, \\ &\dots \\ A_i &= \{p^i x \mid x = 1, \dots, p^{\alpha-i} - 1, p \nmid x\}, \\ &\dots \\ A_{\alpha-1} &= \{p^{\alpha-1} x \mid x = 1, \dots, p - 1, p \nmid x\}. \end{aligned}$$

The vertex set of the graph $\Gamma(\mathbb{Z}_{p^\alpha})$ equals $\bigcup_{i=1}^{\alpha-1} A_i$ with $A_i \cap A_j = \emptyset$ for $i \neq j$, where $i, j \in \{1, \dots, \alpha - 1\}$. Besides, $|A_i|$ means the number of vertices of A_i . We calculate the number of vertices of all zero-divisor sets as $|A_1| = p^{\alpha-1} - p^{\alpha-2}$, $|A_2| = p^{\alpha-2} - p^{\alpha-3}$, ..., $|A_i| = p^{\alpha-i-j}(p-1)$, ..., $|A_{\alpha-1}| = p - 1$. Moreover, the degree of each vertex in these zero-divisor sets can be defined as follows

$$d_u = \begin{cases} p-1, & u \in A_1 \\ p^2-1, & u \in A_2 \\ \dots & \\ p^i-1, & i < \alpha/2 \\ p^i-2, & i \geq \alpha/2 \\ \dots & \\ p^{\alpha-2}-2, & u \in A_{\alpha-2} \\ p^{\alpha-1}-2, & u \in A_{\alpha-1} \end{cases}$$

for all $u \in A_i$. The forgotten topological index of $\Gamma(\mathbb{Z}_{p^\alpha})$ can be attained that

$$\begin{aligned}
 \text{FT}(\Gamma(\mathbb{Z}_{p^\alpha})) &= \sum_{u \in V(\Gamma(\mathbb{Z}_{p^\alpha}))} d_u^3 & (2.1) \\
 &= \sum_{i=1}^{\alpha-1} \sum_{u \in A_i} d_u^3 \\
 &= \sum_{i=1}^{\lfloor \frac{\alpha-1}{2} \rfloor} \sum_{u \in A_i} d_u^3 + \sum_{i=\lceil \frac{\alpha}{2} \rceil}^{\alpha-1} \sum_{u \in A_i} d_u^3 \\
 &= \sum_{i=1}^{\lfloor \frac{\alpha-1}{2} \rfloor} |A_i| (p^i - 1)^3 + \sum_{i=\lceil \frac{\alpha}{2} \rceil}^{\alpha-1} |A_i| (p^i - 2)^3 \\
 &= \sum_{i=1}^{\lfloor \frac{\alpha-1}{2} \rfloor} p^{\alpha-i-1} (p-1) (p^i - 1)^3 + \sum_{i=\lceil \frac{\alpha}{2} \rceil}^{\alpha-1} p^{\alpha-i-1} (p-1) (p^i - 2)^3 \\
 &= p^\alpha (p-1) \left[\sum_{i=1}^{\lfloor \frac{\alpha-1}{2} \rfloor} \frac{(p^i - 1)^3}{p^{i+1}} + \sum_{i=\lceil \frac{\alpha}{2} \rceil}^{\alpha-1} \frac{(p^i - 2)^3}{p^{i+1}} \right].
 \end{aligned}$$

□

Example 2.3 For the graph $\Gamma(\mathbb{Z}_{243})$, we have $p^\alpha = 3^5$. Then, $\text{FT}(\Gamma(\mathbb{Z}_{243})) = 1,089,476$, and the set of zero-divisors can be written as follows:

$$\begin{aligned}
 A_1 &= \{3, 6, 12, 15, 21, 24, 30, 33, 39, 42, 48, 51, 57, 60, 66, 69, 75, 78, 84, 87, 93, 96, 102, \\
 &\quad 105, 111, 114, 120, 123, 129, 132, 138, 141, 147, 150, 156, 159, 165, 168, 174, 177, \\
 &\quad 183, 186, 192, 195, 201, 204, 210, 213, 219, 222, 228, 231, 237, 240\}, \\
 A_2 &= \{9, 18, 36, 45, 63, 72, 90, 99, 117, 126, 144, 153, 171, 180, 198, 207, 225, 234\}, \\
 A_3 &= \{27, 54, 108, 135, 189, 216\}, \\
 A_4 &= \{81, 162\}.
 \end{aligned}$$

Additionally, these sets create the graph which can be shown in Figure 1, where $A_1 = \{3x \mid x = 1, \dots, p^{\alpha-1} - 1, p \nmid x\}$, $A_2 = \{9x \mid x = 1, \dots, p^{\alpha-2} - 1, p \nmid x\}$, $A_3 = \{27x \mid x = 1, \dots, p^{\alpha-3} - 1, p \nmid x\}$, and $A_4 = \{81x \mid x = 1, \dots, p^{\alpha-4} - 1, p \nmid x\}$.

In the next theorem, we give the forgotten index of a zero-divisor graph $\Gamma(\mathbb{Z}_{pq})$ for distinct primes p and q .

Theorem 2.4 Let p and q be prime numbers with $p \neq q$. Then the forgotten index of the graph $\Gamma(\mathbb{Z}_{pq})$ is

$$\text{FT}(\Gamma(\mathbb{Z}_{pq})) = (p-1)(q-1) \left[(q-1)^2 + (p-1)^2 \right].$$




Figure 1. The graph $\Gamma(\mathbb{Z}_{243})$

Proof The graph $\Gamma(\mathbb{Z}_{pq})$ is a complete bipartite graph. The bipartitions of $\Gamma(\mathbb{Z}_{pq})$ are $A_1 = \{px \mid x = 1, 2, \dots, q-1\}$ and $A_2 = \{qx \mid x = 1, 2, \dots, p-1\}$. Since $|A_1| = \varphi(\frac{pq}{p}) = q-1$ and $|A_2| = \varphi(\frac{pq}{q}) = p-1$, then the size of this graph is $(p-1)(q-1)$. It follows that

$$\begin{aligned} \text{FT}(\Gamma(\mathbb{Z}_{pq})) &= \sum_{i=1}^2 \sum_{u \in A_i} d_u^3 \\ &= |A_1|(q-1)^3 + |A_2|(p-1)^3 \\ &= (p-1)(q-1)^3 + (q-1)(p-1)^3 \\ &= (p-1)(q-1) \left[(p-1)^2 + (q-1)^2 \right]. \end{aligned}$$

□

Example 2.5 For the zero-divisor graph of \mathbb{Z}_{119} , we attain $n = pq$ where $p = 7$ and $q = 17$. Then, $\text{FT}(\Gamma(\mathbb{Z}_{119})) = 28,032$, and the set of zero-divisors as follows:

$$\begin{aligned} A_1 &= \{7, 14, 21, 28, 35, 42, 49, 56, 63, 70, 77, 84, 91, 98, 105, 112\}, \\ A_2 &= \{17, 34, 51, 68, 85, 102\}. \end{aligned}$$

These sets give rise to the graph depicted in Figure 2, where $A_1 = \{7x \mid x = 1, 2, \dots, q-1\}$ and $A_2 = \{17x \mid x = 1, 2, \dots, p-1\}$.




Figure 2. The graph $\Gamma(\mathbb{Z}_{119})$

Theorem 2.6 Let $\Gamma(\mathbb{Z}_{p^2q})$ be a zero-divisor graph and p and q be distinct prime numbers. If $n = p^2q$, then the forgotten index of $\Gamma(\mathbb{Z}_{p^2q})$ is

$$FT(\Gamma(\mathbb{Z}_{p^2q})) = (p-1)(q-1) \left[(p-1)^3 + p(q-1)^2 + (p-1)^2(p+1)^3 + \frac{(pq-2)^3}{q-1} \right].$$

Proof Since the proper divisors of $n = p^2q$ are p, p^2, q and pq , then the vertex set can be partitioned as $V(\Gamma(\mathbb{Z}_n)) = A_1 \cup A_2 \cup A_3 \cup A_4$ and $A_i \cap A_j = \emptyset$ for $i \neq j$, where $i, j \in \{1, \dots, 4\}$ and

$$A_1 = \{px \mid x = 1, 2, \dots, pq-1, p \nmid x, q \nmid x\},$$

$$A_2 = \{qx \mid x = 1, 2, \dots, p^2-1, p \nmid x\},$$

$$A_3 = \{p^2x \mid x = 1, 2, \dots, q-1\},$$

$$A_4 = \{pqx \mid x = 1, 2, \dots, p-1\}.$$

We can calculate the number of vertices of all zero-divisor sets as $|A_1| = (p-1)(q-1)$, $|A_2| = p(p-1)$, $|A_3| = (q-1)$, and $|A_4| = (p-1)$. Also, the degree of each vertex in these zero-divisor sets can be determined as

$$d_u = \begin{cases} |A_4|, & u \in A_1 \\ |A_3|, & u \in A_2 \\ |A_2| + |A_4|, & u \in A_3 \\ |A_1| + |A_3| + |A_4| - 1, & u \in A_4 \end{cases}.$$

Then, the forgotten topological index of $\Gamma(\mathbb{Z}_{p^2q})$ can be attained that

$$\begin{aligned}
 \text{FT}(\Gamma(\mathbb{Z}_{p^2q})) &= \sum_{u \in V(\Gamma(\mathbb{Z}_{p^2q}))} d_u^3 \\
 &= \sum_{i=1}^4 \sum_{u \in A_i} d_u^3 \\
 &= |A_1||A_4|^3 + |A_2||A_3|^3 + |A_3|(|A_2| + |A_4|)^3 + |A_4|(|A_1| + |A_3| + |A_4| - 1)^3 \\
 &= (p-1)(q-1)(p-1)^3 + p(p-1)(q-1)^3 + (q-1)(p(p-1) + (p-1))^3 \\
 &\quad + (p-1)\left[(p-1)(q-1) + (q-1) + (p-1) - 1\right]^3 \\
 &= (p-1)^4(q-1) + p(p-1)(q-1)^3 + (q-1)(p^2-1)^3 + (p-1)(pq-2)^3 \\
 &= (p-1)\left[(p-1)^3(q-1) + p(q-1)^3 + (p-1)^2(p+1)^3(q-1) + (pq-2)^3\right] \\
 &= (p-1)(q-1)\left[(p-1)^3 + p(q-1)^2 + (p-1)^2(p+1)^3 + \frac{(pq-2)^3}{q-1}\right].
 \end{aligned}$$

□

Example 2.7 For the graph $\Gamma(\mathbb{Z}_{175})$, we have $n = p^2q$ where $p = 5$ and $q = 7$. Then, $\text{FT}(\Gamma(\mathbb{Z}_{175})) = 232,548$, and the set of zero-divisors can be written as follows:

$$A_1 = \{5, 10, 15, 20, 30, 40, 45, 55, 60, 65, 80, 85, 90, 95, 110, 115, 120, 130, 135, 145, 155, 160, 165, 170\},$$

$$A_2 = \{7, 14, 21, 28, 42, 49, 56, 63, 77, 84, 91, 98, 112, 119, 126, 133, 147, 154, 161, 168\},$$

$$A_3 = \{25, 50, 75, 100, 125, 150\},$$

$$A_4 = \{35, 70, 105, 140\}.$$

Moreover, these sets give rise to the graph which can be shown in Figure 3, where $A_1 = \{5x \mid x = 1, 2, \dots, pq - 1, p \nmid x, q \nmid x\}$, $A_2 = \{7x \mid x = 1, 2, \dots, p^2 - 1, p \nmid x\}$, $A_3 = \{25x \mid x = 1, 2, \dots, q - 1\}$, $A_4 = \{35x \mid x = 1, 2, \dots, p - 1\}$.

Theorem 2.8 Let $\Gamma(\mathbb{Z}_{p^2q^2})$ be a zero-divisor graph and p and q be distinct prime numbers. If $n = p^2q^2$, then the forgotten index of $\Gamma(\mathbb{Z}_{p^2q^2})$ is

$$\begin{aligned}
 \text{FT}(\Gamma(\mathbb{Z}_{p^2q^2})) &= (p-1)(q-1)\left[(p-1)^3q + p(q-1)^3 + (p-1)^2(p+1)^3q\right. \\
 &\quad \left.+ p(q-1)^2(q+1)^3 + (pq-2)^3 + \frac{(p^2q-2)^3}{p-1} + \frac{(pq^2-2)^3}{q-1}\right].
 \end{aligned}$$

Proof Since the proper divisors of $n = p^2q^2$ are p, q, p^2, q^2, pq, p^2q and pq^2 , then the vertex set can be




Figure 3. The graph $\Gamma(\mathbb{Z}_{175})$

partitioned into seven subsets as $V(\Gamma(\mathbb{Z}_n)) = \bigcup_{i=1}^7 A_i$ and $A_i \cap A_j = \emptyset$ for $i \neq j$, where $i, j \in \{1, \dots, 7\}$ and

$$\begin{aligned} A_1 &= \{px \mid x = 1, 2, \dots, pq^2 - 1, p \nmid x, q \nmid x\}, \\ A_2 &= \{qx \mid x = 1, 2, \dots, p^2q - 1, p \nmid x, q \nmid x\}, \\ A_3 &= \{p^2x \mid x = 1, 2, \dots, q^2 - 1, q \nmid x\}, \\ A_4 &= \{q^2x \mid x = 1, 2, \dots, p^2 - 1, p \nmid x\}, \\ A_5 &= \{pqx \mid x = 1, 2, \dots, pq - 1, p \nmid x, q \nmid x\}, \\ A_6 &= \{p^2qx \mid x = 1, 2, \dots, q - 1\}, \\ A_7 &= \{pq^2x \mid x = 1, 2, \dots, p - 1\}. \end{aligned}$$

We can calculate the number of vertices of all zero-divisor sets as $|A_1| = (p-1)q(q-1)$, $|A_2| = p(p-1)(q-1)$, $|A_3| = q(q-1)$, $|A_4| = p(p-1)$, $|A_5| = (p-1)(q-1)$, $|A_6| = (q-1)$ and $|A_7| = (p-1)$. Also, the degree of each vertex in these zero-divisor sets can be determined as

$$d_u = \begin{cases} |A_7|, & u \in A_1 \\ |A_6|, & u \in A_2 \\ |A_4| + |A_7|, & u \in A_3 \\ |A_3| + |A_6|, & u \in A_4 \\ |A_5| + |A_6| + |A_7| - 1, & u \in A_5 \\ |A_2| + |A_4| + |A_5| + |A_6| + |A_7| - 1, & u \in A_6 \\ |A_1| + |A_3| + |A_5| + |A_6| + |A_7| - 1, & u \in A_7 \end{cases}$$

Then, the forgotten index of $\Gamma(\mathbb{Z}_{p^2q^2})$ can be attained that

$$\begin{aligned}
\text{FT}(\Gamma(\mathbb{Z}_{p^2q^2})) &= \sum_{u \in V(\Gamma(\mathbb{Z}_{p^2q^2}))} d_u^3 \\
&= \sum_{i=1}^7 \sum_{u \in A_i} d_u^3 \\
&= |A_1||A_7|^3 + |A_2||A_6|^3 + |A_3|(|A_4| + |A_7|)^3 + |A_4|(|A_3| + |A_6|)^3 \\
&\quad + |A_5|(|A_5| + |A_6| + |A_7| - 1)^3 \\
&\quad + |A_6|(|A_2| + |A_4| + |A_5| + |A_6| + |A_7| - 1)^3 \\
&\quad + |A_7|(|A_1| + |A_3| + |A_5| + |A_6| + |A_7| - 1)^3 \\
&= (p-1)q(q-1)(p-1)^3 + p(p-1)(q-1)(q-1)^3 \\
&\quad + q(q-1)(p-1)^3(p+1)^3 + p(p-1)(q-1)^3(q+1)^3 \\
&\quad + (p-1)(q-1) \left[(p-1)(q-1) + (q-1) + (p-1) - 1 \right]^3 \\
&\quad + (q-1) \left[p(p-1)(q-1) + p(p-1) + (p-1)(q-1) + (q-1) + (p-1) - 1 \right]^3 \\
&\quad + (p-1) \left[(p-1)q(q-1) + q(q-1) + (p-1)(q-1) + (q-1) + (p-1) - 1 \right]^3 \\
&= (p-1)^4q(q-1) + p(p-1)(q-1)^4 + (p-1)^3(p+1)^3q(q-1) \\
&\quad + p(p-1)(q-1)^3(q+1)^3 + (p-1)(q-1)(pq-2)^3 \\
&\quad + (q-1)(p^2q-2)^3 + (p-1)(pq^2-2)^3 \\
&= (p-1)(q-1) \left[(p-1)^3q + p(q-1)^3 + (p-1)^2(p+1)^3q \right. \\
&\quad \left. + p(q-1)^2(q+1)^3 + (pq-2)^3 + \frac{(p^2q-2)^3}{p-1} + \frac{(pq^2-2)^3}{q-1} \right].
\end{aligned}$$

□

Example 2.9 For the graph $\Gamma(\mathbb{Z}_{225})$, we have $n = p^2q^2$ where $p = 3$ and $q = 5$. Then, $\text{FT}(\Gamma(\mathbb{Z}_{225})) =$

1, 208, 678, and the set of zero-divisors can be written as follows.

$$\begin{aligned}
 A_1 &= \{3, 6, 12, 21, 24, 33, 39, 42, 48, 51, 57, 66, 69, 78, 84, 87, 93, 96, 102, 111, 114, \\
 &\quad 123, 129, 132, 138, 141, 147, 156, 159, 168, 174, 177, 183, 186, 192, 201, 204, \\
 &\quad 213, 219, 222\}, \\
 A_2 &= \{5, 10, 20, 35, 40, 55, 65, 70, 80, 85, 95, 110, 115, 130, 140, 145, 155, 160, 170, \\
 &\quad 185, 190, 205, 215, 220\}, \\
 A_3 &= \{9, 18, 27, 36, 54, 63, 72, 81, 99, 108, 117, 126, 144, 153, 162, 171, 189, 198, \\
 &\quad 207, 216\}, \\
 A_4 &= \{25, 50, 100, 125, 175, 200\}, \\
 A_5 &= \{15, 30, 60, 105, 120, 165, 195, 210\}, \\
 A_6 &= \{45, 90, 135, 180\}, \\
 A_7 &= \{75, 150\}.
 \end{aligned}$$

These sets build the graph $\Gamma(\mathbb{Z}_{225})$, where $A_1 = \{3x \mid x = 1, 2, \dots, pq^2 - 1, p \nmid x, q \nmid x\}$, $A_2 = \{5x \mid x = 1, 2, \dots, p^2q - 1, p \nmid x, q \nmid x\}$, $A_3 = \{9x \mid x = 1, 2, \dots, q^2 - 1, q \nmid x\}$, $A_4 = \{25x \mid x = 1, 2, \dots, p^2 - 1, p \nmid x\}$, $A_5 = \{15x \mid x = 1, 2, \dots, pq - 1, p \nmid x, q \nmid x\}$, $A_6 = \{45x \mid x = 1, 2, \dots, q - 1\}$, and $A_7 = \{75x \mid x = 1, 2, \dots, p - 1\}$.

In the next theorem, the relation of the forgotten topological index of $\Gamma(\mathbb{Z}_{pqr})$ is represented.

Theorem 2.10 *Let $\Gamma(\mathbb{Z}_{pqr})$ be a zero-divisor graph and p, q and r be distinct prime numbers. Then, the forgotten topological index of $\Gamma(\mathbb{Z}_{pqr})$ is*

$$\begin{aligned}
 FT(\Gamma(\mathbb{Z}_{pqr})) &= (p-1)(q-1)(r-1) \left[(p-1)^2 + (q-1)^2 + (r-1)^2 \right. \\
 &\quad \left. + \frac{(pq-1)^3}{(p-1)(q-1)} + \frac{(pr-1)^3}{(p-1)(r-1)} + \frac{(qr-1)^3}{(q-1)(r-1)} \right].
 \end{aligned}$$

Proof Since the proper divisors of $n = pqr$ are p, q, r, pq, pr and qr , then the vertex set can be partitioned as $V(\Gamma(\mathbb{Z}_n)) = A_1 \cup A_2 \cup A_3 \cup A_4 \cup A_5 \cup A_6$ and $A_i \cap A_j = \emptyset$ for $i \neq j$, where $i, j \in \{1, \dots, 6\}$ and

$$\begin{aligned}
 A_1 &= \{px \mid x = 1, 2, \dots, qr - 1, q \nmid x, r \nmid x\}, \\
 A_2 &= \{qx \mid x = 1, 2, \dots, pr - 1, p \nmid x, r \nmid x\}, \\
 A_3 &= \{rx \mid x = 1, 2, \dots, pq - 1, p \nmid x, q \nmid x\}, \\
 A_4 &= \{pqx \mid x = 1, 2, \dots, r - 1\}, \\
 A_5 &= \{prx \mid x = 1, 2, \dots, q - 1\}, \\
 A_6 &= \{qrx \mid x = 1, 2, \dots, p - 1\}.
 \end{aligned}$$

The number of vertices of all zero-divisor sets can be calculated as $|A_1| = (q - 1)(r - 1)$, $|A_2| = (p - 1)(r - 1)$, $|A_3| = (p - 1)(q - 1)$, $|A_4| = (r - 1)$, $|A_5| = (q - 1)$, and $|A_6| = (p - 1)$. Besides, the degree of each vertex in these zero-divisor sets can be determined as

$$d_u = \begin{cases} |A_6|, & u \in A_1 \\ |A_5|, & u \in A_2 \\ |A_4|, & u \in A_3 \\ |A_3| + |A_5| + |A_6|, & u \in A_4 \\ |A_2| + |A_4| + |A_6|, & u \in A_5 \\ |A_1| + |A_4| + |A_5|, & u \in A_6 \end{cases}.$$

Hence, we can obtain the forgotten index of $\Gamma(\mathbb{Z}_{pqr})$ as

$$\begin{aligned} \text{FT}(\Gamma(\mathbb{Z}_{pqr})) &= \sum_{u \in V(\Gamma(\mathbb{Z}_{p^2q}))} d_u^3 \\ &= \sum_{i=1}^6 \sum_{u \in A_i} d_u^3 \\ &= |A_1||A_6|^3 + |A_2||A_5|^3 + |A_3||A_4|^3 + |A_4|(|A_3| + |A_5| + |A_6|)^3 \\ &\quad + |A_5|(|A_2| + |A_4| + |A_6|)^3 + |A_6|(|A_1| + |A_4| + |A_5|)^3 \\ &= (q - 1)(r - 1)(p - 1)^3 + (p - 1)(r - 1)(q - 1)^3 + (p - 1)(q - 1)(r - 1)^3 \\ &\quad + (r - 1)((p - 1)(q - 1) + (q - 1) + (p - 1))^3 \\ &\quad + (q - 1)((p - 1)(r - 1) + (p - 1) + (q - 1))^3 \\ &\quad + (p - 1)((q - 1)(r - 1) + (q - 1) + (r - 1))^3 \\ &= (p - 1)^3(q - 1)(r - 1) + (p - 1)(q - 1)^3(r - 1) + (p - 1)(q - 1)(r - 1)^3 \\ &\quad + (pq - 1)^3(r - 1) + (pr - 1)^3(q - 1) + (qr - 1)(p - 1) \\ &= (p - 1)(q - 1)(r - 1) \left[(p - 1)^2 + (q - 1)^2 + (r - 1)^2 \right. \\ &\quad \left. + \frac{(pq - 1)^3}{(p - 1)(q - 1)} + \frac{(pr - 1)^3}{(p - 1)(r - 1)} + \frac{(qr - 1)^3}{(q - 1)(r - 1)} \right]. \end{aligned}$$

□

Example 2.11 For the graph $\Gamma(\mathbb{Z}_{165})$, we have $n = pqr$ where $p = 3$, $q = 5$ and $r = 11$. Then,

$FT(\Gamma(\mathbb{Z}_{165})) = 483,040$, and the set of zero-divisors can be written as follows:

$$\begin{aligned}
 A_1 &= \{3, 6, 9, 12, 18, 21, 24, 27, 36, 39, 42, 48, 51, 54, 57, 63, 69, 72, 78, 81, 84, 87, 93, \\
 &\quad 96, 102, 108, 111, 114, 117, 123, 126, 129, 138, 141, 144, 147, 153, 156, 159, 162\}, \\
 A_2 &= \{5, 10, 20, 25, 35, 40, 50, 65, 70, 80, 85, 95, 100, 115, 125, 130, 140, 145, 155, 160\}, \\
 A_3 &= \{11, 22, 44, 77, 88, 121, 143, 154\}, \\
 A_4 &= \{15, 30, 45, 60, 75, 90, 105, 120, 135, 150\}, \\
 A_5 &= \{33, 66, 99, 132\}, \\
 A_6 &= \{55, 110\}.
 \end{aligned}$$

Additionally, these sets give rise to the graph which can be shown in Figure 4, where $A_1 = \{3x \mid x = 1, 2, \dots, qr - 1, q \nmid x, r \nmid x\}$, $A_2 = \{5x \mid x = 1, 2, \dots, pr - 1, p \nmid x, r \nmid x\}$, $A_3 = \{11x \mid x = 1, 2, \dots, pq - 1, p \nmid x, q \nmid x\}$, $A_4 = \{15x \mid x = 1, 2, \dots, r - 1\}$, $A_5 = \{33x \mid x = 1, 2, \dots, q - 1\}$, $A_6 = \{55x \mid x = 1, 2, \dots, p - 1\}$.




Figure 4. The graph $\Gamma(\mathbb{Z}_{165})$

2.1. Matlab code for determining the forgotten topological index of $\Gamma(\mathbb{Z}_n)$

In this subsection, we give an algorithm for calculating the forgotten topological index of $\Gamma(\mathbb{Z}_n)$ when entering an integer n .

```

1 n=input("$"enter n for Z_n:"$)
2 Vert=strings(1,n-2);
3 Adj=zeros(n-2);
4 Deg=zeros(1,n-2);
5 for i=2:n-1

```

```

6 Vert(i-1)=int2str(i);
7 for j=2:n-1
8 if (i==j), continue, end
9 if mod(i*j,n)==0
10 Adj(i-1,j-1)=1;
11 Deg(i-1)=Deg(i-1)+1;
12 end
13 end
14 end
15 for i=size(Deg,2):-1:1
16 if (Deg(i)==0)
17 Adj(i,:)=[];
18 Adj(:,i)=[];
19 Vert(i)=[];
20 Deg(i)=[];
21 end
22 end
23 fi=0;
24 for i=1:size(Deg,2)
25 $fi = fi + Deg(i)^3;$
26 end
27 fprintf("Forgotten Index: %d\n",fi);

```

In the lines 1-4 of the algorithm, an integer n for \mathbb{Z}_n is requested, and the vertex set ($Vert$), the adjacency matrix (Adj) and the degree array (Deg) are initialized. Next, in line 7, all possible vertices in the graph are inserted into the set. Then, as long as the condition $i \cdot j \equiv 0 \pmod{n}$, the adjacency matrix is filled and the degree array is updated in lines 11-12. After that, vertices having no neighbors are removed from vertex set, degree array and adjacency matrix in lines 17-24. To compute the forgotten index of a zero divisor graph, the lines 17-24 are not obligatory. However, these lines are used to represent the zero divisor graph properly. Finally, in lines 26-30, the forgotten topological index of graph $\Gamma(\mathbb{Z}_n)$ is calculated and printed out.

In this algorithm, lines 17-24 are not obligatory. However, these lines are used to construct the graph $\Gamma(\mathbb{Z}_n)$ properly.

3. Forgotten topological index of zero-divisor graph of products of rings of integers modulo n

In this section we calculate the forgotten index of the graphs $\Gamma(\mathbb{Z}_p \times \mathbb{Z}_q)$ and $\Gamma(\mathbb{Z}_p \times \mathbb{Z}_q \times \mathbb{Z}_r)$ for distinct prime numbers p , q and r .

The zero-divisor graph of $\mathbb{Z}_p \times \mathbb{Z}_q$ and some graph theoretical properties of it have been studied in [2]. In the following theorem, we give the forgotten index of $\Gamma(\mathbb{Z}_p \times \mathbb{Z}_q)$.

Theorem 3.1 *Let $\Gamma(\mathbb{Z}_p \times \mathbb{Z}_q)$ be a zero-divisor graph and p, q be distinct prime numbers. Then the forgotten index of $\Gamma(\mathbb{Z}_p \times \mathbb{Z}_q)$ is*

$$FT(\Gamma(\mathbb{Z}_p \times \mathbb{Z}_q)) = (p-1)(q-1)((p-1)^2 + (q-1)^2).$$

Proof Let $x \in \mathbb{Z}_p^*$ and $y \in \mathbb{Z}_q^*$, where $x = 1, 2, \dots, p-1$ and $y = 1, 2, \dots, q-1$. Since $(x, 0)(0, y) = (0, 0)$ the edge set of $x \in \mathbb{Z}_p^*$ contains only the edges between the vertices $(x, 0)$ and $(0, y)$.

The graph $\Gamma(\mathbb{Z}_p \times \mathbb{Z}_q)$ is a complete bipartite graph $K_{p-1, q-1}$. Partitions of vertex set of $\Gamma(\mathbb{Z}_p \times \mathbb{Z}_q)$ are

$$A_1 = \{(x, 0) \mid 1 \leq x < p, x \in \mathbb{Z}_p\},$$

$$A_2 = \{(0, y) \mid 1 \leq y < q, y \in \mathbb{Z}_q\}$$

such that $A_1 \cup A_2 = V(\Gamma(\mathbb{Z}_p \times \mathbb{Z}_q))$ and $A_1 \cap A_2 = \emptyset$. Since $|A_1| = p - 1$ and $|A_2| = q - 1$, the size of this graph is $(p - 1)(q - 1)$. Also, $d_u = |A_2|$ for all $u \in A_1$ and $d_v = |A_1|$ for all $v \in A_2$. Hence, we obtain

$$\begin{aligned} \text{FT}(\Gamma(\mathbb{Z}_p \times \mathbb{Z}_q)) &= \sum_{u \in V(\Gamma(\mathbb{Z}_p \times \mathbb{Z}_q))} d_u^2 \\ &= \sum_{i=1}^2 \sum_{u \in A_i} d_u^3 \\ &= |A_1| (q - 1)^3 + |A_2| (p - 1)^3 \\ &= (p - 1)(q - 1)^3 + (p - 1)^3(q - 1) \\ &= (p - 1)(q - 1)((p - 1)^2 + (q - 1)^2). \end{aligned}$$

□

Example 3.2 For zero-divisor graph of $\mathbb{Z}_{11} \times \mathbb{Z}_{17}$, we attain $p = 11$ and $q = 17$. Then, $\text{FT}(\Gamma(\mathbb{Z}_{11} \times \mathbb{Z}_{17})) = 56,960$, and the set of zero-divisors as follows:

$$\begin{aligned} A_1 &= \{(1, 0), (2, 0), (3, 0), (4, 0), (5, 0), (6, 0), (7, 0), (8, 0), (9, 0), (10, 0)\}, \\ A_2 &= \{(0, 1), (0, 2), (0, 3), (0, 4), (0, 5), (0, 6), (0, 7), (0, 8), (0, 9), (0, 10), (0, 11), \\ &\quad (0, 12), (0, 13), (0, 14), (0, 15), (0, 16)\}. \end{aligned}$$

These sets give rise to the graph depicted in Figure 5, where $A_1 = \{(x, 0) \mid 1 \leq x < p, x \in \mathbb{Z}_{11}\}$ and $A_2 = \{(0, y) \mid 1 \leq y < q, y \in \mathbb{Z}_{17}\}$.

Akgunes and Nacaroglu have studied some properties of zero-divisor graph of $\mathbb{Z}_p \times \mathbb{Z}_q \times \mathbb{Z}_r$ [3]. Moreover, they have calculated irregularity index and Zagreb indices of this graph. We obtain the forgotten index of $\Gamma(\mathbb{Z}_p \times \mathbb{Z}_q \times \mathbb{Z}_r)$ in the following theorem.

Theorem 3.3 Let $\Gamma(\mathbb{Z}_p \times \mathbb{Z}_q \times \mathbb{Z}_r)$ be a zero-divisor graph and p, q, r be distinct prime numbers. Then, the forgotten index of $\Gamma(\mathbb{Z}_p \times \mathbb{Z}_q \times \mathbb{Z}_r)$ is

$$\begin{aligned} \text{FT}(\Gamma(\mathbb{Z}_p \times \mathbb{Z}_q \times \mathbb{Z}_r)) &= (p - 1)(q - 1)(r - 1) \left[(p - 1)^2 + (q - 1)^2 + (r - 1)^2 \right. \\ &\quad \left. + \frac{(pq - 1)^3}{(p - 1)(q - 1)} + \frac{(pr - 1)^3}{(p - 1)(r - 1)} + \frac{(qr - 1)^3}{(q - 1)(r - 1)} \right]. \end{aligned}$$

Proof We divide the vertex set of $\Gamma(\mathbb{Z}_p \times \mathbb{Z}_q \times \mathbb{Z}_r)$ into six subsets such that $V(\Gamma(\mathbb{Z}_p \times \mathbb{Z}_q \times \mathbb{Z}_r)) = \cup_{i=1}^6 A_i$




Figure 5. The graph $\Gamma(\mathbb{Z}_{11} \times \mathbb{Z}_{17})$

and $A_i \cap A_j = \emptyset$, where $i = 1, 2, \dots, 5$ and $j = i + 1, \dots, 6$. We show that these vertex subsets as follows:

$$A_1 = \{(x, 0, 0) \mid 1 \leq x < p, x \in \mathbb{Z}_p\},$$

$$A_2 = \{(0, y, 0) \mid 1 \leq y < q, y \in \mathbb{Z}_q\},$$

$$A_3 = \{(0, 0, z) \mid 1 \leq z < r, z \in \mathbb{Z}_r\},$$

$$A_4 = \{(0, y, z) \mid 1 \leq y < q, 1 \leq z < r, y \in \mathbb{Z}_q, z \in \mathbb{Z}_r\},$$

$$A_5 = \{(x, 0, z) \mid 1 \leq x < p, 1 \leq z < r, x \in \mathbb{Z}_p, z \in \mathbb{Z}_r\},$$

$$A_6 = \{(x, y, 0) \mid 1 \leq x < p, 1 \leq y < q, x \in \mathbb{Z}_p, y \in \mathbb{Z}_q\}.$$

The number of vertices of all zero-divisor sets can be calculated as $|A_1| = (p - 1)$, $|A_2| = (q - 1)$, $|A_3| = (r - 1)$, $|A_4| = (q - 1)(r - 1)$, $|A_5| = (p - 1)(r - 1)$, and $|A_6| = (p - 1)(q - 1)$. Moreover, the degree of each vertex in these zero-divisor sets can be determined as

$$d_u = \begin{cases} |A_2| + |A_3| + |A_4|, & u \in A_1 \\ |A_1| + |A_3| + |A_5|, & u \in A_2 \\ |A_1| + |A_2| + |A_6|, & u \in A_3 \\ |A_1|, & u \in A_4 \\ |A_2|, & u \in A_5 \\ |A_3|, & u \in A_6 \end{cases}.$$

According to these subsets, we calculate the forgotten index of the graph $\Gamma(\mathbb{Z}_p \times \mathbb{Z}_q \times \mathbb{Z}_r)$ as follows:

$$\begin{aligned}
\text{FT}(\Gamma(\mathbb{Z}_p \times \mathbb{Z}_q \times \mathbb{Z}_r)) &= \sum_{u \in V(\Gamma(\mathbb{Z}_{p^2q}))} d_u^3 \\
&= \sum_{i=1}^6 \sum_{u \in A_i} d_u^3 \\
&= |A_1|(|A_2| + |A_3| + |A_4|)^3 + |A_2|(|A_1| + |A_3| + |A_5|)^3 \\
&\quad + |A_3|(|A_1| + |A_2| + |A_6|)^3 + |A_4||A_1|^3 + |A_5||A_2|^3 + |A_6||A_3|^3 \\
&= (p-1)((q-1)(r-1) + (q-1) + (r-1))^3 \\
&\quad + (q-1)((p-1)(r-1) + (p-1) + (r-1))^3 \\
&\quad + (r-1)((p-1)(q-1) + (p-1) + (q-1))^3 \\
&\quad + (q-1)(r-1)(p-1)^3 + (p-1)(r-1)(q-1)^3 \\
&\quad + (p-1)(q-1)(r-1)^3 \\
&= (p-1)(q-1)(r-1) \left[(p-1)^2 + (q-1)^2 + (r-1)^2 \right. \\
&\quad \left. + \frac{(pq-1)^3}{(p-1)(q-1)} + \frac{(qr-1)^3}{(q-1)(r-1)} + \frac{(pr-1)^3}{(p-1)(r-1)} \right].
\end{aligned}$$

Example 3.4 For zero-divisor graph of $\mathbb{Z}_3 \times \mathbb{Z}_5 \times \mathbb{Z}_{13}$, we have $p = 3$, $q = 5$ and $r = 13$. Then, $\text{FT}(\Gamma(\mathbb{Z}_3 \times \mathbb{Z}_5 \times \mathbb{Z}_{13})) = 792,448$, and the following are the set of zero-divisors of this ring. The zero-divisor graph of this ring can be seen in Figure 6: □

$$A_1 = \{(1, 0, 0), (2, 0, 0)\},$$

$$A_2 = \{(0, 1, 0), (0, 2, 0), (0, 3, 0), (0, 4, 0)\},$$

$$A_3 = \{(0, 0, 1), (0, 0, 2), (0, 0, 3), (0, 0, 4), (0, 0, 5), (0, 0, 6), (0, 0, 7), (0, 0, 8), (0, 0, 9), \\ (0, 0, 10), (0, 0, 11), (0, 0, 12)\},$$

$$A_4 = \{(0, 1, 1), (0, 1, 2), (0, 1, 3), (0, 1, 4), (0, 1, 5), (0, 1, 6), (0, 1, 7), (0, 1, 8), (0, 1, 9), \\ (0, 1, 10), (0, 1, 11), (0, 1, 12), (0, 2, 1), (0, 2, 2), (0, 2, 3), (0, 2, 4), (0, 2, 5), (0, 2, 6), \\ (0, 2, 7), (0, 2, 8), (0, 2, 9), (0, 2, 10), (0, 2, 11), (0, 2, 12), (0, 3, 1), (0, 3, 2), (0, 3, 3), \\ (0, 3, 4), (0, 3, 5), (0, 3, 6), (0, 3, 7), (0, 3, 8), (0, 3, 9), (0, 3, 10), (0, 3, 11), (0, 3, 12), \\ (0, 4, 1), (0, 4, 2), (0, 4, 3), (0, 4, 4), (0, 4, 5), (0, 4, 6), (0, 4, 7), (0, 4, 8), (0, 4, 9), \\ (0, 4, 10), (0, 4, 11), (0, 4, 12)\},$$

$$A_5 = \{(1, 0, 1), (1, 0, 2), (1, 0, 3), (1, 0, 4), (1, 0, 5), (1, 0, 6), (1, 0, 7), (1, 0, 8), (1, 0, 9), \\ (1, 0, 10), (1, 0, 11), (1, 0, 12), (2, 0, 1), (2, 0, 2), (2, 0, 3), (2, 0, 4), (2, 0, 5), (2, 0, 6), \\ (2, 0, 7), (2, 0, 8), (2, 0, 9), (2, 0, 10), (2, 0, 11), (2, 0, 12)\},$$

$$A_6 = \{(1, 1, 0), (1, 2, 0), (1, 3, 0), (1, 4, 0), (2, 1, 0), (2, 2, 0), (2, 3, 0), (2, 4, 0)\}.$$

By the above arguments, we give the following corollary.




Figure 6. The graph $\Gamma(\mathbb{Z}_3 \times \mathbb{Z}_5 \times \mathbb{Z}_{13})$

Corollary 3.5 Let $\Gamma(\mathbb{Z}_p \times \mathbb{Z}_q)$, $\Gamma(\mathbb{Z}_p \times \mathbb{Z}_q \times \mathbb{Z}_r)$, $\Gamma(\mathbb{Z}_{pq})$, and $\Gamma(\mathbb{Z}_{pqr})$ be zero-divisor graphs where p , q , and r are distinct prime numbers. Then the followings hold:

- (i) $FT(\Gamma(\mathbb{Z}_p \times \mathbb{Z}_q)) = FT(\Gamma(\mathbb{Z}_{pq}))$,
- (ii) $FT(\Gamma(\mathbb{Z}_p \times \mathbb{Z}_q \times \mathbb{Z}_r)) = FT(\Gamma(\mathbb{Z}_{pqr}))$.

References

- [1] Aijaz M, Pirzada S. Annihilating-ideal graphs of commutative rings. Asian-European Journal of Mathematics 2020; 13 (07): 1-12. doi: 10.1142/S1793557120501211
- [2] Akgunes N, Togan M. Some graph theoretical properties over zero-divisor graphs of special finite commutative rings. Advanced Studies in Contemporary Mathematics (Kyungshang) 2012; 22 (2): 305-315.
- [3] Akgunes N, Nacaroglu Y. Some properties of zero divisor graph obtained by the ring $\mathbb{Z}_p \times \mathbb{Z}_q \times \mathbb{Z}_r$. Asian-European Journal of Mathematics 2019; 12 (06): 1-10. doi: 10.1142/S179355712040001X
- [4] Akbari S, Kiani D, Mohammadi F, Moradi S. The total graph and regular graph of a commutative ring. Journal of Pure and Applied Algebra 2009; 213 (12): 2224-2228. doi: 10.1016/j.jpaa.2009.03.013
- [5] Anderson DF, Livingston PS. The zero-divisor graph of a commutative ring. Journal of Algebra 1999; 217 (2): 434-447. doi: 10.1006/jabr.1998.7840
- [6] Anderson DF, Badawi A. The total graph of a commutative ring. Journal of Algebra 2008; 320 (7): 2706-2719. doi: 10.1016/j.jalgebra.2008.06.028

- [7] Asir T, Rabikka V. The Wiener index of the zero-divisor graph of Zn. *Discrete Applied Mathematics* 2021; 1-11. doi: 10.1016/j.dam.2021.02.035
- [8] Asok A, Kureethara JV. Characterization of the forgotten topological index and the Hyper-Zagreb index for the unicyclic graphs. *MATCH Communications in Mathematical and in Computer Chemistry* 2020; 84 (3): 739-751.
- [9] Beck I. Coloring of commutative rings. *Journal of Algebra* 1988; 116 (1): 208-226. doi: 10.1016/0021-8693(88)90202-5
- [10] Das KC, Gutman I. Some properties of the second Zagreb index. *MATCH Communications in Mathematical and in Computer Chemistry* 2004; 52 (1): 103-112.
- [11] Das KC, Çevik AS, Cangul IN, Shang Y. On sombor index. *Symmetry* 2021; 13 (1): 1-12. doi: 10.3390/sym13010140
- [12] Furtula B, Gutman I. A forgotten topological index. *Journal of Mathematical Chemistry* 2015; 53 (4): 1184-1190. doi: 10.1007/s10910-015-0480-z
- [13] Gao W, Wang WF. The fifth geometric-arithmetic index of bridge graph and carbon nanocones. *Journal of Difference Equations and Applications* 2017; 23 (1-2): 100-109. doi: 10.1080/10236198.2016.1197214
- [14] Gao W, Wang W, Farahani MR. Topological indices study of molecular structure in anticancer drugs. *Journal of Chemistry* 2016; 1-8. doi: 10.1155/2016/3216327
- [15] Gao W, Farahani MR. Computing the reverse eccentric connectivity index for certain family of nanocone and fullerene structures. *Journal of Nanotechnology* 2016; 1-6. doi: 10.1155/2016/3129561
- [16] Gao W, Farahani MR. Degree-based indices computation for special chemical molecular structures using edge dividing method. *Applied Mathematics and Nonlinear Sciences* 2016; 1 (1): 99-122. doi: 10.21042/AMNS.2016.1.00009
- [17] Gao W, Shi L. Szeged related indices of unilateral polyomino chain and unilateral hexagonal chain. *IAENG International Journal of Applied Mathematics* 2015; 45 (2): 1-13.
- [18] Gao W, Wang W. Second atom-bond connectivity index of special chemical molecular structures. *Journal of Chemistry* 2014. doi: 10.1155/2014/906254
- [19] Gao W, Wang W. The vertex version of weighted Wiener number for bicyclic molecular structures. *Computational and Mathematical Methods in Medicine* 2015; 1-10. doi: 10.1155/2015/418106
- [20] Gao W, Wang W. The eccentric connectivity polynomial of two classes of nanotubes. *Chaos, Solitons & Fractals* 2016; 89: 290-294. doi: 10.1016/j.chaos.2015.11.035
- [21] Gao W, Farahani MR, Shi L. Forgotten topological index of some drug structures. *Acta Medica Mediterranea* 2016; 32 (1): 579-585.
- [22] Ghanbari N, Alikhani S. Sombor index of certain graphs. *Iranian Journal of Mathematical Chemistry* 2021; 12 (1): 27-37. doi: 10.22052/ijmc.2021.242106.1547
- [23] Gutman I, Trinajstić N. Graph theory and molecular orbitals. Total φ -electron energy of alternant hydrocarbons. *Chemical Physics Letters* 1972; 17 (4): 535-538. doi: 10.1016/0009-2614(72)85099-1
- [24] Gutman I, Trinajstić N, Ruscic B, Wilcox CF. Graph theory and molecular orbitals. XII. Acyclic polyenes. *The Journal of Chemical Physics* 1975; 62: 33-99. doi: 10.1063/1.430994
- [25] Gutman I, Furtula B, Toropov AA, Toropova AP. The graph of atomic orbitals and its basic properties. 2. Zagreb indices. *MATCH Communications in Mathematical and in Computer Chemistry* 2005; 53 (1): 225-230.
- [26] Gutman I. Geometric approach to degree-based topological indices: Sombor indices. *MATCH Communications in Mathematical and in Computer Chemistry* 2021; 86 (1): 11-16.
- [27] Gutman I. Some basic properties of Sombor indices. *Open Journal of Discrete Applied Mathematics* 2021; 4 (1): 1-3. doi: 10.30538/psrp-odam2021.0047

- [28] Hansen P, Mélot H, Gutman I. Variable neighborhood search for extremal graphs 12. A note on the variance of bounded degrees in graphs. *MATCH Communications in Mathematical and in Computer Chemistry* 2005; 54 (1): 221-232.
- [29] Havare OC, Havare AK. Computation of the forgotten topological index and co-index for carbon base nanomaterial. *Polycyclic Aromatic Compounds* 2020; 1-13. doi: 10.1080/10406638.2020.1866621
- [30] Khaksari A, Ghorbani M. On the forgotten topological index. *Iranian Journal of Mathematical Chemistry* 2017; 8 (3): 327-338. doi: 10.22052/ijmc.2017.43481
- [31] Liu JB, Matejic MM, Milovanovic EI, Milovanovic IZ. Some new inequalities for the forgotten topological index and coindex of graphs. *MATCH Communications in Mathematical and in Computer Chemistry* 2020; 84 (3): 719-738.
- [32] Mehak GE, Bhatti AA. Forgotten topological index of line graphs of some chemical structures in drugs. *Acta Chemica Iasi* 2018; 26 (2): 181-206.
- [33] Naeini H, Kazemi R, Behzadi MH. The modified forgotten topological index of random b-ary recursive trees. *UPB Scientific Bulletin, Series A: Applied Mathematics and Physics* 2018; 80 (4): 125-132.
- [34] Réti T, Došlic T, Ali A. On the Sombor index of graphs. *Contributions to Mathematics* 2021; 3: 11-18. doi: 10.47443/cm.2021.0006
- [35] Salehifar S, Khashyarmansh K, Afkhami M. On the annihilator-ideal graph of commutative rings. *Ricerche di Matematica* 2017; 66 (2): 431-447. doi: 10.1007/s11587-016-0311-y
- [36] Sarkar P, De N, Pal A. The forgotten topological index of graphs based on new operations related to the join of graphs. In *Harmony Search and Nature Inspired Optimization Algorithms*. Springer, Singapore, 2019, pp. 335-345.
- [37] Shaveisi F. Some results on annihilating-ideal graphs. *Canadian Mathematical Bulletin* 2016; 59 (3): 641-651. doi: 10.4153/CMB-2016-016-3
- [38] Singh P, Bhat VK. Adjacency matrix and Wiener index of zero divisor graph $\Gamma(\mathbb{Z}_n)$. *Journal of Applied Mathematics and Computing* 2021; 66 (1): 717-732. doi: 10.1007/s12190-020-01460-2
- [39] Sinha D, Rao AK. A note on co-maximal graphs of commutative rings. *AKCE International Journal of Graphs and Combinatorics* 2018; 15 (1): 112-114. doi: 10.1016/j.akcej.2018.03.003
- [40] Tehranian A, Maimani HR. A study of the total graph. *Iranian Journal of Mathematical Sciences and Informatics* 2011; 6 (2): 75-80.
- [41] Wang HJ. Graphs associated to co-maximal ideals of commutative rings. *Journal of Algebra* 2008; 320 (7): 2917-2933. doi: 10.1016/j.jalgebra.2008.06.020
- [42] Yan L, Gao W, Li J. General harmonic index and general sum connectivity index of polyomino chains and nanotubes. *Journal of Computational and Theoretical Nanoscience* 2015; 12 (10): 3940-3944. doi: 10.1166/jctn.2015.4308
- [43] Ye M, Wu T. Co-maximal ideal graphs of commutative rings. *Journal of Algebra and its Applications* 2012; 11 (06): 1-14. doi: 10.1142/S0219498812501149
- [44] Zhou B. Zagreb indices. *MATCH Communications in Mathematical and in Computer Chemistry* 2004; 52 (1): 13-18.
- [45] Zhou T, Lin Z, Miao L. The Sombor index of trees and unicyclic graphs with given matching number. arXiv preprint arXiv: 2103.04645. 2021.