

Positive and decreasing solutions for higher order Caputo boundary value problems with sign-changing Green's function

Rian YAN^{1,*}, Yige ZHAO², Xuan LENG¹, Yabing LI²

¹School of Mathematics and Computer Science, Hunan City University, Yiyang, China

²School of Mathematical Sciences, University of Jinan, Jinan, China

Received: 09.10.2023

Accepted/Published Online: 06.05.2024

Final Version: ..202

Abstract: In this paper, Caputo boundary value problems of order $3 < \zeta \leq 4$ are investigated on the interval $[0, 1]$. By Guo-Krasnoselskii fixed point theorem, some criteria of existence and multiplicity of positive and decreasing solutions are established. The main novelty of the paper lies in its capability to achieve positive solutions while the corresponding Green's function changes sign. Finally, two examples are provided to illustrate the application of these results.

Key words: Caputo fractional derivatives, sign-changing Green's function, fixed point theorem

1. Introduction

There is currently great interest in fractional differential equations (FDEs), since these equations appear naturally in modelling many real world processes, see [8, 23]. Many interesting works were presented for the study of theoretical knowledge and applications of FDEs, see [5–7, 15, 17, 20, 26, 27], and the references therein.

Boundary value problems (BVPs) for integer or fractional order differential equations with positive solutions arise in many fields of science and engineering, see [2, 13, 24]. Therefore, the solvability of positive solutions constitute a significant class of problems, see [12, 18, 19, 22]. By using the fixed point theorems on cone, Bai and Lü [1] studied the existence of positive solutions for Riemann-Liouville (R-L) two-point BVPs with nonnegative Green's function. In fact, most of the existing papers have been written on positive solutions are based on the condition the corresponding Green's functions are nonnegative, see [25].

Recently, several papers have discussed on the existence of positive solutions while the Green's function changes sign, see [3, 4, 11, 21, 25]. In [14], Ma established some criteria of existence and nonexistence of positive solutions for nonlinear periodic BVPs under the condition the Green's kernel changes sign. In [16], Sun and Zhao discussed the following BVP

$$\begin{aligned}u'''(\chi) &= f(\chi, u(\chi)), \\ u'(0) &= u(1) = u''(\eta) = 0,\end{aligned}$$

where f is a given function, η is a given constant, the corresponding Green's kernel may changes sign on $[0, 1] \times [0, 1]$. By iterative technique, they gave some existence results of positive solutions for such problem.

*Correspondence: yanrian89@163.com

2010 AMS Mathematics Subject Classification: 34A08, 34B08

Inspired by the above works, in present paper we deals with the follwing BVP

$${}^C D_{0+}^\zeta u(\chi) = f(\chi, u(\chi)), \quad \chi \in [0, 1], \quad (1.1)$$

$$u'(0) = u'''(0) = 0, \quad u'''(\eta) + \lambda u''(0) = 0, \quad u(1) - \gamma u(0) = 0, \quad (1.2)$$

where ${}^C D_{0+}^\zeta$ is the Caputo FD, $3 < \zeta \leq 4$, $0 < \eta, \lambda, \gamma < 1$ are constants, $f : [0, 1] \times [0, +\infty) \rightarrow [0, +\infty)$ is continuous.

To the best of our knowledge, although the idea on obtaining positive solutions while the Green's function changes sign has been considered by some papers, very little is known on applying such idea on higher order Caputo BVPs in the literature. We undertake this investigation in the present paper. By Guo-Krasnoselskii fixed point theorem, for any positive integer $n (n \geq 2)$, our target is to establish some criteria of existence of at least $n - 1$ positive and decreasing solutions for BVP (1.1)–(1.2). The most significant feature is that the present paper capability to achieve positive solutions while the corresponding Green's function changes sign.

The present paper is organized as follows. In Section 2, some useful definitions are introduced, and some lemmas are proved. In Section 3, some sufficient conditions for the existence of positive and decreasing solutions are derived. Section 4 presents some experiments to explain the results. In Section 5, the conclusion is given.

2. Preliminaries

Definition 2.1 [8] *Let $\zeta > 0$. Then the R-L fractional integral is*

$$I_{0+}^\zeta f(\chi) = \frac{1}{\Gamma(\zeta)} \int_0^\chi (\chi - \xi)^{\zeta-1} f(\xi) d\xi.$$

Definition 2.2 [8] *Let $\zeta > 0$. Then the Caputo FD is*

$${}^C D_{0+}^\zeta f(\chi) = \frac{d^n}{d\chi^n} \int_0^\chi \frac{(\chi - \xi)^{n-\zeta-1}}{\Gamma(n-\zeta)} \left(f(\xi) - \sum_{k=0}^{n-1} \frac{f^{(k)}(0)}{k!} \xi^k \right) d\xi, \quad (2.1)$$

where $n = [\zeta] + 1$ for $\zeta \notin \mathbb{N}_0$; $n = \zeta$ for $\zeta \in \mathbb{N}_0$, $\mathbb{N}_0 = \{0, 1, \dots\}$. If $f \in AC^n[0, 1]$, then the Caputo FD is

$${}^C D_{0+}^\zeta f(\chi) = \frac{1}{\Gamma(n-\zeta)} \int_0^\chi (\chi - \xi)^{n-\zeta-1} f^{(n)}(\xi) d\xi. \quad (2.2)$$

Lemma 2.3 [8] *Let $\zeta > 0$ and let n be given by Definition 2.2. If $f \in AC^n[0, 1]$, then*

$$I_{0+}^\zeta {}^C D_{0+}^\zeta f(\chi) = f(\chi) - \sum_{k=0}^{n-1} \frac{f^{(k)}(0)}{k!} \chi^k.$$

Lemma 2.4 *The BVP*

$${}^C D_{0+}^\zeta u(\chi) = y(\chi), \quad 3 < \zeta \leq 4, \quad y \in C[0, 1], \quad (2.3)$$

$$u'(0) = u'''(0) = 0, \quad u'''(\eta) + \lambda u''(0) = 0, \quad u(1) - \gamma u(0) = 0, \quad (2.4)$$

has a unique solution

$$u(\chi) = \int_0^1 G(\chi, \xi) y(\xi) d\xi, \quad (2.5)$$

where

$$G(\chi, \xi) = g_1(\chi, \xi) + g_2(\chi, \xi) + g_3(\chi, \xi), \quad (2.6)$$

and

$$\begin{aligned} g_1(\chi, \xi) &= -\frac{1}{(1-\gamma)\Gamma(\zeta)}(1-\xi)^{\zeta-1}, \quad (\chi, \xi) \in [0, 1] \times [0, 1], \\ g_2(\chi, \xi) &= \begin{cases} 0, & 0 \leq \chi \leq \xi \leq 1, \\ \frac{(\chi-\xi)^{\zeta-1}}{\Gamma(\zeta)}, & 0 \leq \xi \leq \chi \leq 1, \end{cases} \\ g_3(\chi, \xi) &= \begin{cases} 0, & \xi \geq \eta, \\ \frac{(\frac{1}{1-\gamma}-\chi^2)(\eta-\xi)^{\zeta-4}}{2\lambda\Gamma(\zeta-3)}, & \xi < \eta. \end{cases} \end{aligned}$$

Proof. By Lemma 2.3, we may transfer (2.3) to the integral equation

$$u(\chi) - \sum_{k=0}^3 \frac{u^{(k)}(0)}{k!} \chi^k = \frac{1}{\Gamma(\zeta)} \int_0^\chi (\chi - \xi)^{\zeta-1} y(\xi) d\xi.$$

Using the condition of $u'(0) = u''(0) = 0$, it follows

$$u(\chi) - u(0) - \frac{u''(0)}{2} \chi^2 = \frac{1}{\Gamma(\zeta)} \int_0^\chi (\chi - \xi)^{\zeta-1} y(\xi) d\xi.$$

According to $u'''(\eta) + \lambda u''(0) = 0$, $u(1) - \gamma u(0) = 0$, we have

$$\begin{aligned} u''(0) &= \frac{-1}{\lambda\Gamma(\zeta-3)} \int_0^\eta (\eta - \xi)^{\zeta-4} y(\xi) d\xi, \\ u(0) &= -\frac{1}{(1-\gamma)\Gamma(\zeta)} \int_0^1 (1 - \xi)^{\zeta-1} y(\xi) d\xi - \frac{1}{2(1-\gamma)} u''(0). \end{aligned}$$

Thus

$$\begin{aligned} u(\chi) &= \frac{1}{\Gamma(\zeta)} \int_0^\chi (\chi - \xi)^{\zeta-1} y(\xi) d\xi - \frac{1}{(1-\gamma)\Gamma(\zeta)} \int_0^1 (1 - \xi)^{\zeta-1} y(\xi) d\xi + \frac{\frac{1}{1-\gamma}-\chi^2}{2\lambda\Gamma(\zeta-3)} \int_0^\eta (\eta - \xi)^{\zeta-4} y(\xi) d\xi \\ &= \int_0^1 \left(g_1(\chi, \xi) + g_2(\chi, \xi) + g_3(\chi, \xi) \right) y(\xi) d\xi \\ &= \int_0^1 G(\chi, \xi) y(\xi) d\xi. \end{aligned}$$

Conversely, if $u(\chi)$ satisfies the integral expression $u(\chi) = \int_0^1 G(\chi, \xi) y(\xi) d\xi$, then

$$u(\chi) = I_{0+}^\zeta y(\chi) + c_0 + c_1 \chi + c_2 \chi^2 + c_3 \chi^3, \quad (2.7)$$

where

$$c_0 = -\frac{1}{1-\gamma} I_{0+}^\zeta y(1) + \frac{1}{2(1-\gamma)\lambda} I_{0+}^{\zeta-3} y(\eta), \quad c_2 = \frac{-1}{2\lambda} I_{0+}^{\zeta-3} y(\eta), \quad c_1 = c_3 = 0. \quad (2.8)$$

Since $\zeta > 3$, we have $u'''(\chi) = I_{0+}^{\zeta-3}y(\chi)$. Thus $u \in C^3[0, 1]$. By (2.1), it follows from the equality

$$\begin{aligned} {}^CD_{0+}^{\zeta}u(\chi) &= \frac{d^4}{d\chi^4} \int_0^{\chi} \frac{(\chi-\xi)^{3-\zeta}}{\Gamma(4-\zeta)} \left(u(\xi) - u(0) - u'(0)\xi - \frac{u''(0)}{2}\xi^2 - \frac{u'''(0)}{6}\xi^3 \right) d\xi \\ &= \frac{d^4}{d\chi^4} \int_0^{\chi} \frac{(\chi-\xi)^{3-\zeta}}{\Gamma(4-\zeta)} \left(I_{0+}^{\zeta}y(\xi) + c_0 + c_2\xi^2 - u(0) - u'(0)\xi - \frac{u''(0)}{2}\xi^2 - \frac{u'''(0)}{6}\xi^3 \right) d\xi \\ &= \frac{d^4}{d\chi^4} I_{0+}^{4-\zeta} I_{0+}^{\zeta}y(\chi) = y(\chi) \end{aligned} \quad (2.9)$$

that ${}^CD_{0+}^{\zeta}u \in C[0, 1]$. From the argument used in Remark 2 of [10], we deduce that $u \in AC^4[0, 1]$. The equality (2.9) also implies $u(\chi)$ satisfies (2.3). Moreover, through (2.7) and (2.8), we can obtain $u(\chi)$ satisfies (2.4). Thus, $u(\chi)$ is a solution of the BVP (2.3)–(2.4).

Remark 2.5 From (2.2) in Definition 2.2, we can also get the fact $u(\chi)$ satisfies (2.3). Since $u \in AC^4[0, 1]$, we have

$$\begin{aligned} {}^CD_{0+}^{\zeta}y(\chi) &= \frac{1}{\Gamma(4-\zeta)} \int_0^{\chi} (\chi-\xi)^{3-\zeta} u^{(4)}(\xi) d\xi = \frac{d}{d\chi} \frac{1}{\Gamma(5-\zeta)} \int_0^{\chi} (\chi-\xi)^{4-\zeta} u^{(4)}(\xi) d\xi \\ &= \frac{d}{d\chi} \frac{1}{\Gamma(5-\zeta)} \left(u'''(\xi)(\chi-\xi)^{4-\zeta} \Big|_0^{\chi} + \int_0^{\chi} (4-\zeta)(\chi-\xi)^{3-\zeta} u'''(\xi) d\xi \right) \\ &= \frac{d}{d\chi} \frac{1}{\Gamma(4-\zeta)} \int_0^{\chi} (\chi-\xi)^{3-\zeta} u'''(\xi) d\xi = \frac{d}{d\chi} \frac{1}{\Gamma(4-\zeta)} \int_0^{\chi} (\chi-\xi)^{3-\zeta} I_{0+}^{\zeta-3}y(\xi) d\xi \\ &= \frac{d}{d\chi} I_{0+}^{4-\zeta} I_{0+}^{\zeta-3}y(\chi) = y(\chi). \end{aligned}$$

Remark 2.6 Let

$$\lambda \leq \gamma\eta^2(\zeta - 3) \quad (2.10)$$

hold. Then $G(\chi, \xi)$ given by (2.6) changes sign.

Proof. From the expression of $g_2(\chi, \xi)$ and $g_3(\chi, \xi)$, it follows that $g_2(\chi, \xi)$ is increasing with respect to χ , and $g_3(\chi, \xi)$ is decreasing with respect to χ for $\xi < \eta$.

For $\xi \geq \eta$, $0 < \eta < 1$, $(\chi, \xi) \in [0, 1] \times [0, 1]$, it implies

$$\begin{aligned} G(\chi, \xi) &= g_1(\chi, \xi) + g_2(\chi, \xi) + g_3(\chi, \xi) = -\frac{(1-\xi)^{\zeta-1}}{(1-\gamma)\Gamma(\zeta)} + g_2(\chi, \xi) \\ &\leq -\frac{(1-\xi)^{\zeta-1}}{(1-\gamma)\Gamma(\zeta)} + g_2(1, \xi) = -\frac{(1-\xi)^{\zeta-1}}{(1-\gamma)\Gamma(\zeta)} + \frac{(1-\xi)^{\zeta-1}}{\Gamma(\zeta)}. \end{aligned}$$

Since $3 < \zeta \leq 4$, $0 < \gamma < 1$, we have

$$G(\chi, \xi) \leq \frac{-(1-\xi)^{\zeta-1} + (1-\gamma)(\chi-\xi)^{\zeta-1}}{(1-\gamma)\Gamma(\zeta)} < \frac{-(1-\xi)^{\zeta-1} + (\chi-\xi)^{\zeta-1}}{(1-\gamma)\Gamma(\zeta)} \leq 0.$$

Next, we consider the case $\xi < \eta$, $0 < \eta < 1$, $(\chi, \xi) \in [0, 1] \times [0, 1]$, we have

$$\begin{aligned} G(\chi, \xi) &\geq -\frac{(1-\xi)^{\zeta-1}}{(1-\gamma)\Gamma(\zeta)} + g_2(\chi, \xi) + g_3(1, \xi) \geq -\frac{(1-\xi)^{\zeta-1}}{(1-\gamma)\Gamma(\zeta)} + \frac{(\frac{1}{1-\gamma}-1)(\eta-\xi)^{\zeta-4}}{2\lambda\Gamma(\zeta-3)} \\ &= \frac{1}{\Gamma(\zeta)} \left(-\frac{(1-\xi)^{\zeta-1}}{(1-\gamma)} + \frac{(\frac{1}{1-\gamma}-1)(\zeta-1)(\zeta-2)(\zeta-3)(\eta-\xi)^{\zeta-4}}{2\lambda} \right) \\ &\geq \frac{1}{\Gamma(\zeta)} \left(-\frac{1}{1-\gamma} + \frac{(\frac{1}{1-\gamma}-1)(\zeta-1)(\zeta-2)(\zeta-3)(\eta-\xi)^{\zeta-4}}{2\lambda} \right). \end{aligned}$$

It follows from (2.10) that

$$G(\chi, \xi) \geq \frac{1}{\Gamma(\zeta)} \left(-\frac{1}{1-\gamma} + \frac{(\zeta-1)(\zeta-2)(\eta-\xi)^{\zeta-4}}{2(1-\gamma)\eta^2} \right) \geq \frac{-\eta^2 + (\eta-\xi)^{\zeta-4}}{(1-\gamma)\eta^2\Gamma(\zeta)} > 0.$$

Thus, it gets $G(\chi, \xi) > 0$, $0 \leq \xi < \eta$, and $G(\chi, \xi) \leq 0$, $\eta \leq \xi \leq 1$.

Lemma 2.7 Let (2.10) hold and

$$K_0 = \{y \in C[0, 1] : y \geq 0 \text{ and } y \text{ is decreasing on } [0, 1]\}.$$

Suppose that $y \in K_0$. Then u given by (2.5) satisfies $u \in K_0$, and u is concave on $[0, \eta]$.

Proof. According to Lemma 2.4, we have $u(\chi) = \int_0^1 G(\chi, \xi)y(\xi)d\xi$. Since $y(\chi) \geq 0$, $\chi \in [0, 1]$, we get

$$\begin{aligned} u''(\chi) &= \frac{1}{\Gamma(\zeta-2)} \int_0^\chi (\chi-\xi)^{\zeta-3}y(\xi)d\xi - \frac{1}{\lambda\Gamma(\zeta-3)} \int_0^\eta (\eta-\xi)^{\zeta-4}y(\xi)d\xi \\ &\leq \frac{1}{\Gamma(\zeta-2)} \int_0^\chi (\chi-\xi)^{\zeta-3}y(\xi)d\xi - \frac{1}{\gamma\eta^2(\zeta-3)\Gamma(\zeta-3)} \int_0^\eta (\eta-\xi)^{\zeta-4}y(\xi)d\xi \\ &\leq \frac{1}{\Gamma(\zeta-2)} \left(\int_0^\chi \left((\chi-\xi)^{\zeta-3} - (\eta-\xi)^{\zeta-4} \right) y(\xi)d\xi - \int_\chi^\eta (\eta-\xi)^{\zeta-4}y(\xi)d\xi \right) \\ &\leq 0, \quad \chi \in [0, \eta]. \end{aligned}$$

This shows u is concave on $[0, \eta]$.

Next we will prove $u \in K_0$. For $\chi \in [0, \eta]$, $u'(\chi)$ is decreasing, then

$$u'(\chi) \leq u'(0) = 0. \quad (2.11)$$

For $\chi \in (\eta, 1]$, in view of $y \in K_0$ and (2.10), it gets

$$\begin{aligned} u'(\chi) &= \frac{1}{\Gamma(\zeta-1)} \int_0^\chi (\chi-\xi)^{\zeta-2}y(\xi)d\xi - \frac{\chi}{\lambda\Gamma(\zeta-3)} \int_0^\eta (\eta-\xi)^{\zeta-4}y(\xi)d\xi \\ &\leq \frac{1}{\Gamma(\zeta-1)} \int_0^1 y(\xi)d\xi - \frac{\eta^2}{\lambda\Gamma(\zeta-3)} \int_0^1 y(\eta\xi)d\xi \\ &\leq \frac{1}{\Gamma(\zeta-1)} \int_0^1 y(\xi)d\xi - \frac{1}{\gamma\Gamma(\zeta-2)} \int_0^1 y(\eta\xi)d\xi \leq 0. \end{aligned} \quad (2.12)$$

As a consequence of (2.11) and (2.12), we have $u'(\chi) \leq 0$, $\chi \in [0, 1]$. This shows that $u(\chi)$ is decreasing on $[0, 1]$. Considering

$$\begin{aligned} u(1) &= \left(\frac{1}{1-\gamma} - 1 \right) \left(\frac{1}{2\lambda\Gamma(\zeta-3)} \int_0^\eta (\eta-\xi)^{\zeta-4}y(\xi)d\xi - \frac{1}{\Gamma(\zeta)} \int_0^1 (1-\xi)^{\zeta-1}y(\xi)d\xi \right) \\ &\geq \left(\frac{1}{1-\gamma} - 1 \right) \left(\frac{\eta}{2\lambda\Gamma(\zeta-3)} \int_0^1 y(\eta\xi)d\xi - \frac{1}{\Gamma(\zeta)} \int_0^1 y(\xi)d\xi \right) \\ &\geq \left(\frac{1}{1-\gamma} - 1 \right) \left(\frac{(\zeta-2)(\zeta-1)}{2\gamma\eta\Gamma(\zeta)} \int_0^1 y(\eta\xi)d\xi - \frac{1}{\Gamma(\zeta)} \int_0^1 y(\xi)d\xi \right) \geq 0, \end{aligned}$$

we get $u(\chi) \geq 0$, $\chi \in [0, 1]$. Hence $u \in K_0$.

Lemma 2.8 Let (2.10) hold and $y \in K_0$. Then u given by (2.5) satisfies

$$\min_{\chi \in [0, \theta]} u(\chi) \geq \theta^* \|u\|,$$

where $\theta \in (0, \eta)$, $\theta^* = \frac{\eta - \theta}{\eta}$, $\|u\| = \max_{\chi \in [0, 1]} |u(\chi)|$.

Proof. By Lemma 2.7, we have

$$u(\chi) \geq \frac{\eta - \chi}{\eta} u(0) + \frac{\chi}{\eta} u(\eta) \geq \frac{\eta - \chi}{\eta} u(0) = \frac{\eta - \chi}{\eta} \|u\|, \quad \chi \in [0, \eta].$$

Consequently, for $\theta \in (0, \eta)$, it gets

$$\min_{\chi \in [0, \theta]} u(\chi) = u(\theta) \geq \frac{\eta - \theta}{\eta} \|u\| = \theta^* \|u\|.$$

Lemma 2.9 [9] (Guo-Krasnoselskii fixed point theorem) Let \mathbb{E} be a Banach space and $K \subseteq \mathbb{E}$ be a cone. Suppose $\Omega_1 \subseteq \mathbb{E}$ and $\Omega_2 \subseteq \mathbb{E}$ are bounded open sets such that $0 \in \Omega_1$, $\bar{\Omega}_1 \subset \Omega_2$, and let $T : K \cap (\bar{\Omega}_2 \setminus \Omega_1) \rightarrow K$ be a completely continuous operator, and T satisfies either

- (1) $\|Tu\| \leq \|u\|$ for $u \in K \cap \partial\Omega_1$ and $\|Tu\| \geq \|u\|$ for $u \in K \cap \partial\Omega_2$, or
- (2) $\|Tu\| \geq \|u\|$ for $u \in K \cap \partial\Omega_1$ and $\|Tu\| \leq \|u\|$ for $u \in K \cap \partial\Omega_2$.

Then T has a fixed point in $K \cap (\bar{\Omega}_2 \setminus \Omega_1)$.

3. Main results

Let $0 < \theta \leq \eta - \frac{2\gamma(\zeta-3)}{\zeta(\zeta-1)(\zeta-2)}\eta^2 < \eta$ and

$$K = \{u \in K_0 : \min_{\chi \in [0, \theta]} u(\chi) \geq \theta^* \|u\|\}.$$

Lemma 3.1 Let (2.10) hold. Define $T : K \rightarrow C[0, 1]$ by

$$Tu(\chi) = \int_0^1 G(\chi, \xi) f(\xi, u(\xi)) d\xi, \quad \chi \in [0, 1].$$

Assume that f satisfies

(H) $f(\chi, u)$ is decreasing with respect to χ , and increasing with respect to u .

Then $T : K \rightarrow K$, and T is completely continuous.

Proof. As $u \in K$ and (H) holds, we have $f(\cdot, u(\cdot)) \in K_0$. Thus, we determine that $T : K \rightarrow K$ via Lemmas 2.7 and 2.8.

Next, we will use the Arzela-Ascoli theorem to testify the consequence. Here, (a) Let $u_n \rightarrow u$ in K . Then

$$\begin{aligned} |Tu_n(\chi) - Tu(\chi)| &\leq \frac{1}{\Gamma(\zeta)} \int_0^\chi (\chi - \xi)^{\zeta-1} |f(\xi, u_n(\xi)) - f(\xi, u(\xi))| d\xi \\ &\quad + \frac{1}{(1-\gamma)\Gamma(\zeta)} \int_0^1 (1 - \xi)^{\zeta-1} |f(\xi, u_n(\xi)) - f(\xi, u(\xi))| d\xi \\ &\quad + \frac{1-\gamma-\chi^2}{2\lambda\Gamma(\zeta-3)} \int_0^\eta (\eta - \xi)^{\zeta-4} |f(\xi, u_n(\xi)) - f(\xi, u(\xi))| d\xi. \end{aligned}$$

Note that f is continuous, and thus, we obtain

$$\|Tu_n - Tu\| \rightarrow 0, \quad n \rightarrow \infty.$$

Thus, T is continuous.

(b) Let $\Omega \subset K$ be bounded. Then, for any $u \in \Omega$, we have

$$\begin{aligned} |Tu(\chi)| &\leq \frac{L}{\Gamma(\zeta)} \int_0^\chi (\chi - \xi)^{\zeta-1} d\xi + \frac{L}{(1-\gamma)\Gamma(\zeta)} \int_0^1 (1 - \xi)^{\zeta-1} d\xi + \frac{\frac{1}{1-\gamma}L}{2\lambda\Gamma(\zeta-3)} \int_0^\eta (\eta - \xi)^{\zeta-4} d\xi \\ &\leq (1 + \frac{1}{1-\gamma}) \frac{L}{\Gamma(\zeta)} + \frac{\frac{1}{1-\gamma}L}{2\lambda\Gamma(\zeta-2)}, \end{aligned}$$

where $L = \max_{0 \leq \chi \leq 1, 0 \leq u \leq M} |f(\chi, u)| + 1$. Hence, T is uniformly bounded.

(c) For any $u \in \Omega$, $\chi_1, \chi_2 \in [0, 1]$, $\chi_1 < \chi_2$, we get

$$\begin{aligned} |Tu(\chi_2) - Tu(\chi_1)| &\leq \left| \frac{1}{\Gamma(\zeta)} \int_0^{\chi_2} (\chi_2 - \xi)^{\zeta-1} f(\xi, u(\xi)) d\xi - \frac{1}{\Gamma(\zeta)} \int_0^{\chi_1} (\chi_1 - \xi)^{\zeta-1} f(\xi, u(\xi)) d\xi \right| \\ &\quad + \frac{\chi_2^2 - \chi_1^2}{2\lambda\Gamma(\zeta-3)} \int_0^\eta (\eta - \xi)^{\zeta-4} f(\xi, u(\xi)) d\xi \\ &\leq \frac{L}{\Gamma(\zeta+1)} (\chi_2^\zeta - \chi_1^\zeta) + \frac{(\chi_2^2 - \chi_1^2)L}{2\lambda\Gamma(\zeta-2)}, \end{aligned} \quad (3.1)$$

which implies that the left-hand side of (3.1) $\rightarrow 0$ if $\chi_1 \rightarrow \chi_2$. Thus, T is equicontinuous in K .

Combining the above three steps (a), (b), (c) with the Arzela-Ascoli theorem, we discern that T is completely continuous.

Lemma 3.2 Let (2.10) and (H) hold. Suppose there is a number $r_1 > 0$ such that

$$f(0, r_1) \leq \frac{r_1}{A},$$

where $A = \frac{1}{(1-\gamma)\lambda\Gamma(\zeta-2)} > 0$ is a constant. Then we have

$$\|Tu\| \leq \|u\|, \quad u \in K \cap \partial\Omega_{r_1}, \quad (3.2)$$

with $\Omega_{r_1} = \{u \in C[0, 1] : \|u\| < r_1\}$.

Proof. From Remark 2.6, it gets

$$0 < G(\chi, \xi) = -\frac{(1-\xi)^{\zeta-1}}{(1-\gamma)\Gamma(\zeta)} + g_2(\chi, \xi) + \frac{(\frac{1}{1-\gamma} - \chi^2)(\eta - \xi)^{\zeta-4}}{2\lambda\Gamma(\zeta-3)}, \quad 0 \leq \xi < \eta,$$

and $G(\chi, \xi) \leq 0$, $\eta \leq \xi \leq 1$. As $3 < \zeta \leq 4$, $0 < \eta, \gamma < 1$ and (2.10) holds, for $0 \leq \xi < \eta$, we can deduce

$$\begin{aligned} G(\chi, \xi) &\leq \frac{(\chi - \xi)^{\zeta-1}}{\Gamma(\zeta)} + \frac{(\frac{1}{1-\gamma} - \chi^2)(\eta - \xi)^{\zeta-4}}{2\lambda\Gamma(\zeta-3)} \\ &\leq \frac{2\gamma\eta^2}{(\zeta-1)(\zeta-2)} \cdot \frac{\chi^{\zeta-1}(1 - \frac{\xi}{\chi})^{\zeta-1}}{2\lambda\Gamma(\zeta-3)} + \frac{(\frac{1}{1-\gamma} - \chi^2)(\eta - \xi)^{\zeta-4}}{2\lambda\Gamma(\zeta-3)} \\ &\leq \frac{2}{(\zeta-1)(\zeta-2)} \cdot \frac{\chi^2(\eta - \xi)^{\zeta-4}}{2\lambda\Gamma(\zeta-3)} + \frac{(\frac{1}{1-\gamma} - \chi^2)(\eta - \xi)^{\zeta-4}}{2\lambda\Gamma(\zeta-3)} \\ &\leq \frac{\frac{1}{1-\gamma}(\eta - \xi)^{\zeta-4}}{2\lambda\Gamma(\zeta-3)} = g_3(0, \xi), \quad \xi \leq \chi, \end{aligned} \quad (3.3)$$

and

$$G(\chi, \xi) \leq \frac{(\frac{1}{1-\gamma} - \chi^2)(\eta - \xi)^{\zeta-4}}{2\lambda\Gamma(\zeta-3)} \leq \frac{\frac{1}{1-\gamma}(\eta - \xi)^{\zeta-4}}{2\lambda\Gamma(\zeta-3)} = g_3(0, \xi), \quad \chi \leq \xi. \quad (3.4)$$

Thus, for $u \in K \cap \partial\Omega_{r_1}$, we have

$$\begin{aligned} 0 \leq Tu(\chi) &= \int_0^1 G(\chi, \xi) f(\xi, u(\xi)) d\xi \\ &= \int_0^\eta G(\chi, \xi) f(\xi, u(\xi)) d\xi + \int_\eta^1 G(\chi, \xi) f(\xi, u(\xi)) d\xi \\ &\leq \int_0^\eta G(\chi, \xi) f(\xi, u(\xi)) d\xi \\ &\leq f(0, u(0)) \cdot \int_0^\eta g_3(0, \xi) d\xi \\ &\leq f(0, r_1) \frac{\frac{1}{1-\gamma}}{2\lambda\Gamma(\zeta-2)} \leq f(0, r_1) A \leq r_1 = \|u\|, \quad \chi \in [0, 1]. \end{aligned}$$

This shows that (3.2) holds.

Lemma 3.3 *Let (2.10) and (H) hold. Suppose there is a number $r_2 > 0$ such that*

$$f(\theta, \theta^* r_2) \geq \frac{r_2}{B}.$$

where $B = \int_0^\theta G(0, \xi) d\xi$. Then we have

$$\|Tu\| \geq \|u\|, \quad u \in K \cap \partial\Omega_{r_2}, \quad (3.5)$$

with $\Omega_{r_2} = \{u \in C[0, 1] : \|u\| < r_2\}$.

Proof. First, we will prove $0 < B < A$. For $\xi < \theta < \eta$, it gets $G(0, \xi) > 0$ via Remark 2.6. Hence, $B > 0$. As (3.3) and (3.4) hold, we have

$$B = \int_0^\theta G(0, \xi) d\xi \leq \int_0^\theta g_3(0, \xi) d\xi < \frac{1}{2\lambda(1-\gamma)\Gamma(\zeta-2)} \leq A.$$

Thus, B is a constant, and satisfies $0 < B < A$. Next, we shall show that $\int_\theta^1 G(0, \xi) f(\xi, u(\xi)) d\xi \geq 0$. Using the condition (H) and $0 < \theta \leq \eta - \frac{2\gamma(\zeta-3)}{\zeta(\zeta-1)(\zeta-2)}\eta^2$, we have

$$\begin{aligned} \int_\theta^1 G(0, \xi) f(\xi, u(\xi)) d\xi &= \int_\theta^\eta G(0, \xi) f(\xi, u(\xi)) d\xi + \int_\eta^1 G(0, \xi) f(\xi, u(\xi)) d\xi \\ &\geq f(\eta, u(\eta)) \left(\int_\theta^\eta G(0, \xi) d\xi + \int_\eta^1 G(0, \xi) d\xi \right) \\ &= f(\eta, u(\eta)) \left(\int_\theta^\eta \frac{\frac{1}{1-\gamma}(\eta-\xi)^{\zeta-4}}{2\lambda\Gamma(\zeta-3)} d\xi - \int_\theta^1 \frac{1}{(1-\gamma)\Gamma(\zeta)} (1-\xi)^{\zeta-1} d\xi \right) \\ &= f(\eta, u(\eta)) \left(\frac{(\eta-\theta)^{\zeta-3}}{2(1-\gamma)\lambda\Gamma(\zeta-2)} - \frac{(1-\theta)^\zeta}{(1-\gamma)\Gamma(\zeta+1)} \right) \\ &\geq \frac{f(\eta, u(\eta))}{(1-\gamma)\Gamma(\zeta+1)} \left(\frac{\zeta(\zeta-1)(\zeta-2)(\eta-\theta)^{\zeta-3}}{2\gamma\eta^2(\zeta-3)} - 1 \right) \geq 0. \end{aligned}$$

Now, it remains to verify (3.5). For $u \in K \cap \partial\Omega_{r_2}$, it gets

$$\begin{aligned}
 \|Tu\| = Tu(0) &= \int_0^1 G(0, \xi) f(\xi, u(\xi)) d\xi \\
 &= \int_0^\theta G(0, \xi) f(\xi, u(\xi)) d\xi + \int_\theta^1 G(0, \xi) f(\xi, u(\xi)) d\xi \\
 &\geq \int_0^\theta G(0, \xi) f(\xi, u(\xi)) d\xi \\
 &\geq f(\theta, u(\theta)) \int_0^\theta G(0, \xi) d\xi \\
 &\geq f(\theta, \theta^* r_2) B \geq r_2 = \|u\|.
 \end{aligned}$$

Thus, (3.5) holds.

Theorem 3.4 Let (2.10) and (H) hold. Suppose that: there are two numbers $r_1 > 0$ and $r_2 > 0$ with $r_1 \neq r_2$, and r_1, r_2 satisfy

$$f(0, r_1) \leq \frac{r_1}{A}, \quad f(\theta, \theta^* r_2) \geq \frac{r_2}{B}.$$

Then the BVP (1.1)–(1.2) has at least one positive and decreasing solution u , and

$$r_1 \leq \|u\| \leq r_2 \quad (\text{or } r_2 \leq \|u\| \leq r_1).$$

Furthermore, $u(\chi)$ is concave on $[0, \eta]$.

Proof. Let $r_1 < r_2$. In view of Lemmas 3.2 and 3.3, we get

$$\|Tu\| \leq \|u\|, \quad u \in K \cap \partial\Omega_{r_1},$$

$$\|Tu\| \geq \|u\|, \quad u \in K \cap \partial\Omega_{r_2}.$$

Hence, it follows the Guo-Krasnoselskii fixed point theorem that T has a fixed point $u \in K \cap (\overline{\Omega}_{r_2} \setminus \Omega_{r_1})$, which is a desired solution of the BVP (1.1)–(1.2).

Theorem 3.5 Let (2.10) and (H) hold. Assume that there exist three positive numbers r_1, r_2, r_3 with $r_1 < r_2 < r_3$, and meet one of the following conditions

$$f(0, r_1) \leq \frac{r_1}{A}, \quad f(\theta, \theta^* r_2) > \frac{r_2}{B}, \quad f(0, r_3) \leq \frac{r_3}{A}, \quad (3.6)$$

$$f(\theta, \theta^* r_1) \geq \frac{r_1}{B}, \quad f(0, r_2) < \frac{r_2}{A}, \quad f(\theta, \theta^* r_3) \geq \frac{r_3}{B}. \quad (3.7)$$

Then the BVP (1.1)–(1.2) has at least two positive and decreasing solutions u_1 and u_2 , and

$$r_1 \leq \|u_1\| < r_2 < \|u_2\| \leq r_3.$$

Proof. We only consider the case when (3.6) is satisfied, as the proof for the case (3.7) is similar. By Lemmas 3.2 and 3.3, we get

$$\|Tu\| \leq \|u\|, \quad u \in K \cap \partial\Omega_{r_1},$$

$$\|Tu\| > \|u\|, \quad u \in K \cap \partial\Omega_{r_2},$$

and

$$\|Tu\| \leq \|u\|, \quad u \in K \cap \partial\Omega_{r_3}.$$

Hence, it follows Guo-Krasnoselskii fixed point theorem that T has two fixed points $u_1 \in K \cap (\bar{\Omega}_{r_2} \setminus \Omega_{r_1})$ and $u_2 \in K \cap (\bar{\Omega}_{r_3} \setminus \Omega_{r_2})$, which are two desired solutions of the BVP (1.1)–(1.2).

Remark 3.6 *Similar to Theorem 3.5, for any positive integer $n(n \geq 2)$, we can obtain the existence of at least $n - 1$ positive and decreasing solutions of the BVP (1.1)–(1.2), where n is the number of $r_i, i = 1, 2, \dots, n$.*

4. Examples

Example 4.1 Consider

$${}^C D_{0+}^{3.5} u(\chi) = \frac{u(\chi)}{625} + 1 - \chi, \quad \chi \in [0, 1], \quad (4.1)$$

$$u'(0) = u'''(0) = 0, \quad u'''(0.5) + \frac{1}{32} u''(0) = 0, \quad u(1) - 0.5u(0) = 0, \quad (4.2)$$

where $\zeta = 3.5$, $\eta = 0.5$, $\lambda = \frac{1}{32}$, $\gamma = 0.5$, $f(\chi, u) = \frac{u}{625} + 1 - \chi$. Then (2.10) and (H) are satisfied.

The Green's function $G(\chi, \xi) = g_1(\chi, \xi) + g_2(\chi, \xi) + g_3(\chi, \xi)$, $(\chi, \xi) \in [0, 1] \times [0, 1]$, where

$$g_1(\chi, \xi) = -\frac{2}{\Gamma(3.5)}(1 - \xi)^{2.5}, \quad (\chi, \xi) \in [0, 1] \times [0, 1],$$

$$g_2(\chi, \xi) = \begin{cases} 0, & 0 \leq \chi \leq \xi \leq 1, \\ \frac{(\chi - \xi)^{2.5}}{\Gamma(3.5)}, & 0 \leq \xi \leq \chi \leq 1, \end{cases} \quad g_3(\chi, \xi) = \begin{cases} 0, & \xi \geq 0.5, \\ \frac{16(2 - \chi^2)(0.5 - \xi)^{-0.5}}{\Gamma(0.5)}, & \xi < 0.5. \end{cases}$$

This shows

$$\begin{aligned} G(\chi, \xi) &= -\frac{2}{\Gamma(3.5)}(1 - \xi)^{2.5} + g_2(\chi, \xi) + \frac{16(2 - \chi^2)(0.5 - \xi)^{-0.5}}{\Gamma(0.5)} \\ &\geq -\frac{2}{\Gamma(3.5)}(1 - \xi)^{2.5} + \frac{16(2 - \chi^2)(0.5 - \xi)^{-0.5}}{\Gamma(0.5)} \\ &\geq -\frac{2}{\Gamma(3.5)} + \frac{16}{\Gamma(0.5)} = 8.4252 > 0, \quad 0 \leq \xi < 0.5, \end{aligned}$$

and

$$\begin{aligned} G(\chi, \xi) &= -\frac{2}{\Gamma(3.5)}(1 - \xi)^{2.5} + g_2(\chi, \xi) + g_3(\chi, \xi) \\ &\leq -\frac{2}{\Gamma(3.5)}(1 - \xi)^{2.5} + \frac{(1 - \xi)^{2.5}}{\Gamma(3.5)} = -\frac{1}{\Gamma(3.5)}(1 - \xi)^{2.5} \leq 0, \quad 0.5 \leq \xi \leq 1. \end{aligned}$$

Thus, Remark 2.6 holds.

Next, we choose $\theta = 0.4$. Through computing, we have

$$\theta^* = \frac{\eta - \theta}{\eta} = \frac{1}{5}, \quad A = \frac{1}{(1 - \gamma)\lambda\Gamma(\zeta - 2)} = \frac{64}{\Gamma(1.5)} = 72.2163,$$

$$B = \int_0^\theta G(0, \xi) d\xi = \int_0^{0.4} \left(-\frac{2}{\Gamma(3.5)}(1-\xi)^{2.5} + \frac{32(\eta-\xi)^{-0.5}}{\Gamma(0.5)} \right) d\xi = 13.9707.$$

Moreover, it is easy to achieve

$$f(0, r_1) = 1 + \frac{r_1}{625} \leq \frac{r_1}{A}, \quad \text{if } r_1 \geq 81.6507,$$

$$f(0.4, 0.2 * r_2) = 0.6 + \frac{r_2}{3125} \geq \frac{r_2}{B}, \quad \text{if } r_2 \leq 8.4201.$$

Thus, we can choose $r_1 = 82$, $r_2 = 8$ in Theorem 3.4. Then the conditions of Theorem 3.4 are satisfied. Consequently the BVP (4.1)–(4.2) has a positive and decreasing solution u satisfying

$$8 \leq \|u\| \leq 82.$$

Example 4.2 Consider

$${}^C D_{0+}^{3.5} u(\chi) = \frac{u^2(\chi)}{20861} + 1 - \chi, \quad \chi \in [0, 1], \quad (4.3)$$

$$u'(0) = u'''(0) = 0, \quad u'''(0.5) + \frac{1}{32} u''(0) = 0, \quad u(1) - 0.5u(0) = 0, \quad (4.4)$$

where $\zeta = 3.5$, $\eta = 0.5$, $\lambda = \frac{1}{32}$, $\gamma = 0.5$, $f(\chi, u) = \frac{u^2}{20861} + 1 - \chi$. Then (2.10) and (H) are satisfied, and Remark 2.6 holds.

From Example 4.1, we get $\theta = 0.4$ and $\theta^* = \frac{1}{5}$, $A = 72.2163$, $B = 13.9707$. By computing, we have

$$f(0.4, 0.2 * r_1) = 0.6 + \frac{0.04 * r_1^2}{20861} \geq \frac{r_1}{B}, \quad \text{if } r_1 \geq 37322 \text{ or } r_1 \leq 8.3843,$$

$$f(0, r_2) = 1 + \frac{r_2^2}{20861} < \frac{r_2}{A}, \quad \text{if } 143.9608 < r_2 < 144.9075.$$

Thus, we can choose $r_1 = 8$, $r_2 = 144$, $r_3 = 37323$ in Theorem 3.5. Then the conditions of Theorem 3.5 are satisfied. Consequently the BVP (4.3)–(4.4) has two positive and decreasing solution u_1, u_2 , and

$$8 \leq \|u_1\| < 144 < \|u_2\| \leq 37323.$$

5. Conclusion

In the paper, for any positive integer $n(n \geq 2)$, the existence of at least $n - 1$ positive and decreasing solutions for Caputo three-point BVPs are studied. The obtained solutions are also proved to be concave on $[0, \eta]$. The main novelty of the paper lies in obtaining positive solutions while the corresponding Green's function changes sign. For future work, one can discuss the positive solutions for other types of FDEs, and can also consider much more difficult research on fractional differential systems.

Acknowledgment

This research was funded by National Natural Science Foundation of China (12201199, 61703180), the Natural Science Foundation of the Department of Education of Hunan Province (2022JJ40021), the Educational Department of Hunan Province of China (21B0722, 21C0660), and Science and Technology Program of University of Jinan (1008399).

References

- [1] Bai Z, Lü H. Positive solutions for boundary value problem of nonlinear fractional differential equation. *Journal of Mathematical Analysis and Applications* 2005; 311 (2): 495-505. <https://doi.org/10.1016/j.jmaa.2005.02.052>
- [2] Benhadri M, Caraballo T. On positive periodic solutions to third-order integro-differential equations with distributed delays. *Turkish Journal of Mathematics* 2023; 47 (6): 1822-1847. <https://doi.org/10.55730/1300-0098.3465>
- [3] Ding Y, Wei Z, Zhao Q. Solutions for a nonlinear fractional boundary value problem with sign-changing Green's function. *Journal of Nonlinear Sciences and Applications* 2015; 8 (5): 650-659. <https://doi.org/10.22436/JNSA.008.05.17>
- [4] Elsanosi MEM. Positive solutions of nonlinear neumann boundary value problems with sign-changing green's function. *Kyungpook Mathematical Journal* 2019; 59 (1): 65-71. <https://doi.org/10.5666/KMJ.2019.59.1.65>
- [5] Feng L, Sun Y, Han Z. Philos-type oscillation criteria for impulsive fractional differential equations. *Journal of Applied Mathematics and Computing* 2020; 62 (1-2): 361-376. <https://doi.org/10.1007/s12190-019-01288-5>
- [6] Ismail M, Saeed U, Alzabut J, Rehman M. Approximate solutions for fractional boundary value problems via Green-CAS wavelet method. *Mathematics* 2019; 7(12): 1164. <https://doi.org/10.3390/math7121164>
- [7] Khaminsou B, Thaiprayoon C, Alzabut J, Sudsutad W. Nonlocal boundary value problems for integro-differential Langevin equation via the generalized Caputo proportional fractional derivative. *Boundary Value Problems* 2020; 2020 (176): 1-30. <https://doi.org/10.1186/s13661-020-01473-7>
- [8] Kilbas AA, Srivastava HM, Trujillo JJ. *Theory and Applications of Fractional Differential Equations*. Amsterdam, The Netherlands: Elsevier B. V., 2006. [https://doi.org/10.1016/S0304-0208\(06\)80001-0](https://doi.org/10.1016/S0304-0208(06)80001-0)
- [9] Krasnosel'skii MA. *Positive Solutions of Operator Equations*. Groningen, The Netherlands: Noordho, 1966.
- [10] Liang H, Styne M. Collocation methods for general Caputo two-point boundary value problems. *Journal of Scientific Computing* 2018; 76 (1): 390-425. <https://doi.org/10.1007/s10915-017-0622-5>
- [11] Li H, Gao C, Dimitrov ND. Existence of positive solutions of discrete third-order three-point BVP with sign-changing Green's function. *Open Mathematics* 2022; 20 (1): 1229-1245. <https://doi.org/10.1515/math-2022-0510>
- [12] Li S, Zhang Z, Jiang W. Multiple positive solutions for four-point boundary value problem of fractional delay differential equations with p -Laplacian operator. *Applied Numerical Mathematics* 2021; 165: 348-356. <https://doi.org/10.1016/j.apnum.2021.03.001>
- [13] Liu T. Parameter estimation with the multigrid-homotopy method for a nonlinear diffusion equation. *Journal of Computational and Applied Mathematics* 2022; 413 (114393): 1-14. <https://doi.org/10.1016/j.cam.2022.114393>
- [14] Ma R. Nonlinear periodic boundary value problems with sign-changing Green's function. *Nonlinear Analysis: Theory, Methods & Applications* 2011; 74 (5): 1714-1720. <https://doi.org/10.1016/j.na.2010.10.043>
- [15] Selvam AGM, Alzabut J, Dhineshabu R, Rashid S, Rehman M. Discrete fractional order two-point boundary value problem with some relevant physical applications. *Journal of Inequalities and Applications* 2020; 2020 (221): 1-19. <https://doi.org/10.1186/s13660-020-02485-8>
- [16] Sun J, Zhao J. Iterative technique for a third-order three-point BVP with sign-changing green's function. *Electronic Journal of Differential Equations* 2013; 2013 (215): 1-9.
- [17] Telli B, Souid MS, Alzabut J, Khan H. Existence and uniqueness theorems for a variable-order fractional differential equation with delay. *Axioms* 2023; 12 (4): 339. <https://doi.org/10.3390/axioms12040339>
- [18] Wang Y, Wang H. Triple positive solutions for fractional differential equation boundary value problems at resonance. *Applied Mathematics Letters* 2020; 106 (106376): 1-7. <https://doi.org/10.1016/j.aml.2020.106376>
- [19] Xu M, Sun S. Positivity for integral boundary value problems of fractional differential equations with two nonlinear terms. *Journal of Applied Mathematics and Computing* 2019; 59 (1-2): 271-283. <https://doi.org/10.1007/s12190-018-1179-7>

- [20] Yan R, Ma Q, Ding X. Existence results for coupled nonlinear fractional differential equations with coupled strip and infinite point boundary conditions. *Fixed Point Theory* 2021; 22 (2): 913-932. <https://doi.org/10.24193/fpt-ro.2021.2.60>
- [21] Yan R, Zhao Y. Positive solutions for periodic boundary value problems of fractional differential equations with sign-changing nonlinearity and Green's function. *Axioms* 2023; 12 (9): 819. <https://doi.org/10.3390/axioms12090819>
- [22] Yue Y, Tian Y, Bai Z. Infinitely many nonnegative solutions for a fractional differential inclusion with oscillatory potential. *Applied Mathematics Letters* 2019; 88: 64-72. <https://doi.org/10.1016/j.aml.2018.08.010>
- [23] Zhang C, Sun S. Sturm-Picone comparison theorem of a kind of conformable fractional differential equations on time scales. *Journal of Applied Mathematics and Computing* 2017; 55 (1-2): 191-203. <https://doi.org/10.1007/s12190-016-1032-9>
- [24] Zhang W, Ni J. New multiple positive solutions for Hadamard-type fractional differential equations with nonlocal conditions on an infinite interval. *Applied Mathematics Letters* 2021; 118 (107165): 1-10. <https://doi.org/10.1016/j.aml.2021.107165>
- [25] Zhang Y, Sun J, Zhao J. Positive solutions for a fourth-order three-point BVP with sign-changing Green's function. *Electronic Journal of Qualitative Theory of Differential Equations* 2018; 2018 (5): 1-11. <https://doi.org/10.14232/ejqtde.2018.1.5>
- [26] Zhao Y. On the existence for a class of periodic boundary value problems of nonlinear fractional hybrid differential equations. *Applied Mathematics Letters* 2021; 121 (107368): 1-6. <https://doi.org/10.1016/j.aml.2021.107368>
- [27] Zhao Y, Sun Y. On the existence for parametric boundary value problems of a coupled system of nonlinear fractional hybrid differential equations. *Turkish Journal of Mathematics* 2022; 46 (2): 612-625. <https://doi.org/10.3906/mat-2107-19>